



# Internet Addiction among Egyptian Teens during COVID-19 Pandemic

Ayman F. Armaneous<sup>1\*</sup>, Hanan Atta<sup>2</sup>, Alshaimaa A. Elkhatib<sup>2</sup>, Alshaimaa F. Kamal<sup>2</sup>, Sara F. Sallam<sup>2</sup>

<sup>1</sup>Department of Child Health, National Research Center, Institute of Medical Research and Clinical studies, Cairo, Egypt;

<sup>2</sup>Department of Child Health, National Research Centre, Cairo, Egypt

## Abstract

**Edited by:** Ksenija Bogoeva-Kostovska  
**Citation:** Armaneous AF, Atta H, Elkhatib AA, Kamal AF, Sallam SF. Internet Addiction among Egyptian Teens during COVID-19 Pandemic. Open Access Maced J Med Sci. 2024 Sep 15; 12(3):388-397. <https://doi.org/10.3889/oamjms.2024.11901>  
**Keywords:** COVID-19 infection; Internet addiction; Teens  
**\*Correspondence:** Ayman F. Armaneous, Department of Child Health, National Research Center, Institute of Medical Research and Clinical studies, Cairo, Egypt. E-mail: fekreyayman@yahoo.com  
**Received:** 24-Mar-2024  
**Revised:** 25-Apr-2024  
**Accepted:** 10-Jun-2024  
**Ahead of print:** 30-Jul-2024  
**Copyright:** © 2024 Ayman F. Armaneous, Hanan Atta, Alshaimaa A. Elkhatib, Alshaimaa F. Kamal, Sara F. Sallam  
**Funding:** This research did not receive any financial support  
**Competing Interests:** The authors have declared that no competing interests exist  
**Open Access:** This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

**BACKGROUND:** Teens' lifestyles have changed significantly during and after COVID-19 epidemic and they have started engaging in risky health behaviors like using smartphone more frequently. Everyone is compelled to use their smartphone more frequently than usual to access daily necessities during the COVID-19 pandemic lockdown.

**AIM:** We aimed to evaluate the problem of internet addiction and the increasing use of smartphone among teenage Egyptian youth during COVID-19 pandemic.

**METHODS:** Online surveys were sent to Egyptian teens (12–18 years old) in several governorates in June 2020, and they were completed by 154 youths over the course of one month. Different teenagers were required to respond to and share in this poll, and versions in both English and Arabic were used. At the outset of the questionnaire, the question about consent was viewed as mandatory. The extent of Internet use was evaluated using Young's Internet Addiction Test (IAT). The survey was divided into four sections and created with the goals of understanding the modes (frequency, patterns, and goals) of internet use, the negative effects, linked parental behaviors, as well as the eventual occurrence and severity of addiction. The same data from the time before the epidemic was also looked into and analyzed. Version 23 of IBM SPSS Statistics was used for the analysis. The p-value threshold for significance was established at 0.05.

**RESULTS:** The mean age of the teens was  $14.38 \pm 2.87$  years and equal gender frequency (males of 50.6% and females of 49.4%). About 79.9% of them live in urban environment, 18.8% suffered from family problems and 40.9% were in the secondary educational level. Increased physical inactivity of participants was noticed after COVID-19 among participants (55.8%). About 87.7% of them used smartphone for online access (87.7%). Half of the subjects (50%) spent 1–3 h online before COVID, while more than half of them (59.1%) spent more than 6 hours online after COVID, with a statistically significant difference. Mean IAT was  $61.91 \pm 16.77$  after COVID versus  $45.26 \pm 12.45$  before COVID. The prevalence of internet addiction was increased from 67.5% before COVID-19 to 77.9% after COVID-19 infection. IAT score was normal, mild, moderate, and severe in 5.8%, 16.2%, 65.6%, and 12.3% after COVID-19, respectively, compared to normal, mild, moderate, and severe IAT score in 26.6%, 5.8%, 61.0%, and 6.5% before COVID, respectively, with significant differences between them before and after COVID ( $p < 0.000$ ). Increasing mother's and father's educational, increasing time spent online, decreasing duration of sleep, and lower the number of days of exercise for more than 30 min/week were associated with higher IAT score.

**CONCLUSIONS:** During the COVID-19 pandemic, internet addiction was extremely common among teenagers (77.9%). The amount of time spent online, the length of sleep, and the number of days that a person exercises for more than 30 min/week were all predictors of addiction.

## Introduction

The adoption of numerous measures, including both pharmacological and non-pharmacological methods, to reduce the extensive coronavirus-19 transmission in the world resulted from an improved understanding of the epidemiology of the pandemic [1]. Following that, many nations implemented countrywide lockdowns by closing all schools and offices [2], [3], [4]. Egypt went under lockdown on March 14, 2020 [5]. Lockdown made teenagers spend more time at home, which increased the amount of time, spent online [6].

Egypt's internet user base increased from 44.94 million in 2019 to 54.74 million in January 2020 (+22%). There were 80.75 million additional online users

at the start of 2023 when the percentage of people using the internet was 72.2%. In early 2023, there were 46.25 million (41.4%) social media users and 105.1 million (93.9%) active mobile phone connections in Egypt [7]. The coronavirus-19 disease (COVID-19) outbreak's after effects have exacerbated the link between technology and addiction. The COVID-19 problem has impacted health and well-being, which has exacerbated smartphone addiction [8], [9], [10], [11], [12], [13].

The use of smartphones has significantly altered daily routines and behavior. Applications enable social interaction, email access, music/video/film enjoyment, gameplay, and scheduling management. According to Hong *et al.* [14], smartphones can broaden horizons, foster safety, reduce stress, preserve connections, and provide essential information. However, improper

smartphone use encourages unintentional time-wasting, and excessive use raises the possibility of smartphone addiction, which can have negative effects on one's physical and mental well-being and lead to dependence. It can also have an impact on routines, habits, social behaviors, family relationships, and interpersonal relationships [15]. Relevant studies have examined whether internet addiction has significantly increased during the COVID-19 home quarantines, particularly among teenagers, without establishing a clear accurate prevalence [9], [10], [11], [16], [17].

### ***Aim of the study***

We aimed to evaluate the problem of internet addiction and the increasing use of smartphone among teenage Egyptian youth during the COVID-19 pandemic.

## **Methods**

### ***Study design***

Observational, cross-sectional online study design was used in this study.

### ***Study participants***

The study constituted adolescents attending secondary schools in different governorates in Egypt.

### ***Eligibility criteria***

(a) Teens (12–18 years), (b) Having an internet account or smartphone cell, and (c) voluntary participation in the survey.

### ***Exclusion criteria***

Subjects younger than 12 years of age, all adolescents who refused to consent, and those whose parents refused to consent were eliminated from the study. In addition, those who were unable to engage in the study due to illness (both physical and mental) were not included.

### ***Sampling methods***

#### ***Determination of sample size***

Using Epi Info 7 software (center for disease control and prevention) Atlanta, Georgia) program for calculation of sample size and based on the expected prevalence of internet addiction of 53.6% [18], 95% confidence level, and 5% confidence limits, the minimum sample size required is 150.

### ***Sampling technique***

Online self-report and anonymous questionnaire was sent to 170 Egyptian teens (12–18 years old) in several governorates (Cairo, Giza, Alexandria, Elbehira, Qaliubia, Dakahlia, Kafr El-Sheikh, Suez, Ismailia, Fayoum, Beni-Suef, Menia, Sohag, Assiut, and Menoufia) over the course of one month (June 2020), through technological means (email, Twitter, Face book, and WhatsApp) along with a detailed explanation of the study's objectives. Different teenagers were required to respond to and share in this poll, and versions in both English and Arabic were used.

We had 154 participants overall after excluding 16 pupils from the study due to incomplete questionnaires. At the outset of the questionnaire, the question about consent was viewed as mandatory. The extent of internet use was evaluated using Young's Internet Addiction Test (IAT).

### ***Tools***

#### ***IAT***

This questionnaire consists of 20 items regarding internet overuse. All items begin with the phrase "How often do you...", for example, "How often do you try to cut down the amount of time you spend online and fail?" Respondents are requested to choose one of the following scores: 5 = always, 4 = often, 3 = frequently, 2 = occasionally, and 1 = rarely. The IAT has been used to measure the severity of internet addiction. The total score of IAT ranges from 20 to 100. In the present study, we classified the level of internet addiction according to the cutoff points previously reported by Young (1998) [19].

- 20–30 points are considered normal internet usage;
- 31–49 points indicate the presence of a mild internet addiction;
- 50–79 points reflect the presence of a **moderate** internet addiction;
- 80–100 points indicate a severe internet addiction.

### ***Ethical considerations***

The Institutional Review Board (IRB) of the National Research Centre in Giza, Egypt, gave its approval to this study. In addition, each participant's informed consent was gained by a statement of agreement at the start of each questionnaire. All procedures were also carried out in accordance with the Declaration of Helsinki and all applicable rules and regulations. Teenagers were emailed a link to an online survey form that was labeled with the stud's goal. The online questionnaire's completion and submission were taken into account as the parent's written approval of his child's involvement in the study.

### Statistical analysis

The statistical program IBM SPSS (Statistical Package for the Social Sciences) version 23 (IBM SPSS Inc., Chicago, IL, USA) was used to process the data. To control the findings, variables pertaining to the respondents' demographic traits, their family structure, and their parents' socioeconomic level were included in the analysis. Bivariate analysis (Chi-square) was used to identify associations between variables, and multivariate analysis (binary logistic regression model) was used to identify the predictors for all variables with a maximum p-value of 0.2. Using two-tailed tests, the level of statistical significance for each analysis was set at = 0.05.

### Outcome measures

- Primary outcome: Prevalence of internet addiction among the participants.
- Data measurement
- Secondary outcomes: Participants' socioeconomic and demographic characteristics that are linked to and predict internet addiction.

## Results

### Baseline and social characteristics of the respondents

The current study was done on 154 participants with a mean age ( $\pm$  SD) of students participating in this study was 14.38 ( $\pm$  2.87), males constituted about 50.6% of participants and 79.9% of them live in urban. Majority of students (68.8%) live in Cairo, Giza, and Alexandria, while 11.7% were living abroad. About 18.8% of participants suffered from family problems. Nearly equal number of students were in international school and language private schools (34.4% each), while 40.9% of participants were in the secondary educational level. The majority of participants' mothers and fathers were graduate (55.8% and 57.8%, respectively), with 76.6% of them live in medium socioeconomic level and the majority (92.2%) were living with two parents. 63.6% of students completed the form without any help from mothers, fathers, or other members. Majority of participants used smartphone for online access (87.7%), 42.2% of them had laptops and 17.5% of them used tablets (Table 1).

### Before and after COVID infection

Increased physical inactivity of participants was noticed after COVID-19 among participants (55.8%), versus 22.1% of inactivity before COVID-19 ( $p = 0.009$ ).

**Table 1: Baseline and social characteristics of enrolled students (n = 154)**

Variables	n (%)
Age (years)	
Mean $\pm$ SD	14.38 $\pm$ 2.87
Range	(12–18)
Gender	
Male	78 (50.6)
Female	76 (49.4)
Residence	
Urban	123 (79.9)
Abroad	31 (20.1)
Languages	
Arabic	105 (68.2)
English	49 (31.8)
Educational level	
Primary	31 (20.1)
Preparatory	40 (26.0)
Secondary	63 (40.9)
Diploma	2 (1.3)
University	18 (11.7)
Type of school	
International school	53 (34.4)
Language private school	53 (34.4)
Arabic private school	19 (12.3)
Language experimental school	10 (6.5)
Arabic experimental school	3 (1.9)
State school	16 (10.4)
Family status	
Normal	125 (81.2)
Family conflicts	11 (7.1)
Divorce	6 (3.9)
Child's negative feeling to parents	8 (5.2)
Parent's negative feeling to parent	8 (5.2)
Single child family	6 (3.9)
Mother's education	
Primary or preparatory	4 (2.6)
Secondary or diploma	17 (11.0)
Graduate	86 (55.8)
Postgraduate	47 (30.5)
Father's Education	
Primary or preparatory	1 (0.6)
Secondary or diploma	15 (9.7)
Graduate	89 (57.8)
Postgraduate	49 (31.8)
Socioeconomic level	
Low socioeconomic level	2 (1.3)
Medium socioeconomic level	118 (76.6)
High socioeconomic level	34 (22.1)
With whom does the student live?	
Two parents	142 (92.2)
Mother only	11 (7.1)
Father only	1 (0.6)
Who completed the form?	
Student	98 (63.6)
Student with mother's help	35 (22.7)
Student with father's help	11 (7.1)
Others	10 (6.5)
You go online using	
Smartphone	135 (87.7)
Laptop	65 (42.2)
Tablet	27 (17.5)
PC	15 (9.7)
Smartwatch	4 (2.6)

Regarding hours of sleep, 66.2% of participants had 6–8 h of sleep before COVID-19, while 53.2% of them had 8–12 h of sleep after COVID-19, with a statistically significant difference. Only 12.3% of participants spent more than 6 hours with their family before COVID-19, versus 39.6% after COVID-19. Half of the subjects (50%) spent 1–3 h online before COVID-19, while more than half of them (59.1%) spent more than 6 h online after COVID-19, with a statistically significant difference. As expected, real social activity had a statistically significant decrease after COVID-19. Most of the students did not use the online learning at all before COVID (57.1%), while more online learning was used at the time of COVID, with 50.6% of students used the internet access for doing homework (Table 2).

There was a highly significant increase in the time spent online for different purposes after COVID-19

**Table 2: Frequency distribution and percentage of number of hours student spend sleeping, with family, doing social activity or online**

Variables	Before COVID		After COVID		p-value
	n	%	n	%	
How many days/week do you exercise for more than 30 minutes?					
None	34	22.1	86	55.8	0.009*
1 day/week	15	9.7	19	12.3	
2 days/week	27	17.5	16	10.4	
3 days/week	41	26.6	23	14.9	
4 days/week	10	6.5	3	1.9	
5 days/week	27	17.5	7	4.5	
How many hours do you sleep per day?					
<4 h	0	0.0	1	0.6	0.000*
4–6 h	15	9.7	11	7.1	
6–8 h	102	66.2	53	34.4	
8–12 h	35	22.7	82	53.2	
More than 12 h	2	1.3	7	4.5	
How many hours you spend with your family per day?					
<1 h	38	24.7	16	10.4	0.000*
1–3 h	66	42.9	43	27.9	
3–6 h	31	20.1	34	22.1	
More than 6 h	19	12.3	61	39.6	
How many hours do you have real social activity?					
<1 h	35	22.7	100	64.9	0.025*
1–3 h	54	35.1	29	18.8	
3–6 h	30	19.5	17	11.0	
More than 6 h	35	22.7	8	5.2	
How many hours do you spend time online					
<1 h	19	12.3	6	3.9	0.000*
1–3 h	77	50.0	13	8.4	
3–6 h	39	25.3	44	28.6	
More than 6 h	19	12.3	91	59.1	
The online learning you used was in the form of					
None	88	57.1	17	11.0	0.000*
Interactive learning	22	14.3	58	37.3	
Homework	46	29.9	78	50.6	
Videos	22	14.3	77	50	
Others	4	2.6	9	5.8	

among students, such as chat, social networking services, gaming, and videos. Other purposes included food channels, TikTok, Instagram, courses, reading, children’s sites, training, religious sites, drawing, and watching series and movies (Table 3). There was a highly significant increase in the frequency of use of various social networking services after COVID-19 than before, especially, the Facebook, Zoom, and TikTok services. Other services used included Netflix, Viber, YouTube, Twitch, and Pinterest (Table 4).

**The prevalence of internet addiction among the respondents**

Total scoring of IAT was higher among students after COVID (61.91 ± 16.77) than before COVID-19 (45.26 ± 12.45), with a highly significant difference (p < 0.001). IAT score was normal, mild, moderate, and severe in 5.8%, 16.2%, 65.6%, and 12.3% after COVID-19, respectively, compared to normal, mild, moderate, and severe IAT score in 26.6%, 5.8%, 61.0%, and 6.5% before COVID-19, respectively, with significant differences between them before and after COVID-19 (p < 0.000). We consider moderate and severe addictions were the categories that significantly relevant, so the prevalence of internet addiction among enrolled teens was increased from 67.5% before COVID-19 to 77.9% of 154 teens after COVID-19 infection (Table 5).

**Table 3: Frequency distribution and percentage of the time spent online for different purposes**

	Before COVID		After COVID		P value*
	n	%	n	%	
Chat					
Never	10	6.5	3	1.9	0.000
Rarely	30	19.5	20	13.0	
Sometimes	51	33.1	27	17.5	
Often	37	24.0	49	31.8	
Always	26	16.9	55	35.7	
Social networking service					
Never	14	9.1	9	5.8	0.001*
Rarely	26	16.9	21	13.6	
Sometimes	50	32.5	25	16.2	
Often	30	19.5	38	24.7	
Always	34	22.1	61	39.6	
Gaming					
Never	15	9.7	14	9.1	0.000*
Rarely	41	26.6	32	20.8	
Sometimes	46	29.9	23	14.9	
Often	28	18.2	43	27.9	
Always	24	15.6	42	27.3	
Music					
Never	13	8.4	14	9.1	0.000*
Rarely	50	32.5	28	18.2	
Sometimes	32	20.8	42	27.3	
Often	28	18.2	29	18.8	
Always	31	20.1	41	26.6	
Video					
Never	3	1.9	4	2.6	0.000*
Rarely	28	18.2	16	10.4	
Sometimes	48	31.2	42	27.3	
Often	46	29.9	38	24.7	
Always	29	18.8	54	35.1	
Web searches					
Never	8	5.2	12	7.8	0.000*
Rarely	47	30.5	19	12.3	
Sometimes	54	35.1	46	29.9	
Often	25	16.2	49	31.8	
Always	20	13.0	28	18.2	
News					
Never	67	43.5	43	27.9	0.000*
Rarely	52	33.8	33	21.4	
Sometimes	18	11.7	37	24.0	
Often	15	9.7	29	18.8	
Always	2	1.3	12	7.8	
Sports sites					
Never	61	39.6	67	43.5	0.000*
Rarely	35	22.7	33	21.4	
Sometimes	28	18.2	28	18.2	
Often	17	11.0	13	8.4	
Always	13	8.4	13	8.4	
TV channels					
Never	47	30.5	42	27.3	0.000*
Rarely	50	32.5	32	20.8	
Sometimes	31	20.1	32	20.8	
Often	10	6.5	29	18.8	
Always	16	10.4	19	12.3	
Others					
Never	77	50.0	76	49.4	0.000*
Rarely	37	24.0	38	24.7	
Sometimes	21	13.6	19	12.3	
Often	9	5.8	8	5.2	
Always	10	6.5	13	8.4	

Chi-square test. \*Significant.

**Scoring of IAT in relation to gender, residence, and social characteristics**

There was no statistically significant difference in IAT scoring before or after COVID-19 regards to gender differences, residence, type of school, and educational level of teens, mother’s and father’s education, socio-economic level and partner whom does the child lives with them (p > 0.05) (Table 6).

**Predictors of internet addiction among the respondents**

By doing linear regression, taking IAT score as the dependent variable and other parameters were



**Table 4: Frequency distribution and percentage of the time of usage of Social Networking Services**

	Before COVID		After COVID		p-value
	n	%	n	%	
Facebook					
Never	25	16.2	19	12.3	0.000*
Rarely	28	18.2	20	13.0	
Sometimes	39	25.3	28	18.2	
Often	36	23.4	35	22.7	
Always	26	16.9	52	33.8	
Twitter					
Never	65	42.2	60	39.0	0.000*
Rarely	43	27.9	38	24.7	
Sometimes	33	21.4	35	22.7	
Often	8	5.2	11	7.1	
Always	5	3.2	10	6.5	
WhatsApp					
Never	7	4.5	3	1.9	0.000*
Rarely	10	6.5	5	3.2	
Sometimes	40	26.0	34	22.1	
Often	33	21.4	38	24.7	
Always	64	41.6	74	48.1	
Telegram					
Never	83	53.9	68	44.2	0.000*
Rarely	41	26.6	45	29.2	
Sometimes	20	13.0	23	14.9	
Often	7	4.5	13	8.4	
Always	3	1.9	5	3.2	
Snapchat					
Never	53	34.4	49	31.8	0.000*
Rarely	40	26.0	40	26.0	
Sometimes	36	23.4	31	20.1	
Often	11	7.1	8	5.2	
Always	14	9.1	26	16.9	
Instagram					
Never	37	24.0	32	20.8	0.000*
Rarely	32	20.8	32	20.8	
Sometimes	31	20.1	25	16.2	
Often	19	12.3	21	13.6	
Always	35	22.7	44	28.6	
Zoom					
Never	93	60.4	32	20.8	0.000*
Rarely	28	18.2	25	16.2	
Sometimes	16	10.4	48	31.2	
Often	10	6.5	25	16.2	
Always	7	4.5	24	15.6	
TikTok					
Never	69	44.8	51	33.1	0.000*
Rarely	31	20.1	25	16.2	
Sometimes	29	18.8	32	20.8	
Often	14	9.1	21	13.6	
Always	11	7.1	25	16.2	
IMO					
Never	100	64.9	84	54.5	0.000*
Rarely	35	22.7	29	18.8	
Sometimes	14	9.1	25	16.2	
Often	2	1.3	10	6.5	
Always	3	1.9	6	3.9	
Others					
Never	103	66.9	87	56.5	0.000*
Rarely	23	14.9	29	18.8	
Sometimes	15	9.7	18	11.7	
Often	7	4.5	7	4.5	
Always	6	3.9	13	8.4	

used as the independent variables, it was noticed that higher IAT score was associated with increasing educational levels of mothers and fathers, increasing time spent online, decreasing duration of sleep and lower number of days of exercise for more than 30 min/week ( $p < 0.05$ ) (Table 7).

**Table 5: Frequency distribution and percentage of internet addiction grades according to the scoring interpretation of IAT**

Total IAT score (Mean ± SD)	Before COVID		After COVID		p-value
	n	%	n	%	
45.26 ± 12.45					0.000*
61.91 ± 16.77					
Scoring IAT interpretation					
Normal internet usage	41	26.6	9	5.8	0.000*
Mild internet addiction	9	5.8	25	16.2	
Moderate internet addiction	94	61.0	101	65.6	
Severe internet addiction	10	6.5	19	12.3	

Chi-square test. \*Significant, IAT: Internet addiction test.

## Discussion

The idea of internet and smartphone addiction has become more important in light of the rising internet use caused by the decline in social connection during the COVID-19 shutdown [20]. As a result, action must be taken to stop or cure this addictive behavior, with the first step being the development of a precise diagnostic criterion [15].

Importantly, all schools were closed in Egypt during the COVID-19 outbreak, and at one time, online classes were developed to keep all types of pupils occupied. Homes were compelled to install internet access. With a highly significant difference ( $p = 0.001$ ), the overall internet addiction test score climbed from  $45.26 \pm 2.45$  before COVID-19 to  $61.91 \pm 16.77$  after COVID-19. IAT score was normal, mild, moderate, and severe in 5.8%, 16.2%, 65.6%, and 12.3% after COVID-19, respectively, compared to normal, mild, moderate, and severe IAT score in 26.6%, 5.8%, 61.0%, and 6.5% before COVID, respectively, with significant differences between them before and after COVID ( $p < 0.000$ ). We consider moderate and severe addictions were the categories that significantly relevant, so the prevalence of internet addiction among enrolled teens was increased from 67.5% before COVID-19 to 77.9% of 154 teens after COVID-19 infection. Similar to current findings, Abou El Wafa *et al.* [5] revealed that during the period of COVID-19 pandemic in Egypt and according to IAT score, the majority (74.65%) were problematic users, and 10.69% were classified as severe internet addicts.

Furthermore, Ilesanmi *et al.* [21] found that internet addiction rose from 7.7% to 64.3% before and after the COVID-19 pandemic, respectively, in a study of adolescents in Nigeria. Internet addiction was recently discovered in 63.2% of respondents in a study by Onukwuli *et al.* [22] (24.9% had mild, 59.6% had moderate, and 3.6% had severe) in southeast Nigeria.

The prevalence at the time was higher than what Duan *et al.* [23] found to be 23.5% (China), Albursan *et al.* [24] found to be 37.4% (Saudi Arabia), Paulus *et al.* found to be 43.69% (Germany) [25], Afolabi *et al.* [26] found to be 44.9% (Nigeria), Mokhtarinia *et al.* found to be 53.3% (Iran), and Eldesokey *et al.* [18], [27]. However, their study did not distinguish between mild, moderate, and severe internet addiction among Egyptian medical students. For instance, in Taiwan, the prevalence rate among adolescents increased from 17.4% to 24.4% during the pandemic (Lin 2020) [28].

Our study found a higher prevalence of internet addiction than some of the earlier studies. The timing of the many research and the populations examined may have contributed to this outcome. In addition,

**Table 6: Scoring of IAT in relation to gender, residence and social characteristics (expressed in Mean  $\pm$  SD)**

Variables	Before COVID score	F	p-value	After COVID score	F	p-value
Gender						
Male	55.60 $\pm$ 16.03	0.077	0.781	63.36 $\pm$ 17.09	1.182	0.279
Female	54.91 $\pm$ 14.93			60.42 $\pm$ 16.41		
Residence						
Urban	55.76 $\pm$ 15.24	0.629	0.429	62.38 $\pm$ 16.10	0.484	0.488
Abroad	53.29 $\pm$ 16.37			60.03 $\pm$ 19.36		
Type of school						
International	54.38 $\pm$ 15	1.216	0.305	59.83 $\pm$ 15.9	1.148	0.338
Language Private	56.57 $\pm$ 16.264			63.43 $\pm$ 17.711		
Arabic Private	56.68 $\pm$ 12.876			65.21 $\pm$ 13.286		
Language Experimental	59 $\pm$ 13.258			66.4 $\pm$ 16.426		
Arabic experimental	36.33 $\pm$ 4.509			45.67 $\pm$ 7.572		
State	53.38 $\pm$ 18.323			60.06 $\pm$ 20.43		
Educational level						
Primary	51.84 $\pm$ 17.365	1.157	0.332	58.42 $\pm$ 19.802	0.957	0.433
Preparatory	55.58 $\pm$ 17.994			61.55 $\pm$ 19.470		
Secondary	55.29 $\pm$ 13.907			61.98 $\pm$ 14.968		
Diploma	48.00 $\pm$ 0.00			67.00 $\pm$ 0.00		
University	61.17 $\pm$ 10.217			67.89 $\pm$ 9.480		
Mother's education						
Primary or Preparatory	46 $\pm$ 20.607	1.264	0.288	52.5 $\pm$ 23.923	1.335	0.265
Secondary or Diploma	46.88 $\pm$ 13.55			56.06 $\pm$ 21.306		
Graduate	55.23 $\pm$ 14.028			62.52 $\pm$ 15.596		
Postgraduate	59.13 $\pm$ 15.812			63.7 $\pm$ 16.332		
Father's education						
Primary or Preparatory	20 $\pm$ 0	1.141	0.321	20 $\pm$ 0	2.490	0.063
Secondary or Diploma	49.40 $\pm$ 16.987			58.2 $\pm$ 19.622		
Graduate	55.19 $\pm$ 14.982			62.42 $\pm$ 16.182		
Postgraduate	57.90 $\pm$ 14.894			62.98 $\pm$ 16.182		
Socioeconomic level						
Low	43.50 $\pm$ 33.234	0.974	0.380	45.0 $\pm$ 35.355	1.952	0.146
Medium	54.82 $\pm$ 14.857			61.15 $\pm$ 16.056		
High	57.47 $\pm$ 16.646			65.53 $\pm$ 17.877		
With whom does the child live?						
Two parents	54.60 $\pm$ 15.326	1.877	0.157	61.38 $\pm$ 17.063	0.979	0.378
Mother only	63.91 $\pm$ 15.915			68.73 $\pm$ 12.001		
Father only	54.00 $\pm$ 0			62 $\pm$ 0		

Independent t-test.

during the mandatory lockdown and after all, schools were closed when our study was conducted in the middle of 2020; people and institutions were required to follow certain COVID-19 safety procedures, which included social separation. Since social media was being utilized at the time to kill boredom and people had not yet undone the lifestyle adjustments they had made during the lockdown, it stands to reason that students' unrestrained internet use during the lockdown had led to addiction. Naturally, this could be the cause of the study's findings regarding the prevalence of internet addiction. There is evidence that the lockdown increased use of social media sites, particularly among young people, who became more dependent on it as it offered a convenient means of connecting with others and the outside world [29].

Similar to five studies [5], [24], [30], [31], [32], the current study found no statistically significant difference in IAT scoring before or after COVID-19 regarding gender differences. According to previous research [33], [34], [35], [36], [37], females consistently

scored on average much higher than males. However, other research [25], [27], [38], [39] have discovered higher scores and a larger chance of addiction among the male population.

Although content watched and motivation/justification may vary, the current results showed similar symptoms in both males and females and equivalent degrees of smartphone addiction [15].

It was intriguing to see from our study that 42.2% of the enrolled youths had laptops, 17.5% used tablets, and 87.7% used smartphones to access the internet. This might be due to the fact that multiple online schools developed during the pandemic and gave teen's access to mobile devices. This was corroborated by the fact that during the COVID-19 lockdown, roughly 50.6% of respondents mostly accessed the internet for assignments. However, a sizeable portion of them utilized the internet for social networking, particularly WhatsApp (72.8%) and Facebook (56.5%). This result is consistent with previous research that found teenagers in Hungary and Saudi Arabia spend a lot

**Table 7: Linear regression analysis of IAT score in relation to other parameters**

Predictor	Unstandardized Coefficients		Standardized Coefficients Beta	t	p-value	95.0% Confidence Interval for B	
	B	SE				Lower Bound	Upper Bound
Age of student (years)	0.687	0.435	0.128	1.580	0.116	-0.172-	1.547
Gender	-0.695-	2.498	-0.023-	-0.278-	0.781	-5.631-	4.241
Residence	-1.233-	1.555	-0.064-	-0.793-	0.429	-4.305-	1.839
Type of School	-0.382-	0.803	-0.039-	-0.476-	0.635	-1.968-	1.204
Socioeconomic level	3.362	2.849	0.095	1.180	0.240	-2.266-	8.991
Mother's Education	5.182	1.718	0.238	3.017	0.003*	1.788	8.576
Father's Education	4.689	1.943	0.192	2.414	0.017*	0.851	8.527
Duration of sleep	-5.423-	2.058	-0.209-	-2.634-	0.009*	-9.489-	-1.356-
Number of days per week of exercise more than 30 min	-2.253-	0.706	-0.251-	-3.193-	0.002*	-3.647-	-0.859-
Time spent online	5.062	1.695	0.232	3.002	0.003*	1.722	9.589
Educational level	1.881	1.055	0.143	1.782	0.077	-0.204-	3.966

of time communicating on social media, particularly Facebook and WhatsApp [26], [40]. The respondents' high internet usage was consistent with the evidence that internet usage among Egyptians had grown steadily from 41.79 million users in January 2020 to 59.66 million users in January 2021 (+18%) [7].

No statistically significant differences in IAT scores between enrolled teens before and after COVID-19 were found in relation to residence, type of school, educational level, mother's and father's education, socioeconomic status, or partner with whom the child resides ( $p > 0.05$ ).

These results were compared to those from the Onukwuli *et al.* 2023 [22] study, which showed that internet addiction in South-east Nigeria was significantly correlated with the respondent's age ( $p = 0.043$ ), mother's education level ( $p = 0.023$ ), family size ( $p = 0.021$ ), place of residence ( $p = 0.035$ ), alcohol intake ( $p = 0.017$ ), smoking ( $p = 0.015$ ), substance use ( $p = 0.001$ ), and length of internet use ( $p < 0.001$ ).

Using linear regression with the IAT score as the dependent variable and other parameters as the independent variables, it was discovered that higher IAT scores were correlated with higher levels of education for the parents, increased online time, shorter sleep duration, and fewer days of exercise lasting longer than 30 min/week ( $p = 0.05$ ).

A 2-h cutoff time interval was employed in previous research conducted in Bangladesh and Hungary [41], [42], according to those studies' findings. Additional research [27], [40], [43], [44] has demonstrated that a lower number of exercise days is an independent risk factor for internet addiction.

According to Abou El Wafa *et al.* [5], 55% of individuals who were classed as internet addicts were between the ages of 15 and 18; this shows that older age is substantially connected with internet addiction.

Internet addiction was recently found to be predicted by male gender, mid- and late teenage age groups of 14–19 years, as well as having used the internet for more than 6 months in a study by Onukwuli *et al.* (2023) [22].

In Paulus *et al.* [25], a multiple regression analysis was performed to predict the addiction score from gender, age, time spent using digital media, and the intensity of negative feelings during the COVID-19 epidemic. Higher levels of internet addiction were linked to factors such as female gender, advancing age, longer time spent using digital media, and higher negative emotion intensity during the COVID-19 pandemic.

This result confirms earlier research by Schimmenti *et al.* [45], which discovered that time spent online was a predictor of internet addiction as well. Although there is a strong correlation between time spent online and internet addiction, the cause and impact of this correlation cannot be determined by a

cross-sectional study. However, this discovery might be useful for both preventing and treating internet addiction. In addition, Lai *et al.* [46] found that the amount of time spent online is a risk factor for developing an internet addiction, even going so far as to assert that just an hour more of online time is enough to raise problematic behavior or addiction. Gentile *et al.* [47] demonstrated in a longitudinal design that higher gaming time is a substantial predictor of a subsequent gaming disorder, independent of the pandemic and exclusive to gaming. On the other hand, Yildiz Durak [48] discovered no statistically significant link between the length of social media use and problematic social media use.

Due to the rising amount of time, kids spend online and the much-increased rates of reliance following lockdown, these findings about digital media usage time become pertinent in the face of the pandemic [49] related to the most advanced forms of digital media (food channels, TikTok, Instagram, courses, reading, kids' websites, training websites, religious websites, drawing, viewing TV shows and movies, Facebook, Zoom, and TikTok services. In the study presented here, Eales *et al.* [50] discovered a significant increase in screen media use and problematic media use among children in the United States during the COVID-19 pandemic. Other services used included Netflix, Viber, YouTube, Twitch, and Pinterest. During COVID-19, Drouin *et al.* [51] observed an upsurge in smartphone usage. The likelihood of developing problematic patterns of use grows when young people spend more time playing games and using smartphones (Paschke *et al.*, 2021) [52].

According to Duan *et al.* [37], throughout the epidemic in China, about 30% of individuals spent more than five hours each day online. The amount of time spent online has increased since the epidemic began. It was shown that there was a mildly favorable correlation between internet and smartphone addiction and depression and anxiety. Later, according to Duan *et al.* [23], daily smartphone usage time, the severity of anxiety symptoms, and the type of coping strategy used all predict smartphone addiction.

The duration of internet use and the frequency of internet/smartphone addictions among teenagers both rose throughout the COVID-19 period, according to a systematic study by Yilmaz *et al.* [20]. However, studies on smartphone and internet addiction found no difference in gender. In addition, it has been discovered that adolescent internet and smartphone addictions are linked to mental illnesses, particularly post-traumatic stress disorder, anxiety disorders, and depression. Internet and smartphone addiction can be viewed as a risk factor for teenagers during the COVID-19 period when these findings are taken into consideration collectively.

To aid in the early detection of internet addiction during pandemics such as COVID-19, parents, and guardians should closely monitor their children's use of

the internet. They should keep in mind that factors like time spent online, sleep duration, and daily exercise may increase the likelihood that internet addiction will develop in a particular population of teenagers. Furthermore, to reduce the danger of developing physical and mental illnesses, parents should think about how they can regulate and track their children's smartphone usage.

### **Limitations of the study**

The cross-sectional aspect of the research, which must be taken into account when interpreting and generalizing the results, is one of the study's weaknesses.

The youngsters' inability to recall exactly how many hours they spent online may have introduced bias. Therefore, the potential for recollection bias may have an impact on our study.

Before participating in the study, individuals were not clinically evaluated to ascertain their mental and physical health; nevertheless, those who had a history of serious health issues were not permitted to take part.

This study's generalizability to broader time periods or lockdown-free situations is constrained because it was carried out during the COVID-19 epidemic.

## **Conclusions**

In Egypt, teens have a high frequency of internet addiction (77.9%), according to our study. The amount of time a youngster spent online, the education level of their parents, how much they slept, how many days they exercised for more than 30 min each week, and how much time they spent online were all linked to internet addiction.

## **Declarations**

### **Ethics approval and consent to participate**

The IRB of the National Research Centre in Giza, Egypt, gave its approval to this study. In addition, each participant's informed consent was gained by a statement of agreement at the start of each questionnaire. All procedures were also carried out in accordance with the Declaration of Helsinki and all applicable rules and regulations. Teenagers were emailed a link to an online survey form that was labeled with the stud's goal. The online questionnaire's completion and submission were taken into account as

the parent's written approval of his child's involvement in the study.

### **Availability of data and material**

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

### **Authors' contributions**

Study design, data collection, data analysis, writing the Manuscript, all these parts were a shared work between all authors with variable but finally equal amounts.

## **References**

1. Scarabel L, Guardascione M, Dal Bo M, Toffoli G. Pharmacological strategies to prevent SARS-CoV-2 infection and treat the early phases of COVID-19. *Int J Infect Dis.* 2021;104:441-51. <https://doi.org/10.1016/j.ijid.2021.01.035> PMID:33476760
2. Guessoum SB, Lachal J, Radjack R, Carretier E, Minassian S, Benoit L, et al. Adolescent psychiatric disorders during the COVID-19 pandemic and lockdown. *Psychiatry Res.* 2020;291:113264. <https://doi.org/10.1016/j.psychres.2020.113264> PMID:32622172
3. Duner P, Salehi A. COVID-19 and possible pharmacological preventive options. *J Clin Med Res.* 2020;12(12):758-72. <https://doi.org/10.14740/jocmr4383> PMID:33447309
4. Raquib A, Raquib R, Jamil S, Hossain A, Al-Mamun F, Mamun MA. Knowledge, attitudes, and practices toward the prevention of COVID-19 in Bangladesh: A systematic review and meta-analysis. *Front Med (Lausanne).* 2022;9:1425. <https://doi.org/10.3389/fmed.2022.856156> PMID:35733876
5. Abou El Wafa HE, Ashry MH, Mekky JF, Elweshahi HM. Internet abuse in children and adolescents during COVID-19 pandemic in Egypt. *Egypt J Psychiatry.* 2021;42(1):36-42. [https://doi.org/10.4103/ejpsy.ejpsy\\_43\\_20](https://doi.org/10.4103/ejpsy.ejpsy_43_20)
6. Evli M, Şimşek N. The effect of COVID-19 uncertainty on internet addiction, happiness and life satisfaction in adolescents. *Arch Psychiatr Nurs.* 2022;41:20-6. <https://doi.org/10.1016/j.apnu.2022.07.008> PMID:36428049
7. Statista. Number of Smartphone Users Worldwide from 2014 to 2023; 2021. Available from: <https://www.statista.com/statistics/330695/number-of-smartphone-users-worldwide> [Last accessed on 2023 Jan 14].
8. Masaeli N, Farhadi H. Prevalence of internet-based addictive behaviors during Covid-19 pandemic: A systematic review. *J Addict Dis.* 2021;39(4):468-88. <https://doi.org/10.1080/10550887.2021.1895962> PMID:33749537
9. Serra G, Lo Scalzo L, Giuffrè M, Ferrara P, Corsello G. Smartphone use and addiction during the coronavirus disease 2019 (COVID-19) pandemic: Cohort study on 184 Italian



- children and adolescents. *Ital J Pediatr.* 2021;47(1):150. <https://doi.org/10.1186/s13052-021-01102-8>  
PMid:34215311
10. Chopdar PK, Paul J, Prodanova J. Mobile shoppers' response to Covid-19 phobia, pessimism and Smartphone addiction: does social influence matter? *Technol Forecast Soc Change.* 2022;174:121249. <https://doi.org/10.1016/j.techfore.2021.121249>  
PMid:36540714
  11. Hu Q, Liu Q, Wang Z. Meaning in life as a mediator between interpersonal alienation and Smartphone addiction in the context of Covid-19: A three-wave longitudinal study. *Comput Hum Behav.* 2022;127:107058. <https://doi.org/10.1016/j.chb.2021.107058>  
PMid:34690416
  12. de Freitas BH, Gaíva MA, Diogo PM, Bortolini J. Self-reported Smartphone addiction among Brazilian adolescents in the COVID-19 pandemic context: A mixed-method study. *Trends Psychol.* 2022:1-20. <https://doi.org/10.1007/s43076-022-00208-0>
  13. Zhao J, Ye B, Luo L, Yu L. The effect of parent phubbing on Chinese adolescents' Smartphone addiction during COVID-19 pandemic: Testing a moderated mediation model. *Psychol Res Behav Manag.* 2022;15:569-79. <https://doi.org/10.2147/PRBM.S349105>  
PMid:35281997
  14. Hong W, Liu RD, Oei TP, Zhen R, Jiang S, Sheng X. The mediating and moderating roles of social anxiety and relatedness need satisfaction on the relationship between shyness and problematic mobile phone use among adolescents. *Comput Hum Behav.* 2019;93:301-8.
  15. Yue H, Yue X, Liu B, Li X, Dong Y, Bao H. Short version of the Smartphone addiction scale: Measurement invariance across gender. *PLoS One.* 2023;18(3):e0283256. <https://doi.org/10.1371/journal.pone.0283256>  
PMid:36947534
  16. Saadeh H, Al Fayed RQ, Al Refaei A, Shewaikani N, Khawaldah H, Abu-Shanab S, *et al.* Smartphone use among university students during COVID-19 quarantine: An ethical trigger. *Front Public Health.* 2021;9:600134. <https://doi.org/10.3389/fpubh.2021.600134>  
PMid:34381747
  17. de Freitas BH, Gaíva MA, Diogo PM, Bortolini J. Relationship between lifestyle and self-reported smartphone addiction in adolescents in the COVID-19 pandemic: A mixed-methods study. *J Pediatr Nurs.* 2022b;65:82-90. <https://doi.org/10.1016/j.pedn.2022.03.001>  
PMid:35331609
  18. Eldesokey S, Gomaa Z, Sabri Y, El-Gilany A, Mohamed Elwasify M. Smartphone addiction among medical students in Mansoura university. *Egypt J Psychiatry.* 2021;42:50-6.
  19. Young KS. Internet addiction: The emergence of a new clinical disorder. *CyberPsychol Behav.* 1998;1:237-44. <https://doi.org/10.1089/cpb.1998.1.237>
  20. Yilmaz YA, Aydogdu F, Kocturk N. A systematic overview of adolescents' smartphone habits and internet addiction during the COVID-19 era. *Curr Approach Psychiatry.* 2022;14(4):510-7. <https://doi.org/10.18863/pgy.1064501>
  21. Ilesanmi OS, Afolabi AA, Adebayo AM. Problematic internet use (PIU) among adolescents during COVID-19 lockdown: A study of high school students in Ibadan, Nigeria. *The African Journal of Information and Communication.* 2021;27:1-22.
  22. Onukwuli VO, Onyinye EN, Udigwe IB, Umeh UM. Internet addiction during the COVID-19 pandemic among adolescents in southeast Nigeria and implications for adolescent care in the post-pandemic era: A cross-sectional study. *SAGE Open Med.* 2023;11:20503121231152763. <https://doi.org/10.1177/20503121231152763>  
PMid:36860685
  23. Duan L, He J, Li M, Dai J, Zhou Y, Lai F, *et al.* Based on a decision tree model for exploring the risk factors of Smartphone addiction among children and adolescents in China during the COVID-19 pandemic. *Front Psychiatry.* 2021;12:652356. <https://doi.org/10.3389/fpsy.2021.652356>  
PMid:34168575
  24. Albusan IS, Al Qudah MF, Al-Barashdi HS, Bakhiet SF, Darandari E, Al-Asqah SS, *et al.* Smartphone addiction among university students in light of the COVID-19 pandemic: Prevalence, relationship to academic procrastination, quality of Life, gender and educational stage. *Int J Environ Res Public Health.* 2022;19(16):10439. <https://doi.org/10.3390/ijerph191610439>  
PMid:36012073
  25. Paulus FW, Joas J, Gerstner I, Kühn A, Wenning M, Gehrke T, *et al.* problematic internet use among adolescents 18 months after the onset of the COVID-19 pandemic. *Children (Basel).* 2022;9(11):1724. <https://doi.org/10.3390/children9111724>  
PMid:36360452
  26. Afolabi AA, Ilesanmi OS, Adebayo AM. Prevalence and pattern of internet addiction among adolescents in Ibadan, Nigeria: A cross-sectional study. *Cureus.* 2022;14(2):e22293. <https://doi.org/10.7759/cureus.22293>  
PMid:35350489
  27. Mokhtarinia HR, Torkamani MH, Farmani O, Biglarian A, Gabe CP. Smartphone addiction in children: Patterns of use and musculoskeletal discomfort during the COVID-19 pandemic in Iran. *BMC Pediatr.* 2022;22(1):681. <https://doi.org/10.1186/s12887-022-03748-7>  
PMid:36435748
  28. Lin MP. Prevalence of internet addiction during the covid-19 outbreak and its risk factors among junior high school students in Taiwan. *Int J Environ Res Public Health.* 2020;17(22):8547. <https://doi.org/10.3390/ijerph17228547>  
PMid:33218018
  29. Fernandes B, Biswas UN, Tan-Mansukhani R, Vallejo A, Essau CA. The impact of COVID-19 lockdown on internet use and escapism in adolescents. *Rev Psicol Clin Ninos Adolesc.* 2020;7:59-65. <https://doi.org/10.21134/rpcna.2020.mon.2056>
  30. Kumar VA, Chandrasekaran V, Brahadeeswari H. Prevalence of Smartphone addiction and its effects on sleep quality: A cross-sectional study among medical students. *Ind Psychiatry J.* 2019;28(1):82-5. [https://doi.org/10.4103/ipj.ipj\\_56\\_19](https://doi.org/10.4103/ipj.ipj_56_19)  
PMid:31879452
  31. Mescollotto FF, Castro EM, Pelai EB, Pertille A, Bigaton DR. Translation of the short version of the Smartphone Addiction Scale into Brazilian Portuguese: Cross-cultural adaptation and testing of measurement properties. *Braz J Phys Ther.* 2019;23(3):250-6. <https://doi.org/10.1016/j.bjpt.2018.08.013>  
PMid:30249438
  32. Harris B, McCredie M, Fields S. Examining the psychometric properties of the Smartphone Addiction Scale and its short version for use with emerging adults in the U.S. *Comput Hum Behav Rep.* 2020;1:100011. <https://doi.org/10.1016/j.chbr.2020.100011>
  33. Kwon M, Kim DJ, Cho H, Yang S. The Smartphone addiction scale: Development and validation of a short version for adolescents. *PLoS One.* 2013;8(12):e83558. <https://doi.org/10.1371/journal.pone.0083558>  
PMid:24391787
  34. Chi X, Hong X, Chen X. Profiles and socio-demographic correlates of Internet addiction in early adolescents in southern

- China. *Addict Behav.* 2020;106:106385. <https://doi.org/10.3389/fpubh.2022.944085>  
PMid:35812472
35. Tateno M, Kim DJ, Teo AR, Skokauskas N, Guerrero AP, Kato TA. Smartphone addiction in Japanese college students: Usefulness of the Japanese version of the smartphone addiction scale as a screening tool for a new form of internet addiction. *Psychiatry Investig.* 2019;16(2):115-20. <https://doi.org/10.30773/pi.2018.12.25.2>  
PMid:30808117
36. Tomaszek K, Muchacka-Cymerman A. Sex differences in the relationship between student school burnout and problematic internet use among adolescents. *Int J Environ Res Public Health.* 2019;16(21):4107. <https://doi.org/10.3390/ijerph16214107>  
PMid:31653105
37. Duan L, Shao X, Wang Y, Huang Y, Miao J, Yang X, et al. An investigation of mental health status of children and adolescents in china during the outbreak of COVID-19. *J Affect Disord.* 2020;275:112-8. <https://doi.org/10.1016/j.jad.2020.06.029>  
PMid:32658812
38. Andrade AL, Kim DJ, Scatena A, Enes CC, Enumo SR, De Micheli D. Validity and reliability of the Brazilian version of the smartphone addiction scale-long version (SAS-LV). *Trends Psychol.* 2020;29:302-19. <https://doi.org/10.1007/s43076-020-00046-y>
39. Chatterjee S, Kar SK. Smartphone addiction and quality of sleep among Indian medical students. *Psychiatry.* 2021;84(2):182-91. <https://doi.org/10.1080/00332747.2021.1907870>  
PMid:33856961
40. Kapus K, Nyulas R, Nemeskeri Z, Zadori I, Muity G, Kiss J, et al. Prevalence and risk factors of internet addiction among Hungarian high school students. *Int J Environ Res Public Health.* 2021;18(13):6989. <https://doi.org/10.3390/ijerph18136989>  
PMid:34208800
41. Mamun M, Griffiths MD. The assessment of internet addiction in Bangladesh: Why are prevalence rates so different? *Asian J Psychiatry.* 2019;40:46.
42. Chandrima RM, Kircaburun K, Kabir H, Riaz BK, Kuss DJ, Griffiths MD, et al. Adolescent problematic internet use and parental mediation: A Bangladeshi structured interview study. *Addict Behav Rep.* 2020;12:100288. <https://doi.org/10.1016/j.abrep.2020.100288>  
PMid:32637564
43. Cha SS, Seo BK. Smartphone use and Smartphone addiction in middle school students in Korea: Prevalence, social networking service, and game use. *Health Psychol Open.* 2018;5(1):2055102918755046. <https://doi.org/10.1177/2055102918755046>  
PMid:29435355
44. Tenzin K, Dorji T, Choeda T, Wangdi P, Oo MM, Tripathy JP, et al. Internet addiction among secondary school adolescents A mixed methods study. *JNMA J Nepal Med Assoc.* 2019;57(219):344-51. <https://doi.org/10.31729/jnma.4292>  
PMid:32329462
45. Schimmenti A, Musetti A, Costanzo A, Terrone G, Maganuco N, Rinella C, et al. The unfabulous four: Maladaptive personality functioning, insecure attachment, dissociative experiences, and problematic internet use among young adults. *Int J Ment Health Addict.* 2021;19:447-61. <https://doi.org/10.1007/s11469-019-00079-0>
46. Lai CM, Mak KK, Watanabe H, Ang RP, Pang JS, Ho RC. Psychometric properties of the internet addiction test in Chinese adolescents. *J Pediatr Psychol.* 2013;38(7):794-807. <https://doi.org/10.1093/jpepsy/jst022>  
PMid:23671059
47. Gentile DA, Choo H, Liau A, Sim T, Li D, Fung D, et al. Pathological video game use among youths: A two-year longitudinal study. *Pediatrics.* 2011;127(2):e319-29. <https://doi.org/10.1542/peds.2010-1353>  
PMid:21242221
48. Yıldız Durak H. Modeling of variables related to problematic internet usage and problematic social media usage in adolescents. *Curr Psychol.* 2020;39:1375-87.
49. Sun Y, Li Y, Bao Y, Meng S, Sun Y, Schumann G, et al. Brief report: increased addictive internet and substance use behavior during the COVID-19 pandemic in China. *Am J Addict.* 2020;29:268-70. <https://doi.org/10.1111/ajad.13066>  
PMid:32500608
50. Eales L, Gillespie S, Alstat RA, Ferguson GM, Carlson SM. Children's screen and problematic media use in the United States before and during the COVID-19 pandemic. *Child Dev.* 2021;92:e866-82.
51. Drouin M, McDaniel BT, Pater J, Toscos T. How parents and their children used social media and technology at the beginning of the COVID-19 pandemic and associations with anxiety. *Cyberpsychol Behav Soc Netw.* 2020;23(11):727-36. <https://doi.org/10.1089/cyber.2020.0284>  
PMid:32726144
52. Paschke K, Austermann M, Simon-Kutscher K, Thomasius R. Adolescent gaming and social media usage before and during the COVID-19 pandemic interim results of a longitudinal study. *Sucht.* 2021;67:13-22.