

# Ethanollic Extract of *Emilia sonchifolia* Leaves Possess Erythropoietic and Hepatoprotective Effect in Mice Infected with *Plasmodium Berghei Berghei*

Innocent A. Edagha<sup>1\*</sup>, Koofreh G. Davies<sup>2</sup>, Blessing C. Akpan<sup>3</sup>, Christopher C. Mbadugha<sup>1</sup>, Wonderful U. Udoiso<sup>2</sup>

<sup>1</sup>Department of Anatomy, Faculty of Basic Medical Sciences, University of Uyo, P.M.B 1017, Uyo, Nigeria; <sup>2</sup>Department of Physiology, Faculty of Basic Medical Sciences, University of Uyo, P.M.B 1017, Uyo, Nigeria; <sup>3</sup>Chemical Pathology Unit, Laboratory Department, Federal Psychiatric Hospital Calabar, P.M.B 1052; Calabar, Nigeria

## Abstract

**Citation:** Edagha IA, Davies KG, Akpan BC, Mbadugha CC, Udoiso WU. Ethanollic Extract of *Emilia Sonchifolia* Leaves Possess Erythropoietic and Hepatoprotective Effect in Mice Infected with *Plasmodium Berghei Berghei*. OA Maced J Med Sci. 2014 Mar 15; 2(1):11-17.  
http://dx.doi.org/10.3889/oamjms.2014.002

**Key words:** *Plasmodium berghei berghei*, *Emilia sonchifolia*, Haematology, Liver, Mice.

**\*Correspondence:** Mr. Innocent Asuquo Edagha. Department of Anatomy, Faculty of Basic Medical Sciences, University of Uyo, P.M.B 1017, Uyo, Nigeria. E-Mail: iedagha@yahoo.com

**Received:** 08-Nov-2013; **Revised:** 19-Dec-2013; **Accepted:** 20-Dec-2013; **Online first:** 23-Jan-2014

**Copyright:** © 2014 Edagha et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Competing Interests:** The authors have declared that no competing interests exist.

**Aim:** This study was designed to investigate the effect of ethanollic extracts of the leaves of *Emilia sonchifolia* on the haematological parameters and histomorphology of the liver of male Swiss albino mice infected with *Plasmodium berghei berghei* (*Pbb*).

**Material and Methods:** 35 mice were divided into; Group 1 (control) given normal saline 0.3 ml, Group 2 passaged with *Pbb* only, Group 3 passaged with *Pbb*, and then treated with Coartem®, Group 4 treated with *E. sonchifolia* 325 mg/kg only, Group 5 treated with *E. sonchifolia* 650 mg/kg only, Group 6 passaged with *Pbb* then treated with *E. sonchifolia* 325 mg/kg, while Group 7 was passaged with *Pbb* then treated with *E. sonchifolia* 650 mg/kg. *Pbb* was passaged intraperitoneally, while the test drug and extracts was given via orogavage once daily.

**Results:** The result showed significantly ( $P < 0.001$ ) reduced RBC parameters at in Group 5 treated with 650 mg/kg similar with Group 2 compared to Group 1, while there was significant ( $P < 0.01$ ) increased WBC and differentials in Parasitized groups compared with Group 1. The micrographs showed slightly inflamed nuclei in Group 4, with few nuclei shrinkage Group 5, whereas in the parasitized groups treated with the extract there appeared to be hepatoprotection compared to Group 2.

**Conclusion:** In conclusion, the extract promotes erythropoiesis at 325 mg/kg, but was haemolytic at 650 mg/kg, and exerts its effect possibly through an agonistic and a synergistic activity of its rich bioactive ingredients. It showed mild toxic effect in the histomorphology of the non-parasitized mice at 325 mg/kg and 650 mg/kg, and also appeared to offer hepatoprotection in parasitized mice compared to the parasitized group that had no treatment.

## Introduction

*Emilia sonchifolia* (Lin.) is a bushy annual herb distributed mainly in Asian countries [1]. It has been traditionally used as important medicinal plant in most tropical and subtropical countries, including in the South-South region of Akwa Ibom State, Nigeria. Globally, an estimated 3.3 billion people were at risk of malaria in 2011, with populations living in sub-Saharan Africa having the highest risk of acquiring malaria [2]. *Emilia sonchifolia* has also been reported to possess anti-fever activities [3, 4] antimicrobial activity [5] analgesic and anti-inflammatory activities [6-8]; anticancer activities [9-11]; antioxidant activities [12-15]; anti-diabetic [16]; anti-cataract activities [17-20]; anticonvulsant activity [21]; antinociceptive effect [22].

Compounds like simiral, beta-sitosterol, stigmasterol, palmitic acid and honey acid were obtained from the whole plant of *E. sonchifolia* [23].

Few pyrrolizidine alkaloids, senkirkine and doronine were isolated from the aerial parts of *E. sonchifolia* [24].

There is a growing disillusionment with modern medicine and also the misconception that herbal remedy, being natural may be devoid of adverse and toxic effects often associated with allopathic medicines [25]. The dangers associated with the potential toxicity of herbal therapies demand that herbal practitioners be kept abreast of the report on renal and hepatic toxicity resulting from ingestion of medicinal herbs [26]. Researchers with interest in natural products have intensified their efforts towards scientific evaluation of traditional medicines [27].

There has been no report on the effect of *Emilia sonchifolia* on haematological indices as well as histomorphology of the liver in a parasitized mice model; hence this study was designed to contribute this information.

## Materials and Methods

### Experimental animals

Thirty five male Swiss albino mice were obtained from the animal holding facility of Faculty of Basic Medical Sciences, University of Uyo, Uyo, Nigeria and acclimatized for two weeks before the start of the experiment. They were allowed access to water and feed *ad libitum*. All procedures involving animals in this study conformed to the guiding principles in the care and use of animals [28] and the institution's code of ethics for the use of laboratory animals.

### Plant collection

Fresh leaves of *Emilia sonchifolia* was obtained at the medicinal farm of the Department of Pharmacology and Toxicology, University of Uyo during the October period of 2012. They were identified and authenticated by the Curator at the Herbarium with voucher numbers UUH/10(e) for *Emilia sonchifolia* deposited.

### Plant extraction

The extraction was done with 700 g of fresh leaves of *Emilia sonchifolia*, macerated in 96% ethanol (Sigma Aldrich, Germany) in a flat bottom flask and were kept for 72 hours at room temperature. The macerated leaves were then filtered and the filtrate concentrated in water-bath at 45°C to dryness with a yield of 15.71 g.

### Parasite inoculation

Each mouse used in the experiment was inoculated intraperitoneally with 0.3 ml of infected blood containing about  $1 \times 10^7$  *Plasmodium berghei berghei* parasitized erythrocytes. The inoculum consisted of  $5 \times 10^7$  *Plasmodium berghei berghei* erythrocytes per ml. This was prepared by determining both the percentage parasitaemia and the erythrocytes count of the donor mouse and diluting the blood with isotonic saline in proportions indicated by both determinations [29]. The origin of the parasites was from 3 donor mice of a Chloroquine sensitive strain of *Plasmodium berghei berghei* (ANKA) obtained from the National Institute of Medical Research (NIMR) Yaba, Lagos, and was maintained via cycles of subpassage at the Department of Pharmacology and Toxicology, University of Uyo, where they were monitored until the scheduled time when experimental mice were infected.

### Dosage

All extracts dosage was determined after toxicity test (LD<sub>50</sub>) Median lethal dose, which used the

modified Lorke's method [30]. The 10% and 20% of the LD<sub>50</sub> of the extracts was administered as low and medium dose. Eighteen mice divided into six groups of three mice each were administered with the ethanolic extract (1000 mg/kg, 3000 mg/kg, 3500 mg/kg, 4000 mg/kg, 4500 mg/kg and 5000 mg/kg respectively), and the mice were fasted for 24 hours prior to the administration of the extract and 3 hours before testing, drinking water was removed. The manifestation of physical signs of toxicity such as; writhing, restlessness, decreased motor activity, aggressiveness, weakness, gasping for air and possible death was recorded within 24 hours.

### Drug

Coartem® Dispersible tablets, an anti-malarial agent (Artemether 20 mg/Lumefantrine 120 mg) manufactured by Novartis Pharmaceutical Corporation Suffern, New York, USA for Novartis Pharma AG Basle, Switzerland under licence from the PRC with NAFDAC REG. NO: A4-1680 was purchased from a Pharmacy in Uyo metropolis, and single tablets on each successive days was dissolved in normal saline, then administered in the test group based on body weight.

### Phytochemical screening

The preliminary phytochemical constituents of the leaves was determined [31]

### Experimental design

Group 1 served as control and the other 6 groups served as experimental groups. Group 1-which served as the Control and was given (0.3ml) normal saline (11 days), Group 2-were passaged with *Plasmodium berghei berghei* (11 days), Group 3-were passaged with *Plasmodium berghei berghei* (6 days) and then treated with Coartem® (5 days), Group 4-were administered with *E. sonchifolia* 325 mg/kg (11 days), Group 5-were administered with *E. sonchifolia* 650 mg/kg (11 days), Group 6-were passaged with *Plasmodium berghei berghei* (6 days) then treated with *E. sonchifolia* 325 mg/kg (5 days), Group 7-were passaged with *Plasmodium berghei berghei* (6 days), then treated with *E. sonchifolia* 650 mg/kg (5 days). Parasitized animals were passaged once intraperitoneally lasting for duration of 6 days before treatment with extract commenced once daily for 5 days, except in group 2 where there was no treatment.

### Determination of haematological parameters

Blood was collected from the left ventricle of each animal in a vial containing 0.5 M EDTA. Haematological indices were determined after day 11

of treatment using an Automated Mindray BC-5300 Haematolog Analyzer Made in China at the University of Uyo Teaching Hospital.

#### Tissue collection

Each mouse was humanely sacrificed by chloroform inhalation and the liver was dissected, immediately weighed and rinsed with normal saline and fixed in 10% neutral buffered formaldehyde for light microscopy investigation [31].

#### Statistical analysis

One way analysis of variance (ANOVA) was applied to compare the relationship of the groups, and Dunnett post-hoc test was used to compare the experimental groups and the control. All values were presented as mean  $\pm$  standard error of mean (SEM), and values were considered significant at  $p < 0.05$ .

## Results

#### Phytochemical constituents of Emilia sonchifolia

The results of the preliminary phytochemical screening showed that *E. sonchifolia* was positive for the presence of alkaloids, flavonoids, saponins, tannins, terpenes, and cardiac glycosides at varying degrees of lowly, moderately and highly present (Table 1).

**Table 1: Phytochemical Constituents of Emilia sonchifolia.**

Phytochemical	Emilia sonchifolia
Alkaloids	+++
Flavonoids	++
Phlobatannins	-
Saponin (Frothings' test)	+
Saponins (Fehlings' solution + $\text{Na}_2\text{CO}_3$ )	++
Saponins (Fehlings' solution)	++
Tannins	++
Terpenes	+
Anthraquinones	-
Deoxy-sugar	-
Cardiac glycosides	++

Method: Trease and Evans, 1989 [32]; Key: + = lowly present, ++ = moderately present, +++ = highly Present, - = absent

#### Effect on haematological indices

The result from Table 3 shows that the RBC in Group 2 passaged with *Plasmodium berghei berghei* was significantly ( $P < 0.001$ ) reduced compared to the Group 1 (control), and a similar value was obtained for other RBC parameters like HGB, PCV, MCH and MCHC. Group 5 treated with *E. sonchifolia* at 650mg/kg was also significantly ( $P < 0.001$ ) reduced compared to the Group 1 (control). However, the WBC in Group 2 passaged with *Plasmodium berghei berghei* showed a significantly ( $P < 0.01$ ) increased value compared to Group 1 (control), and this was similar in Group 5 treated with *E. sonchifolia* at 650 mg/kg, while Group 3 passaged

with *Plasmodium berghei berghei*, then treated with Coartem® had a WBC significantly increased at ( $P < 0.05$ ).

**Table 2: Acute Toxicity Test ( $\text{LD}_{50}$ ) for Emilia sonchifolia.**

Groups (n= 3)	Dosage (mg/kg)	Mortality	% Mortality
Group 1	1000	0/3	0
Group 2	3000	0/3	0
Group 3	3500	3/3	100
Group 4	4000	3/3	100
Group 5	4500	3/3	100
Group 6	5000	3/3	100

$\text{LD}_{50} = \sqrt{ab}$ ; a = 3000; b = 3500;  $\text{LD}_{50} = \sqrt{3000 \times 3500}$  (mg/kg) = 3250mg/kg. 10% of 3250.00 mg/kg = 325.00 mg/kg = low dose; 20% of 3250.00 mg/kg = 650.00 mg/kg = Medium dose.

Neutrophils was only significantly ( $P < 0.05$ ) increased in Group 2 passaged with *Plasmodium berghei berghei* compared to Group 1 (control), whereas Lymphocytes was only significantly ( $P < 0.05$ ) increased in Group 4 treated with *E. sonchifolia* at 325 mg/kg compared to Group 1 (control). The result of Platelet indicated that Group 2 passaged with *Plasmodium berghei berghei* was significantly ( $P < 0.001$ ) reduced compared to Group 1 (control); while Group 6 passaged with *Plasmodium berghei berghei*, and then treated with *E. sonchifolia* 325 mg/kg was significantly ( $P < 0.01$ ) reduced compared to Group 1 (control), whereas Groups 4 and 5 were both significantly reduced compared to Group 1 (control) at ( $P < 0.05$ )

#### Effects on the histology of the liver

Group 1 showed normal histological architecture of the liver with the Central vein (Cv), the plates of hepatic cells (Hc), Portal vein (Pv), Endothelial cells (Ec), Sinusoids (S) all appearing unaffected (Figure 1A). Group 2 passaged with *Plasmodium berghei berghei* only showed; Hypertrophy, Inflammation, Numerous Karyorrhectic hepatic cells (Kc) and Necrosis (N), and was strongly affected. Central vein (Cv), the Portal vein (Pv), plates of hepatic cells (Hc), sinusoids (S) (Figure 1B). Group 3 passaged with *Plasmodium berghei berghei*, and then treated with + Coartem® showed the Portal vein (Pv), Central vein (Cv), plates of hepatic cells (Hc), Sinusoids (S), sparse karyorrhectic cells (Kc) and Necrosis (N), few inflamed areas (If) and is strongly affected (Figure 1C), compared to the control. Group 4 treated with *Emilia sonchifolia* 325mg/kg showed slightly Inflamed nuclei, the Central vein (Cv), portal vein (Pv), Bile duct (Bd), Hepatic artery (Ha), Binucleate cells (Figure 1D), mildly affected compared to the control. Group 5 treated with *Emilia sonchifolia* 650mg/kg showed the Central Vein (Cv), Portal Vein (Pv), Hepatic artery (Ha), Bile duct (Bd), Lymphatic vessels (Lv), Sinusoid (S), plate of hepatic cells (Hc).


**Table 3: Effect of the ethanolic extract of *Emilia sonchifolia* on haematological parameters in male mice infected with *Plasmodium berghei berghei*.**

Groups	RBC	HGB	PCV	MCV	MCH	MCHC	WBC	NEU	LYM	PLT
Group 1 Normal Saline 0.3 ml	7.21±0.35	111.00±2.49	40.10±1.20	55.28±0.51	15.40±0.12	277.00±2.24	7.27±0.58	33.24±3.98	61.32±4.52	1086.28±30.70
Group 2 PBB only	2.39±0.73***	41.00±8.79***	17.26±4.11***	73.50±2.43***	18.04±0.99**	243.40±5.06**	36.37±11.32**	48.06±6.07*	50.56±6.66	441.00±106.07***
Group 3 PBB+Coartem®	6.72±0.86	98.20±10.15	35.12±2.23	54.14±5.03	14.36±0.65	275.40±14.15	28.16±13.48*	38.64±8.78	60.00±9.36	955.00±93.24
Group 4 ES 325 g/kg	7.60±0.18	116.20±1.74	40.04±0.81	52.36±1.04	15.20±0.38	290.20±1.32	16.22±3.19	18.22±3.63	80.12±4.36*	807.40±31.58*
Group 5 ES650 mg/kg	2.31±0.22***	39.40±1.89***	17.40±0.97***	83.62±4.17***	18.70±0.92***	226.40±2.89***	42.29±4.69**	30.44±2.42	69.20±2.42	732.00±95.08*
Group 6 PBB+ ES 325 g/kg	5.52±1.06	85.40±14.08	30.24±4.94	55.54±1.75	15.56±0.51	281.40±1.44	21.24±6.89	32.40±3.52	65.58±4.26	757.00±148.10**
Group 7 PBB+ ES 650 mg/kg	7.36±0.58	113.40±10.59	38.66±3.40	52.50±1.09	15.28±0.55	290.20±3.50	13.10±2.27	29.48±2.76	68.18±2.95	944.20±113.67

Values in means + S.E. (Standard error), n=5, \*P<0.05, \*\*P<0.01, \*\*\*P<0.001, when compared with control.

It showed few nuclei shrinkage and some inflamed nuclei (Figure 1E), compared to the control. Group 6 passaged with *Plasmodium berghei berghei*, and then treated with *E. sonchifolia* 325 mg/kg showed the Portal vein (Pv), Bile duct (Bd), Hepatic

artery (Ha), few karyorrhectic cells among healthy hepatic cells (Kc), few inflammation (If) and clumping cells, and was strongly affected (Figure 1F), compared to the control.



**Figure 1: Photomicrographs showing histological sections of the liver of (A) Group 1 served as Control administered (0.3 ml) normal, (B) Group 2 were passaged with *Plasmodium berghei berghei* only, (C) Group 3 were passaged with *Plasmodium berghei berghei* and then treated with Coartem®, (D) Group 4 were administered with *E. sonchifolia* 325 mg/kg only, (E) Group 5 were administered with *E. sonchifolia* 650 mg/kg only, (F) Group 6 were passaged with *Plasmodium berghei berghei* then treated with *E. sonchifolia* 325 mg/kg only, (G) Group 7 were passaged with *Plasmodium berghei berghei* then treated with *E. sonchifolia* 650 mg/kg, at magnification of 400 X stained with Haematoxylin and Eosin.**

Group 7 *Plasmodium berghei berghei* and then treated with *E. sonchifolia* 650 mg/kg showed the portal vein (Pv), Bile duct (Bd), few karyorrhectic cells (Kc), Clumping nuclei (Cn) and Inflammation (If) was strongly affected (Figure 1G), compared to the control.

## Discussion

This study was designed to determine the LD<sub>50</sub> of the plant, establish the phytochemical constituents, and evaluate the effect of *Emilia sonchifolia* on haematological parameters and the histomorphology of the liver of Swiss albino male mice infected with *Plasmodium berghei berghei*. The result from the acute toxicity test indicated that 3250 mg/kg in mice, however 2874.02 mg/kg has been reported [5] obtained using similar method, the reason for this differences may be due to the season and the location in which both plants were cultivated and obtained for the different experiments. The preliminary phytochemical constituent in our study was similar to that reported by [5], although they did not report the degree of the bioavailability of these constituents in the leaves.

The result from the haematological indices in Table 3 indicates that in Group 5 treated with *E. sonchifolia* 650 mg/kg, showed RBC parameters that were significantly ( $P < 0.001$ ) reduced indicating haemolytic activity compared with the Group 1 (control). This may be due to the rich presence of saponins in the extract, see Table 1. Saponins promote hemolysis of RBC by increasing the water transport by the water channel aquaporin rather than by acting on the lipid phase [33]. It acts through structural changes in the membrane of RBC, by causing a decrease in the level cholesterol which thus affects the susceptibility of RBC membrane [34]. Interestingly, in Group 4 treated with *E. sonchifolia* 325mg/kg, there appeared to be erythropoietic effect compared to Group 1 (control), and similarly in Group 7 treated with (*E. sonchifolia* 650 mg/kg), which was parasitized before treatment. Reasoning that, perhaps the extract only promotes erythropoiesis at a low dose or in the presence of the berghei parasite, by a mechanism not well understood. However, flavonoid has been reported in vitro studies to have antidiarrheal and antioxidant activity [35], anti-allergic, anti-inflammatory [36], antimicrobial (antibacterial) [37-39], anti-cancer [40], antiviral [37, 39, 41], antifungal [37, 39]. Hence there appears to be an active interplay of both agonistic and synergistic effect from the extract. Alkaloid is also richly present in the extract and most plants contain several alkaloids. Their mixture is extracted first and then individual alkaloids are separated [41]. Many alkaloids are still used in medicine, usually in the form of salts, including the following: Quinine as antipyretics, antimalarial; Morphine as analgesic; Reserpine as antihypertensive; Codeine as cough medicine,

analgesic; Ergot alkaloids as sympathomimetic, vasodilator, antihypertensive; Caffeine as Stimulant, diuretic, Adenosine receptor antagonist just to mention but a few, with most of the known functions of alkaloids related to protection [42, 43].

The WBC shown in Table 3, indicates that Groups 2, 3, and 5 had significantly increased values ( $P < 0.01$ ) compared to Group 1 (control). Therefore it may imply that, at the dose 650 mg/kg, *E. sonchifolia* triggered sufficient lysing of RBC to activate the elevated levels of WBC present in circulation. WBCs are the mobile units of the body's protective system, and acting together, these cells provide the body with powerful defenses against tumors and viral, bacterial, and parasitic infections [44]. The changes in the Platelet values in the treatment groups compared with the control may reflect the animals' response at the start of bleeding from needle prick, to obtain thin blood smear for parasitemia (data not reported).

Ethanol extract of *Emilia sonchifolia* has been reported to cause splenotoxicity in a dose-dependent manner in mice [45]. The photomicrographs shows that the liver of the treated groups showed varying degree of adaptive responses which consisted of inflammation, hyperplasia, hypertrophy of the hepatocytes with reduced sinusoidal sizes, nuclei shrinkage and pyknotic nuclei especially in the parasitized groups. Group 2 with Figure 1B had poor staining intensity compared to the Group 1 with Figure 1A; this perhaps resulted from the severity of the parasite trauma to the parenchyma of the liver. This is closely observed in Group 3 parasitized and treated with Coartem®. In the non-parasitized Groups; 4 and 5 (Figures 1D and E) there was presence of few inflamed nuclei, but in the parasitized Groups 6 and 7 (Figures 1F and G), the parenchyma was severely affected. These changes may be indicative of an underlying cellular trauma and morphological change in the tissue cytoarchitecture a normal reaction of the liver tissue to insults [46].

In conclusion, oral administration of *Emilia sonchifolia* has a Median lethal dose of about 3250 mg/kg in mice. The extract is very rich in the presence of alkaloids as well as flavonoids and saponins, and promotes erythropoiesis at 325 mg/kg, but is haemolytic at 650 mg/kg, hence possess a dose-dependent negative effect, possibly through an agonistic and a synergistic activity of its rich bioactive ingredients. It showed mild toxic effect to the histomorphology of the non-parasitized mice at the low and medium doses, and also appeared to offer hepatoprotection in the parasitized mice compared to the group that was parasitized but without treatment.

## References

1. Yoga Latha L, Darah I, Sasidharan S, Jain K. Antimicrobial Activity of *Emilia sonchifolia* DC., *Tridax procumbens* L. and *Vernonia cinerea* L. of Asteraceae Family: Potential as Food Preservatives. *Malaysia Journal Nutrition*. 2009; 15(2):223-

- 231.
2. Essien GE, Nwido LL, Nwafor PA. Anti-Inflammatory and Analgesic Potential of Methanolic Extract of Emilia Sonchifolia (Compositae) Leaves in Rodents. *African Journal of Biomedical Research*. 2009; (12):3,199-207.
  3. Rahman A, Akter N, Rashid H, Ahmed NU, Uddin N, Islam S. Analgesic and anti-inflammatory effect of whole *Ageratum conyzoides* and *Emilia sonchifolia* alcoholic extracts in animal models. *African Journal of Pharmacy and Pharmacology*. 2012; 6(20):1469-1476.
  4. Muko, K.N. and Ohiri, F.C., A preliminary study on the anti-inflammatory properties of *Emilia sonchifolia* leaf extracts. *Fitoterapia*, (2000), 71 (1), 65-68.
  5. Shylesh, BS, Padikkala J. In vitro cytotoxic antitumour property of *E. Sonchifolia* (L) in mice. *Journal of Ethnopharmacology*. 2000; 73:495-500.
  6. Cibin TR, Srinivas G, Gayathri Devi D, Srinivas P, Lija Y, Abraham, A. Antioxidant and Antiproliferative Effects of Flavonoids from *Emilia sonchifolia* Linn on Human Cancer Cells. *International Journal of Pharmacology*. 2006; 2(5):520-524.
  7. Jiny Varghese K, Anila J, Nagalekshmi R, Resiya S, Sonu J. Dasapushpam: The Traditional Uses and The Therapeutic Potential of Ten Sacred Plants Of Kerala State In India. *International Journal of Pharmaceutical Sciences and Research*. 2010; 1(10): 50-59
  8. Shyura L, Tsunga J, Chenb J, Chiua C, Lo C. Antioxidant Properties of Extracts from Medicinal Plants Popularly Used in Taiwan. *International Journal of Applied Science and Engineering*. 2005;3(3):195-202.
  9. Guha G, Rajkumar V, Mathew L, Kumar RA. The antioxidant and DNA protection potential of Indian tribal medicinal plants. *Turkish Journal Biology*. 2011; (35):233-242.
  10. Sophia D, Ragavendran P, Arulraj C, Gopalakrishnan VK. In vitro antioxidant activity and HPTLC determination of n-hexane extract of *Emilia sonchifolia* (L.)DC. *Journal of Basic and Clinical Pharmacy*. 2011;002 (004):179-183.
  11. Raj M. Natural Antioxidant (Flavone Glycoside) From *Emilia Sonchifolia* Dc. and Its Potential Activity. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2012;4(1):159-162.
  12. Monago CC, Gozie GC, Joshua PE. Antidiabetic and Antilipidemic Effects of Alkaloidal Extract of *Emilia sonchifolia* in Rat. *Research Journal of Science and Technology*. 2010;51.
  13. Lija Y, Biju PG, Reeni A, Cibin TR, Sahasranamam V, Abraham A. Modulation of selenite cataract by the flavonoid fraction of *Emilia sonchifolia* in experimental animal model. *Phytotherapy Research*. 2006;20:1091-1095.
  14. Patel DK, Prasad SK, Kumar R, Hemalatha S. Cataract: A major secondary complication of diabetes, its epidemiology and an overview on major medicinal plants screened for anti-cataract activity. *Asian Pacific Journal of Tropical Disease*. 2011;323-329.
  15. Patel P, Jivani N, Malaviya S, Gohil T, Bhalodia Y. Cataract: A major secondary diabetic complication. *International Current Pharmaceutical Journal*. 2012;1(7):180-185.
  16. Singh B, Kumar D, Singh R. Phytotherapeutics for management and prevention of cataract generation. *Phytopharmacology*. 2012;3(1)93-110.
  17. Asije O, Adelusi SA, Usifoh CO. Anticonvulsant Activity of *Emilia sonchifolia* Leaf Extracts. *Pakistan Journal of Scientific and Industrial Research*. 2006;49(4):269.
  18. Couto VM, Vilela FC, Dias DF, Dos Santos MH, Soncini R, Nascimento CG, Giusti-Paiva A (2011). Antinociceptive effect of extract of *Emilia sonchifolia* in mice. *J. Ethnopharmacol.*, 134(2): 348-353.
  19. Gao JJ, Cheng DL, Liu XP (1993). Chemical constituents of *Emilia sonchifolia* L. DC. *Zhongguo Zhong Yao Za Zhi*, 18(2):102-3, 127.
  20. Cheng D, Röder E (1986). Pyrrolizidine Alkaloids from *Emilia sonchifolia*. *Planta Med.*, (6): 484- 486.
  21. Ogbonnia SO, Mbaka GO, Anyika EN, Emordi JE, Nwakakwa N. An Evaluation of Acute and Subchronic Toxicities of a Nigerian Polyherbal Tea Remedy. *Pakistan Journal of Nutrition*. 2011; 10(11): 1022-1028
  22. Tedong LP, Dzuefiet DD, Dimo J, Asongalem EA, Sokeng SN, Flejou JI, Callard P, Kamtchouring P. Acute and subchronic Toxicity of *Anacardium occidentale* Leaves Extract in Mice. *African Journal of Traditional Alternative Medicine*. 2007;(4)140-147
  23. Taheri S, Zarei A, Ashtiyani SC, Rezaei A, Zaheiri S. Evaluation of the Effects of Hydroalcoholic extract of *Berberis vulgaris* rot on the Activity of Liver enzymes in Male Hypercholesterolaemic rats. *Journal of Phytomedicine*. 2012;2(3):153-161
  24. American Physiological Society. (2002) Guiding principles for research involving animals and human beings. *Am. J Physiol. Regul. Integr. Comp. Physiol.* 283:281-283.
  25. Odetola A, Basir O. (1980) Evaluation of antimalarial properties of some Nigerian Medicinal Plants. In: Sofowora A, Editor. *Proceeding of African Bioscience Network, Federal Ministry of Science and Technology, Nigerian Society of Pharmacology and Drug Research and Production unit, University of Ife organized Workshop, Ife.* 275-283.
  26. Lorke D. (1983) A New Approach to practical Acute Toxicity Testing. *Arch Toxicol.* 54:275-287.
  27. Trease GE, Evans WC. (1989) *A Textbook of Pharmacognosy*. (13th ed). Bailliere Tinnall Ltd, London.
  28. Conn HJ. (1946). *Biological Stains: A Handbook on the Nature and Uses of the Dyes Employed in the Biological Laboratory*. 5th edition, Geneva. Biotech publication, pp 24-29.
  29. Winter WP. (1994) Mechanism of saponin induced red cell hemolysis: Evidence for the involvement of aquaporin CHIP28. *Blood* 84: Suppl. 1 to 10, Abstr. 445.
  30. Segal R, Milo-Goldstein I. (1978) The susceptibility of cholesterol-depleted erythrocytes to saponin and sapogenin hemolysis. *Biochim Biophys Acta.* 512: 223-226.
  31. Bagchi M, Milnes M, Williams C, Balmoori J, Ye X, Stohs S, Bagchi D. (1999) Acute and chronic stress-induced oxidative gastrointestinal injury in rats, and the protective ability of a novel grape seed proanthocyanidin extract. *Nutrition Research*. 19 (8): 1189-1199.
  32. Yamamoto Y, Gaynor RB. (2001) Therapeutic potential of inhibition of the NF-kB pathway in the treatment of inflammation and cancer. *Journal of Clinical investigation* 107 (2): 135-42.

33. Cushnie TPT, Lamb AJ. (2005) Antimicrobial activity of flavonoids. *International Journal of Antimicrobial Agents*. 26 (5): 343-356.
34. Cushnie TPT, Lamb AJ. (2011) Recent advances in understanding the antibacterial properties of flavonoids. *International Journal of Antimicrobial Agents*. 38 (2): 99-107.
35. Friedman M. (2007) Overview of antibacterial, antitoxin, antiviral, and antifungal activities of tea flavonoids and tea. *Molecular Nutrition and Food research*. 51 (1):116-134.
36. de Sousa RR, Queiroz KC, Souza AC, Gurgueira SA, Augusto AC, Miranda MA, Peppelenbosch MP, Ferreira CV, Aoyama H. (2007) Phosphoprotein levels, MAPK activities and NF $\kappa$ B expression are affected by fisetin. *Journal of Enzyme Inhib Med Chem*. 22 (4): 439 – 444.
37. Grinkevich NI, Safronich LN (1983). *The Chemical Analysis of Medical Plants: Proc. Allowance for Pharmaceutical Universities*. p.132
38. Meyers RA. (2002). *Encyclopedia of Physical Science and Technology – Alkaloids*, 3rd Edition. ISBN 0-12-227411-3
39. Hesse M. (2002). *Alkaloids: Nature's Curse or Blessing?* Wiley-VCH ISBN 978-3-906390-24-6
40. Ganong WF. (2003) *Review of medical physiology* (21 ed.). New York: Lange Medical Books/McGraw-Hill. p. 518.
41. Edagha IA, Ekandem GJ, Ekanemesang UM, Ekanem TB, Isinam BF. (2013) Histopathological effect of *Nauclea latifolia* and *Emilia sonchifolia* on the spleen of Swiss mice infected with *Plasmodium berghei berghei*. *Clinical Anatomy* 26:913.
42. Kumar V, Abbas AK, Fausto N. (2005) *Robbins and Cotran: Pathologic Basis of Disease*, 7th edition. Elsevier Saunders. p 2-45.