



Parents' Acceptance to Alveolar and Nasoalveolar Molding Appliances during Early Cleft Lip and Palate Care: A Call for High-Quality Research

Mohamed Abd El-Ghafour, Sherif A. Elkordy, Mona M. Salah Fayed, Amr Ragab El-Beialy, Faten Hussein Kamel Eid

Department of Orthodontics, Faculty of Dentistry, Cairo University, Cairo, Egypt

Abstract

Edited by: Sasho Stoleski
Citation: Abd El-Ghafour M, Elkordy SA, Fayed MMS, El-Beialy AR, Eid FTK. Parents' Acceptance to Alveolar and Nasoalveolar Molding Appliances during Early Cleft Lip and Palate Care: A Call for High-Quality Research. Open Access Maced J Med Sci. 2020 May 09; 8(A):58-64. https://doi.org/10.3889/oamjms.2020.3856
Keywords: Cleft Lip and Palate; Nasoalveolar Molding; Parents' Acceptance; Quality of Life
***Correspondence:** Mohamed Abd El-Ghafour, Department of Orthodontics, Faculty of Dentistry, Cairo University, Cairo, Egypt.
E-mail: m.abdelghafour@dentistry.cu.edu.eg
Received: 07-Oct-2019
Revised: 11-Feb-2020
Accepted: 08-Apr-2020
Copyright: © 2020 Mohamed Abd El-Ghafour, Sherif A. Elkordy, Mona M. Salah Fayed, Amr Ragab El-Beialy, Faten Hussein Kamel Eid
Funding: This research did not receive any financial support
Competing Interests: The authors have declared that no competing interests exist
Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

AIM: Acceptance and compliance of the parents are an essential pillar in the success of pre-surgical infant orthopedic (PSIO) treatment. The aim of this systematic review is to evaluate the burden of care associated with the alveolar molding (AM) and nasoalveolar molding (NAM) appliances as experienced by the parents with unilateral complete cleft lip and palate (UCLP) infants.

METHODS: An electronic search was carried on by two reviewers in eight search engines, as well as a manual search till July 2019. Randomized controlled trials (RCTs) comparing AM/NAM appliances to controls in infants with UCLP were selected. Risk of bias was evaluated using Cochrane risk of bias assessment tool for RCTs.

RESULTS: One RCT was included in the qualitative analysis. Non-significant differences were found in the amount of mothers' satisfaction between the intervention and control groups.

CONCLUSIONS: Insufficient low-quality evidence is available regarding the effects of AM and NAM on parents' satisfaction and burden of care. No conclusions can be withdrawn from the existing studies. High-quality research is needed to elucidate the degree of parents' acceptance to the molding appliances. PROSPERO registration number: CRD42016043174.

Introduction

Rationale

Cleft lip and palate (CLP) is considered the most common craniofacial anomaly in different populations [1]. Management of patients suffering from CLP can start early at infancy [2] with treatment goals of lip segments approximation, nostrils symmetry achievement, increase columellar length, and alveolar segments alignment [3]. In other words, the aim at infancy is to help the surgeons to achieve better surgical result by decreasing the severity of the cleft defect [4]. It was assumed that pre-surgical infant orthopedics (PSIO) might help in achieving these goals.

PSIO appliances were introduced in the orthodontic literature, including passive plates [5], active plates [6], pin-retained Latham's appliance [7], and nasoalveolar molding (NAM) [3]. In 1978, Hotz et al. [8] used an intraoral plate to mold the maxillary segments before the surgical lip repair, this was only by the concept of "alveolar molding" (AM). Several years after, Grayson et al. [3] introduced the NAM concept in 1993. A nasal stent was added the intraoral plate to

mold the nasal cartilage into a normal form [3] taking a further step toward achieving the PSIO goals. Both Hotz's and Grayson's intraoral plates were activated by addition of soft acrylic on the fitting surface at the pressure areas and grinding at the relief areas.

From all the steps of CLP management protocol, the step of PSIO is considered as the most doubtful. The researchers' recommendations of PSIO usage were swinging between strong promoters [9], [10], [11], [12] and heavy opponents [5], [13], [14], [15], [16]. According to the latest systematic reviews [17], [18], [19], [20], NAM was considered as the most effective type of PSIO, depending on the available low-quality evidence.

Because of the handicapping nature of the infants, the success of PSIO depends mainly on the parents' positive interaction and commitment. Several studies [16], [21], [22], [23], [24], [25], [26], [27] evaluated the parents' satisfaction while dealing with the PSIO. They measured the amount of load on the parents, while carrying out the procedures of taking care of their child in this period. Some of those studies [25], [26] found that there is an increased burden on the parents, while others [16], [24] did not support this finding. In addition, some authors [21], [23] found that the parents were

willing to withstand this added load to help their children; however, no conclusive results could be withdrawn.

None of the previously published systematic reviews [17], [18], [19], [20] evaluated the amount of burden and satisfaction experienced by the parents during handling of the NAM appliance. Moreover, two of the found systematic reviews [17], [18], [19], [20] did not specify the type of the PSIO included in their evaluation.

The current systematic review focused on the degree of parents' acceptance to the AM and NAM alone. Evaluation of parents' satisfaction and avoidance of previous methodological problems were planned to find an answer about the ability of the parent to deal with these appliances.

Objectives

The aim of this systematic review is to answer the following question; in infants with unilateral CLP; what are the responses of the parents with the usage of AM and NAM appliances, in terms of their ability to handle the appliances and the burden of care in comparison to the untreated controls?

The PICOS format is: Population: Parents with unilateral CLP infants, Intervention: AM and NAM appliances, Comparator: Untreated control (only surgical lip repair), Outcomes: Parents' satisfaction, Study designs included: Randomized controlled trials.

Materials and Methods

Protocol registration

The presented systematic review was performed following a preset protocol following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement [27]. The review protocol was registered at the PROSPERO (International prospective register of systematic reviews) with registration number: CRD42016043174.

Information sources, search strategy, and study selection

Electronic search was done in eight databases till July 2019. These eight databases included PubMed,

Cochrane library central, Wiley online library, LILACS, ScienceDirect, Web of Science, Scopus, and Ovid. The used keywords are mentioned in Table 1. Manual search was done to all the online available issues till July 2019 in Cleft Palate-Craniofacial Journal, Plastic and Reconstructive Surgery Journal, Journal of Maxillofacial and Oral Surgery, American Journal of Orthodontics and Dentofacial Orthopedics and Angle Orthodontist. The unpublished literature was searched in the libraries of national universities. Two reviewers did the search separately (M.A. and S.K.).

Each reviewer screened the titles and abstracts for all the found studies after duplicates removal using Mendeley Desktop software (version 1.13.8). Next, each reviewer read the full text of the previously selected studies to select the included ones. Any incongruity between reviewers was solved by a discussion with the third reviewer (M.F.) to reach an agreement.

Eligibility criteria

The inclusion and exclusion criteria of the studies to be included are mentioned in Table 2. The important eligibility criteria were; the presence of control group, prospective nature, unilateral cases, infants age at start of the treatment <30 days, and studies written only in English language.

Data items and collection process

Data extraction sheets were constructed (Table 3) to the included study. The tables were divided into two categories; first; including study design, study settings, total number of patients, the used alveolar or NAM design, follow-up period, types of records collected, and time of lip repair; second the results table containing the results and conclusions.

Risk of bias/quality assessment in individual studies

The Cochrane risk of bias assessment tool [28] was used. The Cochrane tool encompasses seven items; the first two items are random sequence generation and allocation concealment used for detection of selection bias. The third item is blinding of participants and researchers for performance bias assessment. Blinding of outcome assessment is the fourth item to rule out detection bias. The fifth item is incomplete outcome data

Table 1: The electronic databases searched, the search strategies used, and the corresponding results

| Electronic database | Date | Search strategy used | Results |
|----------------------|-----------|--|---------|
| PubMed | July 2019 | (Cleft lip and palate OR cleft lip OR cleft palate OR unilateral OR cleft alveolus | 860 |
| Cochrane library | July 2019 | OR cleft gap OR alveolar notch OR alveolar cleft) AND (Nasoalveolar molding | 107 |
| LILACS | July 2019 | OR nasoalveolar molding OR NAM OR Pre-surgical Nasoalveolar molding OR | 51 |
| Scopus | July 2019 | nasoalveolar molding OR PNAM OR Pre-surgical appliance OR Pre-surgical | 175 |
| Wiley online Library | July 2019 | device OR Pre-surgical orthopedics OR Pre-surgical appliance OR Nasal stents | 23 |
| Web of Science | July 2019 | OR preoperative orthopedics OR alveolar molding OR alveolar molding OR | 429 |
| Ovid | July 2019 | nasal alveolar molding OR nasal alveolar molding) | 8 |
| ScienceDirect | July 2019 | | 237 |

to detect attrition bias, while the sixth item is selective reporting for the reporting bias recognition. Finally, the seventh item is to evaluate any other sources of bias found by the reviewers; mostly the absence of sample size calculation was considered in this item. Three decisions to be made; either low, unclear, or high risk of bias and if a study receives a single unclear or high risk of bias the whole study takes the same evaluation.

Summary measures and synthesis of results

According to the Cochrane Handbook for systematic Reviews of interventions [29], the possible heterogeneity between the included studies was assessed in its three forms; clinical, methodological, and statistical heterogeneity. Clinical heterogeneity was assessed by comparing the demographic data of the included studies. Methodological heterogeneity was evaluated by appraisal of the followed AM/NAM protocol. In the current review, statistical heterogeneity was not assessed due to the inability to perform a meta-analysis.

Table 2: Inclusion and exclusion criteria

| Category | Inclusion criteria | Exclusion criteria |
|----------------------|--|---|
| Participants | a. Patients younger than 1 month at start of treatment. b. Infants with unilateral complete cleft lip and palate. | a. Patients older than 30 days at start of treatment. b. Infants with unilateral incomplete cleft lip and palate. c. Bilateral cleft lip and palate cases. |
| Intervention | Any technique of nasoalveolar molding | All other pre-surgical infant orthopedic devices including reversed expansion screws, pins (Latham appliance). |
| Comparator | Presence of no molding control group. | Absence of no molding control group. |
| Outcomes | Parents' burden of care, including; their ability to handle the appliance and their psychological status during NAM treatment. | Any other outcomes. |
| Study design | a. Randomized controlled trials (RCTs). b. Quasi randomized controlled trials (quasi-RCTs). | a. Prospective controlled clinical trials (CCTs). b. Retrospective studies. c. Case reports. d. Case series. e. Expert opinion. f. Letters to the editor. g. Systematic reviews. h. Narrative reviews. |
| Language restriction | Only studies written in English language. | Studies written in other languages rather than the English language. |

Table 3: Data extraction sheet of the included study

| Study | Study Design and setting | Total number of patients | Patients' age and Characteristics | NAM Technique | AM or NAM (Presence of nasal stent) | Follow-up period | Types of Records | Pre-surgical records | Lip closure | Post-surgical records | Outcome |
|--------------------------|--|---|---|------------------------------|-------------------------------------|------------------|------------------|----------------------|-------------|------------------------------|-----------------------|
| 1 Prael et al. [16] 2008 | RCT/ In three participating academic cleft palate centers in the Netherlands: Nijmegen, Amsterdam and Rotterdam. | 48 | 2 weeks/ Complete UCLP, infants born at term, both parents Caucasian and fluent in the Dutch language, and trial entrance within 2 weeks after birth. | Combined Hard and soft plate | No | 12 months | Questionnaire | T1': at 6 wk. | 18 weeks | T3: at 25 wk. T4': at 58 wk. | Parents' Satisfaction |
| Study | Results | | Conclusions | | | | | | | | |
| 1 Prael et al. [16] 2008 | I | C | Results from this study show that infant orthopedics, with a passive plate during the first year of life, in children with a unilateral cleft lip and palate has no influence on the mothers' satisfaction in motherhood. | | | | | | | | |
| | T1'' 1.60 (0.29) T3 1.48 (0.24) T4' 1.53 (0.24) | T1'' 1.50 (0.22) T3 1.45 (0.19) T4' 1.43 (0.20) | | | | | | | | | |

T1: Before molding, T1': At 3 weeks, T1'': at 6 weeks, T2: After molding and directly before Lip repair, T3: Directly after lip repair, T4: At 1 year, T4': At 58 weeks, T5: At 1.5 year, T6: At 6 years.

Time points were standardized between the studies, where; T1: Before molding, T1': At 3 weeks, T1'': at 6 weeks, T2: After molding and directly before lip repair, T3: Directly after lip repair, T4: At 1 year, T4': At 58 weeks, T5: At 1.5 year, T6: At 6 years.

Risk of bias across studies

For assessment of publication bias, standard funnel plots and contoured enhanced funnel plots were planned to be used, only when more than ten studies included in the meta-analysis [29].

The Grades of Recommendation, Assessment, Development, and Evaluation (GRADE) [30], [31] was the tool used to assess the overall quality of evidence for each of the main outcomes. Evaluation of the methodological quality of the studies, the directness of evidence, the inconsistency, the precision of effect estimates, and the risk of publication bias were assessed using the GRADE profiler. The certainty of evidence was interpreted in four categories; very low quality, very uncertain about the effect's estimate. Low quality, where more research is very likely to have an important impact on the confidence in the estimate of effect and is likely to change the estimate. Moderate quality, further research will have an important impact on the confidence in the estimate and may change it. High quality, the confidence in the estimate is high and new research is very unlikely to change this estimate.

Additional analysis

No additional analyses were performed in the presented systematic review.

Results

Study selection and characteristics

The electronic search resulted in 1917, while the manual search produced 27 studies (Figure 1). After




Figure 1: PRISMA flowchart of article retrieval

duplicates removal using Mendeley Desktop software (version 1.13.8), 1183 articles were subjected to screening by title and abstract. After 1178 exclusions, five studies were read in full text. As a result of full text screening, four articles [21], [22], [23], [24] were excluded with reasons (Table 4) and one study [16] met the inclusion criteria and were included in the qualitative analysis.

The included study [16] was measuring the mother’s satisfaction while dealing with the Hotz molding plate.

Risk of bias within studies

The Cochrane risk of bias assessment tool [28] was used for the included RCT [16] (Table 5). Seven

Table 4: Excluded papers with reasons

| Article | Reason for exclusion |
|----------------------------|--|
| Sischo et al. [21] 2015 | No control group |
| Sischo et al. [22] 2016 | Mixed unilateral and bilateral CLP |
| Broder et al. [23] 2016 | Mixed unilateral and bilateral CLP |
| Hopkin et al. [24] in 2016 | No control group and retrospective study |

CLP: Cleft lip and palate

criteria were evaluated for the included RCT. For the random sequence generation, blinding of outcome assessment, incomplete outcome data and selective reporting, the included RCT was with low risk of bias. For the allocation concealment, blinding of outcome assessment and other risks, it recorded unclear risk of bias. The overall risk of bias of the included RCT was reached to be unclear risk of bias.

Results of individual studies, meta-analysis, and additional analyses

The parents' satisfaction was reported in only one RCT [16] and they found no difference between molding and no molding groups. The range of the mean scores for the individual items on the questionnaires for both groups ranged between 1.1 and 2.4. Mothers appear to be satisfied in motherhood, least satisfied with the available time for themselves, and very satisfied with hugging and walking their babies. No differences were found between groups. In the current systematic review, no meta-analyses were performed.

Risk of bias across studies

The GRADE approach for rated the available evidence as low quality for the assessed outcome (Table 6). Low quality evidence was found for parents' satisfaction at T2 and T3.

Discussion

Controversies exist regarding the inclusion of PSIO in the followed CLP treatment protocols in the cleft centers around the world [2]. Management of infants in their 1st weeks in life is very difficult on both the parents and healthcare providers. Since that the parents are the main performers in the NAM treatment, the aim of this systematic review are to evaluate their responses with such a treatment.

A preplanned inclusion criteria were set. Studies discussed both AM and NAM appliances were included, as alveolar molding is a common step in both types. To increase the validity of this systematic review's results, only RCTs were included. An important

inclusion criterion was strictly followed, was the presence of control group for comparison.

Summary of evidence

After the systematic search, only one RCT [16] was found fitting into the inclusion criteria. This RCT made on the mothers of 48 infants with UCLP divided into 24 using Hotz plate versus 24 controls. In this study, nasal stents were not used.

In the solely found RCT [16], no differences were found between groups regarding the mother's response to her child with and without molding. The presence of one study was insufficient to conclude the real effect of the appliances' usage on the parents especially that, this single article did not discuss the burden of appliance posed itself, the parents' stress and anxiety of the parents nor the effect on the fathers.

The four excluded studies [21], [22], [23], [24] evaluated the caregiver responses to the NAM appliance treatment phase. Unfortunately, these studies mixed the unilateral and bilateral cleft infants, in addition to the lack of the control group in two studies [21], [24] and that is why they were excluded from the current systematic review.

Despite of the indifferent results between the treated and the control groups in the included study [16], the excluded studies [21], [22], [23], [24] reached different conclusions. The excluded studies found a positive impact on the parents with children treated with NAM appliances. They concluded that completing the NAM treatment was often associated with positive factors such as increased empowerment, self-esteem, and bonding with their child [21], [22], in addition to more acceptable esthetic outcomes [23].

Surprisingly, none of available articles evaluated the ability of the parents to handle the

Table 5: Assessment of risk of bias for the included RCT using Cochrane risk of bias assessment tool

| Study | Random sequence generation | Allocation concealment | Blinding of participants and researchers | Blinding of outcome assessment | Incomplete outcome data | Selective reporting | Other bias | Overall risk of bias |
|------------------------|----------------------------|------------------------|--|--------------------------------|-------------------------|---------------------|------------|----------------------|
| Prahl et al. [16] 2008 | Low | Unclear | Low | Unclear | Low | Low | Unclear | Unclear |

Table 6: GRADE summary of findings table for the main outcomes of the systematic review

| Outcomes | Anticipated absolute effects ^a (95% CI) Risk with Alveolar and Nasoalveolar Molding | No of participants (studies) | Certainty of the evidence (GRADE) | Comments |
|---|--|------------------------------|-----------------------------------|---|
| Parent Satisfaction (Parent Satisfaction) assessed with: Questionnaire Scale from: 1 to 4 follow-up: up to 58 weeks | The mean parent Satisfaction in the intervention group was 0.03 – higher (0.09 lower to 0.15 higher) | 48 (1 RCT) | ⊕⊕○○ LOW ^{a,b} | Only one study found on parents' satisfaction outcome discussing only the mother's satisfaction and found no difference between the groups. |

Explanations

^aUnclear risk of bias in allocation concealment. ^bOnly 48 patients included. ^cOne study has high risk of bias. ^dI²>75%. ^eVery wide confidence interval. ^fConfidence interval includes no effect. ^gVery wide confidence interval and including no effect. ^hNon-randomized study. ⁱOnly 28 patients were included.

NAM appliances nor measuring the degree of their acceptance to the hectic appliance. All the found studies [16], [21], [22], [23], [24] evaluated the psychological side only of the parents, missing a former step of evaluating the parents' acceptance to the appliance itself at the very beginning.

Limitations of the available evidence

The included RCTs were of low quality and with unclear risk of bias. No meta-analyses were performed due to the scarcely data. It seems that the degree of parents' satisfaction was not in the interest of most of the researchers.

Generalizability and applicability

A knowledge gap still exists regarding the ability of the parents to handle the appliance. More high-quality studies are strongly recommended to explore this vague point.

This systematic review spots the light on a missed outcome in the literature, which is parents' satisfaction while dealing with AM and NAM appliances. This article will motivate the researchers to assess an unexplored outcome.

Conclusions

Based on the available low-quality evidence, no conclusions can be withdrawn for the effect of AM or NAM on parents' burden of care and NAM appliance acceptance. Insufficient evidence is currently available regarding the effects of AM and NAM on infants with UCLP, especially for parents' satisfaction.

Implications for research

- 1- Further well-designed high quality RCTs are needed to evaluate parents' satisfaction with and without NAM.
- 2- Post-surgical records should be available to determine the effect on the parents after lip repair.
- 3- The presence of control group is strongly advisable to reach true results about the burden of care.
- 4- Standardization of evaluation time points and the used questionnaires is important factors to allow for future pooling of data for the unevaluated outcomes.

References

1. Mossey PA, Modell B. Epidemiology of oral clefts 2012: An international perspective. *Front Oral Biol.* 2012;16:1-8. PMID:22759666
2. Colbert SD, Green B, Brennan PA, Mercer N. Contemporary management of cleft lip and palate in the United Kingdom. Have we reached the turning point? *Br J Oral Maxillofac Surg.* 2015;53(7):594-8. <https://doi.org/10.1016/j.bjoms.2015.06.010> PMID:26130590
3. Grayson BH, Cutting C, Wood R. Preoperative columella lengthening in bilateral cleft lip and palate. *Plast Reconstr Surg.* 1993;92(7):1422-3. PMID:8248436
4. Rubin MS, Clouston S, Ahmed MM, M Lowe K, Shetye PR, Broder HL, et al. Assessment of presurgical clefts and predicted surgical outcome in patients treated with and without nasoalveolar molding. *J Craniofac Surg.* 2015;26(1):71-5. <https://doi.org/10.1097/scs.0000000000001233> PMID:25534051
5. Prah C, Kuijpers-Jagtman AM, van't Hof MA, Prah-Andersen B. A randomised prospective clinical trial into the effect of infant orthopaedics on maxillary arch dimensions in unilateral cleft lip and palate (Dutchcleft). *Eur J Oral Sci.* 2001;109(5):297-305. <https://doi.org/10.1034/j.1600-0722.2001.00056> PMID:11695749
6. McNEIL CK. Orthodontic procedures in the treatment of congenital cleft palate. *Dent Rec (London).* 1950;70(5):126-32. PMID:24537837
7. Latham RA, Kusy RP, Georgiade NG. An extraorally activated expansion appliance for cleft palate infants. *Cleft Palate J.* 1976;13:253-61. <https://doi.org/10.1097/00006534-197803000-00085> PMID:780004
8. Hotz MM, Gnoinski WM, Nussbaumer H, Kistler E. Early maxillary orthopedics in CLP cases: Guidelines for surgery. *Cleft Palate J.* 1978;15(4):405-11. PMID:281285
9. El-Haddad A. Evaluation of Presurgical Oral Orthopedic Appliances in Early Management of Unilateral Cleft Lip and Palate. PhD Thesis, Cairo University; 2005. p. 1-160.
10. Adachi K, Togashi S, Yanagawa T, Ishibashi N, Goto T, Yamagata K, et al. Presurgical orthopedic treatment ameliorates postoperative nasal deformity after cheiloplasty. *Ann Plast Surg.* 2013;71(2):170-5. <https://doi.org/10.1097/sap.0b013e318246814e> PMID:23123612
11. Karube R, Sasaki H, Togashi S, Yanagawa T, Nakane S, Ishibashi N, et al. A novel method for evaluating postsurgical results of unilateral cleft lip and palate with the use of Hausdorff distance: Presurgical orthopedic treatment improves nasal symmetry after primary cheiloplasty. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2012;114(6):704-11. <https://doi.org/10.1016/j.oooo.2012.01.042> PMID:22906581
12. Shetty V, Agrawal RK, Sailer HF. Long-term effect of presurgical nasoalveolar molding on growth of maxillary arch in unilateral cleft lip and palate: Randomized controlled trial. *Int J Oral Maxillofac Surg.* 2017;46(8):977-87. <https://doi.org/10.1016/j.ijom.2017.03.006> PMID:28416097

13. PrahI C, Kuijpers-Jagtman AM, Van 't Hof MA, PrahI-Andersen B. A randomized prospective clinical trial of the effect of infant orthopedics in unilateral cleft lip and palate: Prevention of collapse of the alveolar segments (Dutchcleft). *Cleft Palate Craniofac J.* 2003;40(4):337-42. [https://doi.org/10.1597/1545-1569\(2003\)040<0337:arpcto>2.0.co;2](https://doi.org/10.1597/1545-1569(2003)040<0337:arpcto>2.0.co;2)
PMid:12846598
14. PrahI C, Kuijpers-Jagtman AM, Van 't Hof MA, PrahI-Andersen B. Infant orthopedics in UCLP: Effect on feeding, weight, and length: A randomized clinical trial (Dutchcleft). *Cleft Palate Craniofac J.* 2005;42(2):171-7. <https://doi.org/10.1597/03-111.1>
PMid:15748108
15. PrahI C, PrahI-Andersen B, van 't Hof MA, Kuijpers-Jagtman AM. Infant orthopedics and facial appearance: A randomized clinical trial (Dutchcleft). *Cleft Palate Craniofac J.* 2006;43(6):659-64. <https://doi.org/10.1597/05-139>
PMid:17105328
16. PrahI C, PrahI-Andersen B, Van't Hof MA, Kuijpers-Jagtman AM. Presurgical orthopedics and satisfaction in motherhood: A randomized clinical trial (Dutchcleft). *Cleft Palate Craniofac J.* 2008;45(3):284-8. <https://doi.org/10.1597/07-045.1>
PMid:18452361
17. Papadopoulos MA, Koumridou EN, Vakalis ML, Papageorgiou SN. Effectiveness of pre-surgical infant orthopedic treatment for cleft lip and palate patients: A systematic review and meta-analysis. *Orthod Craniofac Res.* 2012;15(4):207-36. <https://doi.org/10.1111/j.1601-6343.2012.01552.x>
PMid:23020693
18. Abbott MM, Meara JG. Nasoalveolar molding in cleft care: Is it efficacious? *Plast Reconstr Surg.* 2012;130(3):659-66.
PMid:22929251
19. van der Heijden P, Dijkstra PU, Stellingsma C, van der Laan BF, Korsten-Meijer AG, Goorhuis-Brouwer SM. Limited evidence for the effect of presurgical nasoalveolar molding in unilateral cleft on nasal symmetry: a call for unified research. *Plast Reconstr Surg.* 2013;131(1):62e-71e. <https://doi.org/10.1097/prs.0b013e318267d4a5>
PMid:23271555
20. Uzel A, Alparslan ZN. Long-term effects of presurgical infant orthopedics in patients with cleft lip and palate: A systematic review. *Cleft Palate Craniofac J.* 2011;48(5):587-95. <https://doi.org/10.1597/10-008>
PMid:20858135
21. Sischo L, Broder HL, Phillips C. Coping with cleft: A conceptual framework of caregiver responses to nasoalveolar molding. *Cleft Palate Craniofac J.* 2015;52(6):640-50. <https://doi.org/10.1597/14-113>
PMid:25225840
22. Sischo L, Clouston SA, Phillips C, Broder HL. Caregiver responses to early cleft palate care: A mixed method approach. *Health Psychol.* 2016;35(5):474-82. <https://doi.org/10.1037/hea0000262>
PMid:26280177
23. Broder HL, Flores RL, Clouston S, Kirschner RE, Garfinkle JS, Sischo L, et al. Surgeon's and Caregivers' Appraisals of Primary Cleft Lip Treatment with and without Nasoalveolar Molding: A Prospective Multicenter Pilot Study. *Plast Reconstr Surg.* 2016;137(3):938-45. <https://doi.org/10.1097/01.prs.0000479979.83169.57>
PMid:26910677
24. Hopkins EE, Gazza E, Marazita ML. Parental experience caring for cleft lip and palate infants with nasoalveolar moulding. *J Adv Nurs.* 2016;72(10):2413-22. <https://doi.org/10.1111/jan.12994>
PMid:27144651
25. Chen YF, Liao YF. A modified nasoalveolar molding technique for correction of unilateral cleft nose deformity. *J Craniomaxillofac Surg.* 2015;43(10):2100-5. <https://doi.org/10.1016/j.jcms.2015.10.003>
PMid:26541749
26. Rau A, Ritschl LM, Mücke T, Wolff KD, Loeffelbein DJ. Nasoalveolar molding in cleft care—experience in 40 patients from a single centre in Germany. *PLoS One.* 2015;10(3):e0118103. <https://doi.org/10.1371/journal.pone.0118103>
PMid:25734535
27. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *PLoS Med.* 2009;6(7):e1000100. <https://doi.org/10.1371/journal.pmed.1000100>
PMid:19621070
28. Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ.* 2011;343:d5928. <https://doi.org/10.1136/bmj.d5928>
PMid:22008217
29. Higgins J, Green S. *Cochrane Handbook for Systematic Reviews of Interventions.* Oxford: The Cochrane Collaboration and John Wiley and Sons Ltd.; 2006.
30. Guyatt GH, Oxman AD, Schünemann HJ, Tugwell P, Knottnerus A. GRADE guidelines: A new series of articles in the *Journal of Clinical Epidemiology.* *J Clin Epidemiol.* 2011;64(4):380-2. <https://doi.org/10.1016/j.jclinepi.2010.09.011>
PMid:21185693
31. Schünemann H, Brožek J, Guyatt G, Andrew O. *GRADE Handbook;* 2013. Available from: <https://www.gdt.gradepro.org/app/handbook/handbook.html>.