



Evaluation of Body Mass Index of 3–10-year-old Children with Adenotonsillar Hypertrophy, who Referred to Khatamolambia Hospital during 2017–2018

Gholam-Ali Dashti Khavidaki, Reza Gharibi*

Department of ENT Disorder, AL-Zahra Hospital, Zahedan University of Medical Sciences, Zahedan, Iran

Abstract

Edited by: Sasho Stoleski
Citation: Khavidaki GD, Gharibi R. Evaluation of Body Mass Index (BMI) of 3–10-year-old Children with Adenotonsillar Hypertrophy, who Referred to Khatamolambia Hospital during 2017–2018. Open Access Maced J Med Sci. 2020 Jun 17; 8(B):324-328. https://doi.org/10.3889/oamjms.2020.4393
Keywords: Body mass index; 3–10-year-old children; Adenotonsillar hypertrophy
***Correspondence:** Reza Gharibi, Department of ENT Disorder, AL-Zahra Hospital, Zahedan University of Medical Sciences, Zahedan, Iran. Tel.: +98- 543 322 4991. E-mail: Hessam.gharibi@gmail.com
Received: 04-Feb-2020
Revised: 17-May-2020
Accepted: 21-Apr-2020
Copyright: © 2020 Gholam-Ali Dashti Khavidaki, Reza Gharibi
Funding: This research did not receive any financial support.
Competing Interests: The authors have declared that no competing interests exist.
Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

BACKGROUND: Adenotonsillar hypertrophy is a common disorder in the children's population that can lead to growth disorders. Chronic adenotonsillar hypertrophy can cause a break in the normal growth of children, weight loss, and decreased growth hormone secretion.

AIM: The purpose of this study was to evaluate adenotonsillar hypertrophy in the growth rate of children with height, weight, and body mass index (BMI).

METHODS: In this descriptive-analytical cross-sectional study, after examining 312, 3–10-year-old children with adenotonsillar hypertrophy, who met the inclusion criteria, growth scales were evaluated and entered into the patient evaluation form.

RESULTS: There was a significant relationship between BMI, height, weight, and severity of tonsillar hypertrophy at Brodsky's scale ($P < 0.001$). Our studies showed well that with increasing severity of adenotonsillar hypertrophy disease, the growth indicators are also decreased in children.

CONCLUSION: Adenotonsillar hypertrophy decreases growth indicators in children. There was also a relationship between growth indicators and severity of adenotonsillar hypertrophy in patients.

Introduction

Adenotonsillar hypertrophy is a common disorder in the children population that causes an obstruction in the nasal route and eustachian tube and prevents drainage of nasal mucus [1]. The common clinical signs of adenotonsillar hypertrophy include nasal congestion, which results in sleep disorders from snoring to obstructive sleep apnea, rhinorrhea, oral respiration, hyponasal speech, and cough [2].

Obstructive sleep apnea, as one of the most damaging disorders of adenotonsillar hypertrophy, leads to frequent hypoxia and sleep disturbances, which can have adverse effects on the cognitive and behavioral nervous evolution of children as well as cardiovascular disorders. Another disorder that patients with adenotonsillar hypertrophy and airway obstruction may encounter is developmental disorders. Chronic adenotonsillar hypertrophy and subsequent upper airway obstruction appear to cause a disruption in the normal growth of children [3], [4], [5] and weight loss [6]. Several studies have also indicated that adenotonsillar hypertrophy and its subsequent sleep apnea are along with slow growth and weight gain in infants and babies [7], [8], [9]. These disorders

are due to a number of factors such as decreased growth hormone secretion in the sleep phase, decreased insulin-like growth factor 1 (IGF-I) [3], [10], as well as decreased energy intake due to difficulty swallowing and increased energy consumption during the sleep [11].

Due to the importance of the proper growth and weighting process in children and its close relationship to children's health, responsible health systems, especially in developed countries are spending large amounts of money on monitoring and controlling this process in children to identify and resolve the factors that have a negative impact on that [12].

The purpose of this study was to investigate the relationship between body mass index (BMI) and adenotonsillar hypertrophy in children.

Materials and Methods

The study population

In this study, all referral children with adenotonsillar hypertrophy to the clinics of

Khatamolanbia Hospital during the 2017–2018 were evaluated. Inclusion criteria were adenotonsillar hypertrophy, 3–10 years old, and patient satisfaction, and no history of any other concomitant disease to participate in the study.

Procedure

This study was a descriptive-analytical observational study and included 312 children with the age of 3–10 years with adenotonsillar hypertrophy, whose parents were consenting to participate in the study. Age and sex information were collected by asking from parents. The intensity of tonsillar hypertrophy was determined based on Brodsky's scale, where; zero: Tonsil did not affect airway; +1 intensity: Affect the airway <25%; +2 intensity: Affect the airway between 25 and 50%; +3 intensity: Affect the airway between 50-75%, and + 4 intensity: Affect the airway over 75%. In addition, the weight of these children with lightweight clothing without shoes was measured by a pointer scale with a measuring error of 100 g, and their height without shoes was also measured with a 5 mm accuracy height meter.

Statistical analysis

The obtained information was recorded in data collection forms. After completing the sampling, data were analyzed in SPSS software version 22.

Results

BMI

The mean BMI was (85/87±38/43) in the evaluation of patients based on BMI percentile, which among them, the mean BMI for boys was calculated (37/38±42/34) and for girls (01/30±13/66). There was no significant relationship between BMI percentile and age of patients ($p < 0.151$), but there was a significant relationship between this variable and sex ($p < 0.001$).

Statistical analysis also showed a significant difference between the severity of adenotonsillar hypertrophy (Brodsky's scale) and BMI ($p < 0.001$). It should be noted that the correlation between the severity of adenotonsillar hypertrophy and BMI was $R = -0.69$, which indicating a moderate and inverse correlation. Table 1 summarizes the results of statistical analyzes on BMI.

Height growth

The calculated relationship between age and height growth showed that with increasing age, the

Table 1: Impact of age, gender, and severity of adenotonsillar hypertrophy on BMI (by percentile) of children with adenotonsillar hypertrophy, who referred to Khatamolanbia Hospital during 2017–2018

Variable	Mean BMI by percentile	Standard deviation	Value of p	
Age (year)				
3	45/59	44/20	$p < 0/151$	
4	71/07	28/87		
5	39/70	0		
6	27/56	35/64		
7	0/36	0/25		
8	3/6	0		
9	4	0		
10	68/94	25/72		
Sex				$p < 0/001 <$
Male	34/42	38/37		
Female	66/13	30/01		
Adenotonsillar hypertrophy by Brodsky scale			$p < 0/001$	
0	18/60	2/17		
1	20/08	3/48		
2	15/38	1/14		
3	14/30	1/45		
4	14/11	32/35		
Total	43/87	38/85		

BMI: Body mass index.

mean height also shows an increasing and significant correlation between age and height of people ($p < 0.001$) and strong and direct correlation ($R = 0.96$).

Further reviews on height showed that the mean height in males was (83/16±47/113) and in females was (48/15±55/106), and there was a significant relationship between sex and height of patients ($p < 0.002$).

There was no significant relationship between height index and intensity of Tonsillar hypertrophy based on Brodsky's scale ($p = 0.07$) and there was a positive and weak correlation between these two criteria

Table 2: Impact of age, gender, and severity of adenotonsillar hypertrophy on the height growth process of children with adenotonsillar hypertrophy, who referred to Khatamolanbia Hospital during 2017–2018

Variable	Mean height	Standard deviation	Value of p	
Age (year)				
3	90/23	2/10	$p < 0/001$	
4	97/23	4/55		
5	106	0		
6	107/42	2/47		
7	114/93	1/01		
8	124	0		
9	126	0		
10	130/51	10/12		
Sex				$p < 0/002$
Male	113/47	16/83		
Female	106/55	15/48		
Adenotonsillar hypertrophy by Brodsky scale			$p < 0/07$	
0	109/87	24/27		
1	109/05	21/82		
2	114/86	11/70		
3	112/29	12/68		
4	110/77	13/46		
Total	111/41	16/72		

($R = 0.21$). In Table 2 was showed the relationship between age, sex, and severity of adenotonsillar hypertrophy with height growth.

Weight: Similar to the relationship between age and height, there was a significant relationship between age and weight of participants ($p < 0.001$) and the correlation between them ($R = 0.88$) indicated a strong and direction relationship.

Table 3: Impact of age, gender, and severity of adenotonsillar hypertrophy on the weight rate of children with adenotonsillar hypertrophy, who referred to Khatamolanbia Hospital during 2017–2018

Variable	Mean weight	Standard deviation	Value of p	
Age (year)				
3	13/36	1/99	p<0/001	
4	16	0		
5	17	0		
6	16/95	0/82		
7	15/6	1/52		
8	21	0		
9	22	0		
10	33	9/18		
Sex				
Male	21/51	10/54		p=0/34
Female	19/69	5/86		
Adenotonsillar hypertrophy by Brodsky scale				
0	24/21	13/16	p<0/001	
1	25/66	13/53		
2	20/67	5/25		
3	18/44	4/95		
4	17/65	5/63		
Total	20/97	9/42		

The calculated weighted mean in males was (54/10±51/21) and (86/5±69/19) in females. There was no significant relationship between sex and weight of patients ($p = 0.34$) (Table 3). Calculations showed that weight and intensity of Tonsillar hypertrophy based on Brodsky's scale a significant relationship between inverse and mean correlation ($p < 0.001$) ($R = 0.22$).

Discussion

The nose and oral area has a growing lymphoid tissue [13]. Tonsillar hypertrophy is the most common cause of laryngeal obstruction and sleep-related respiratory disorders. Obstructive sleep apnea is a common disease in children with adenotonsillar hypertrophy syndrome. Some of the factors caused by adenotonsillar hypertrophy including too loud snoring, chronic oral breathing, frequent waking at night, excessive drowsiness, nightmares, poor educational performance, and dysphagia. Children with obstruction caused by tonsillar hypertrophy usually eat less and have a lower weight percentage due to loss of appetite [13].

Among the 312 studied patients, 70.2% were male and 29.8% were female. Obviously, in different studies, the proportion of these communities is also different [12], [14], [15], and the reason for this difference is the sampling in different study communities, the population under study, and the male and female populations of different geographic areas. As noted in the findings section, sex was significantly associated with BMI percentile and height. However, there was no relationship between sex and weight of patients. The reason for this can be attributed to the heterogeneity of male and female populations in this study.

The relationship between age and adenotonsillar hypertrophy is somehow that most

studies have been stated that age before 15 is the age of diagnosis. The mean age of the study population was (6/2±53/72) and statistical analysis showed a significant relationship between age and weight and height growth so that the correlation between age and these two variables showed their direct and strong relationship, and in both male and female, by weight increasing and despite adenotonsillar hypertrophy, the height and weight of the child also increase.

In evaluating the relationship of adenotonsillar hypertrophy with BMI, it was found that BMI in growth percentiles was very low in 40.1% of patients, and also 40.4% had proportional weight to age, while 5.1% was overweight, and 14.4% had too much weight (obese). In previous studies, BMI percentiles below 2 were malnourished, between 2 and below 91 were normal, and above 91 considered obese, whereas in our study, based on the latest reference changes, the percentiles below 5% were low weight, 50% to 85% were proportional, and 85% to 95% were overweight and above 95% were obese [16].

Our study indicated that a large population of people with adenotonsillar hypertrophy did not have adequate weight gain. The cause of this can be traced to the effect of adenotonsillar hypertrophy on metabolism. In the Lumeng and Herrmann experiments, it has been demonstrated that chronic adenotonsillar hypertrophy can increase calorie intake and energy loss and consequently impair the development of children [17], [18].

The mean height of 312 patients in our study was 111 cm by percentile (41/21±54/16). About 29.8% of patients had short height, 65.1% moderate height, and 5.1% taller than normal. The mean weight of children in this study was 20 kg which was calculated (19/33±32/27) by percentile. This data indicate growth delay in the studied groups in terms of BMI and height and weight. Thus, it can be observed that, are consistent with other studies, chronic adenotonsillar hypertrophy can impair the normal growth process [8], [19].

It has been proven today that obstructive apnea caused by adenotonsillar hypertrophy, decreases the secretion of growth hormone IGF [20].

IGF-1 is a growth factor that has endocrine and paracrine effects and its secretion is controlled by growth hormone. Most of the existing IGF-1 carries the bond protein that it is called IGF binding protein-3 (IGF BP3).

IGF BP3 synthesis is also controlled by growth hormone. Levels of IGF-1 are applied, so serum levels of IGF BP3 and IGF-1 are considered to evaluate growth hormone levels [8].

Park tests showed that after adenotonsillar surgery, the serum IGF1 levels [21] and height were increased, and the increasing of weight is obvious [15].

In addition to the effect of adenotonsillar hypertrophy on the growth IGF-1 hormone secretion,

it seems that it can provide a suitable substrate for the growth of pathogenic bacteria, especially beta-lactamase-producing groups in tonsillar and adenoid tissue. Growth of bacteria along with obstructive apnea in chronic adenotonsillar hypertrophy can increase metabolism and calorie consumption and subsequently impaired child development [17], [18].

In the reviews of 61 adenotonsillectomized children and comparing them with other 31 healthy children found that troubleshooting such as sleep apnea, nocturnal hypoxia, and sleep disorders, with adenotonsillar surgery, led to an increase in all children.

In the study of growth phenomena and its chemical markers [22], [23] in children with obstructive sleep apnea due to adenotonsillar hypertrophy, 70 children were studied as patients and 35 children as a control group for IGF BP3 and IGF-1. This study showed that during the post-operative phase, the ratio of weight to height and secretion of IGF-1 and IGF-BP3 increased [24].

Conclusion

In this study, we observed a decrease in growth indicators in children with adenotonsillar hypertrophy (height, weight, and BMI). Furthermore, there was a relationship between growth indicators and severity of adenotonsillar hypertrophy in patients.

References

- Hellings PW, Fokkens WJ. Allergic rhinitis and its impact on otorhinolaryngology. *Allergy*. 2006;61(6):656-64. <https://doi.org/10.1111/j.1398-9995.2006.01109.x> PMID:16677233
- Choudhary B, Patil R, Bhatt GC, Dhingra B, Pakhare AP, Goyal A, *et al*. Association of sleep disordered breathing with mono-symptomatic nocturnal enuresis: A study among school children of central India. *PLoS One*. 2016;11(5):e0155808 <https://doi.org/10.1371/journal.pone.0155808> PMID:27191620
- Bonuck KA, Freeman K, Henderson J. Growth and growth biomarker changes after adenotonsillectomy: Systematic review and meta-analysis. *Arch Dis Child*. 2009;94(2):83-91. <https://doi.org/10.1136/adc.2008.141192> PMID:18684748
- Brodsky L, Koch RJ. Bacteriology and immunology of normal and diseased adenoids in children. *Arch Otolaryngol Head Neck Surg*. 1993;119(8):821-9. <https://doi.org/10.1001/archotol.1993.01880200021003> PMID:8343242
- Brodsky L, Radomski K, Gendler J. The effect of post-operative instructions on recovery after tonsillectomy and adenoidectomy. *Int J Pediatr Otorhinolaryngol*. 1993;25(1-3):133-40. [https://doi.org/10.1016/0165-5876\(93\)90046-6](https://doi.org/10.1016/0165-5876(93)90046-6) PMID:8436457
- Bland RM, Bulgarelli S, Ventham JC, Jackson D, Reilly JJ, Paton JY. Total energy expenditure in children with obstructive sleep apnoea syndrome. *Eur Respir J*. 2001;18(1):164-9. <https://doi.org/10.1183/09031936.01.99104401> PMID:11510789
- Capdevila OS, Kheirandish-Gozal L, Dayyat E, Gozal D. Pediatric obstructive sleep apnea: Complications, management, and long-term outcomes. *Proc Am Thorac Soc*. 2008;5(2):274-82. <https://doi.org/10.1513/pats.200708-138mg> PMID:18250221
- Williams EF 3rd, Woo P, Miller R, Kellman RM. The effects of adenotonsillectomy on growth in young children. *Otolaryngol Head Neck Surg*. 1991;104(4):509-16. <https://doi.org/10.1177/019459989110400415> PMID:1903865
- Leach J, Olson J, Hermann J, Manning S. Polysomnographic and clinical findings in children with obstructive sleep apnea. *Arch Otolaryngol Head Neck Surg*. 1992;118(7):741-4. <https://doi.org/10.1001/archotol.1992.01880070071013> PMID:1627296
- Walker MP, Diaugustine RP, Zeringue E, *et al*. An IGF1/insulin receptor substrate-1 pathway stimulates a mitotic kinase (cdk1) in the uterine epithelium during the proliferative response to estradiol. *J Endocrinol*. 2010;207(2):225-35. <https://doi.org/10.1677/joe-10-0102> PMID:20798132
- Greg R, Raymond C. Creative tourism. *ATLAS News* 2000;23:16-20.
- Richart H. *Tarascon Pocket Pharmacopoeia 2015 Deluxe Lab-coat Edition*. United States: Jones & Bartlett Learning; 2015. p. 349.
- Izu SC, Itamoto CH, Pradella-Hallinan M, Pizarro GU, Tufik S, Pignatari S, *et al*. Obstructive sleep apnea syndrome (OSAS) in mouth breathing children. *Braz J Otorhinolaryngol*. 2010;76(5):552-6. PMID:20963335
- Daar G, Sari K, Gencer ZK, Ede H, Aydın R, Saydam L. The relation between childhood obesity and adenotonsillar hypertrophy. *Eur Arch Otorhinolaryngol*. 2016;273(2):505-9. <https://doi.org/10.1007/s00405-015-3554-4> PMID:25876003
- Farmarzi M, Shishegar M, Heydari ST, Haghghi A, Sharouny H. Effects of adenotonsillectomy on serum levels of IGF-1 and IGFBP-3 and growth indices in children with adenotonsillar hypertrophy or recurrent tonsillitis. *Iran J Otorhinolaryngol*. 2016;28(88):329-35. PMID:27738609
- Egeli E, Belli SB, Oghan F, Ozturk O. Relation of isolated tonsillar hypertrophy with body mass index. *ORL J Otorhinolaryngol Relat Spec*. 2006;68(4):228-31. <https://doi.org/10.1159/000092124> PMID:16549938
- Hable KA, Washington JA 2nd, Herrmann EC Jr. Bacterial and viral throat flora. Comparison of findings in children with acute upper respiratory tract disease and in healthy controls during winter. *Clin Pediatr (Phila)*. 1971;10(4):199-203. <https://doi.org/10.1177/000992287101000405> PMID:4396432
- Lumeng JC, Chervin RD. Epidemiology of pediatric obstructive sleep apnea. *Proc Am Thorac Soc*. 2008;5(2):242-52. PMID:18250218
- Schiffmann R, Faber J, Eidelman AI. Obstructive hypertrophic adenoids and tonsils as a cause of infantile failure to thrive:

- Reversed by tonsillectomy and adenoidectomy. *Int J Pediatr Otorhinolaryngol.* 1985;9(2):183-7. [https://doi.org/10.1016/s0165-5876\(85\)80019-7](https://doi.org/10.1016/s0165-5876(85)80019-7)
PMid:4030240
20. Ahlqvist-Rastad J, Hultcrantz E, Melander H, Svanholm H. Body growth in relation to tonsillar enlargement and tonsillectomy. *Int J Pediatr Otorhinolaryngol.* 1992;24(1):55-61. [https://doi.org/10.1016/0165-5876\(92\)90066-x](https://doi.org/10.1016/0165-5876(92)90066-x)
PMid:1399304
21. Park DY, Choi JH, Kang SY, Han J, Park HY, Hwang JS, *et al.* Correlations between pediatric obstructive sleep apnea and longitudinal growth. *Int J Pediatr Otorhinolaryngol.* 2018;106:41-5. PMid:29447889
22. Gibb AG. Unusual complications of tonsil and adenoid removal. *J Laryngol Otol.* 1969;83(12):1159-74. <https://doi.org/10.1017/s0022215100071486>
PMid:5360031
23. Nieminen P, Löppönen T, Tolonen U, Lanning P, Knip M, Löppönen H. Growth and biochemical markers of growth in children with snoring and obstructive sleep apnea. *Pediatrics.* 2002;109(4):e55. <https://doi.org/10.1542/peds.109.4.e55>
PMid:11927728
24. Bar A, Tarasiuk A, Segev Y, Phillip M, Tal A. The effect of adenotonsillectomy on serum insulin-like growth factor-I and growth in children with obstructive sleep apnea syndrome. *J Pediatr.* 1999;135(1):76-80. [https://doi.org/10.1016/s0022-3476\(99\)70331-8](https://doi.org/10.1016/s0022-3476(99)70331-8)
PMid:10393608