



Secretory Carcinoma: A Silent Mass Increasing in the Parotid Gland

Cheng-Chieh Lu¹, Chien-Jui Cheng^{2,3*}, Yu-Chien Kao⁴, Mei-Chien Chen¹

¹Department of Otolaryngology, Head and Neck Surgery, Shuang Ho Hospital, Taipei Medical University, Taipei, Taiwan; ²Department of Pathology, School of Medicine, College of Medicine, Taipei Medical University, Taipei, Taiwan; ³Department of Pathology, Taipei Medical University Hospital, Taipei, Taiwan; ⁴Department of Pathology, Shuang Ho Hospital, Taipei Medical University, Taipei, Taiwan

Abstract

Edited by: Ksenija Bogoeva-Kostovska
Citation: Lu C, Cheng C, Kao Y, Chen M. Secretory Carcinoma: A Silent Mass Increasing in the Parotid Gland. Open Access Maced J Med Sci. 2020 Oct 29; 8(C):191-194. <https://doi.org/10.3889/oamjms.2020.4603>
Keywords: Mammary analog secretory carcinoma; Secretory carcinoma; Salivary gland neoplasms; Salivary gland cancer
***Correspondence:** Dr. Cheng-Chieh Lu, Departments of Otorhinolaryngology: Head and Neck Surgery, Taipei Medical University-Shuang Ho Hospital, No. 291, Zhongzheng Rd., Zhonghe District, New Taipei City, 23561, Taiwan.
E-mail: doc.lulala@gmail.com
Received: 10-Mar-2020
Revised: 15-Oct-2020
Accepted: 19-Oct-2020
Copyright: © 2020 Cheng-Chieh Lu, Chien-Jui Cheng, Yu-Chien Kao, Mei-Chien Chen
Funding: This research did not receive any financial support
Competing Interests: The authors have declared that no competing interests exist
Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

BACKGROUND: Secretory carcinoma (SC) of the salivary gland, also known as mammary analog secretory carcinoma, is a rare tumor in the parotid gland. This kind of tumor is characterized by generally indolent clinical behavior and expression of a break in the *ETV6* gene.

CASE REPORT: We present a unique case of secretory carcinoma and show its favorable prognoses.

CONCLUSION: Secretory carcinoma of the salivary gland is a low-grade carcinoma with a favorable prognosis. It has low regional lymph node and distant metastasis potential. Due to the possibility of misdiagnosis, immunohistochemical studies and FISH are suggested. The most effective treatment is complete surgical excision with negative surgical margins.

Introduction

Secretory carcinoma (SC), formerly known as mammary analog secretory carcinoma (MASC), was renamed by WHO in the Classification of Head and Neck Tumors in 2017 [1]. In the beginning, this kind of tumor was first noticed in the breast and was called juvenile breast carcinoma and secretory carcinoma of the breast [2]. In 2010, Skálová found it occurring in the salivary gland and then named it MASC in salivary glands [3]. Secretory carcinoma is a low-grade malignant tumor and usually occurs in salivary glands, especially in the parotid gland (68%) [4].

It can be confused with acinic cell carcinoma (AciCC) and distinguished by immunohistochemical studies and fluorescence in situ hybridization (FISH). Complete surgical excision of secretory carcinoma is the treatment of choice, while gene fusion data can give us a better understanding of target therapy treatment.

We report a case of secretory carcinoma of the left parotid gland diagnosed and treated in our department. Left superficial parotidectomy was

performed and the FISH study revealed a break-apart of *ETV6* gene. No local recurrence was observed on physical examination and ultrasound on follow-up for 2 years.

Case Report

A 37-year-old man presented with a history of a left parotid mass that was gradually increasing in size for 6 months since October, 2017. No tenderness or facial numbness was noticed. On examination, he presented a firm, fixed, and non-tender mass on the left parotid region of about 6 × 5 cm². Ultrasound showed a cystic mass and the result of fine-needle aspiration cytology report presented atypical cellular change. Contrast-enhanced computed tomography (CT) was performed for pre-surgical operation planning. One 4.1 cm cystic mass with irregular wall thickening at the left parotid gland was noticed (Figure 1).

On the basis of radiological and cytological findings, the left side superficial parotidectomy was




Figure 1: Computed tomography scan of head and neck showing a 4.1 cm cystic mass (arrow) with irregular wall thickening at the left parotid gland

performed under facial nerve monitoring (Figure 2). Histopathology revealed secretory carcinoma composed of cuboid or columnar epithelial cells (Figure 3). Immunohistochemical studies were positive for S-100 and CK7 and negative for p63, CK5/6, and DOG-1. FISH study revealed a break-apart of *ETV6* gene (Figure 4).




Figure 2: Photo of the surgery demonstrating left superficial parotidectomy with preservation of facial nerve

No local recurrence was observed on physical examination and ultrasound on follow-up for 2 years.




Figure 3: Histopathology revealed secretory carcinoma composed of cuboid or columnar epithelial cells




Figure 4: Fluorescence in situ hybridization study showed positive for *ETV6* gene rearrangement with loss of 3' signals (green)

Discussion

Secretory carcinoma of the salivary gland is a rare type of tumor with generally indolent clinical behavior [4]. Its incidence is <0.5% in parotid malignant neoplasms [5] and usually presents as an asymptomatic mass in the parotid gland with slight male predilection (1.5:1) and the age span for SC is wide (5–77 years) [4]. It has also been reported in the nasal cavity, vulva, lacrimal gland [6], eyelid [7], skin [8], thyroid [9], and lung [10]. Most patients with SC present with a slowly growing painless tumor.

The diagnostic plan for SC includes ultrasound and image study (head and neck CT/magnetic resonance imaging). Although fine-needle aspiration may be performed during ultrasound evaluation, the cytology diagnosis for SC is poor. According to Kai *et al.*, only 2 of 109 cytologists identified SC [11]. Immunohistochemistry and FISH studies using pathologic specimen can yield a definite diagnosis. In immunohistochemistry, SC is positive for S100, mammaglobin, and CK7 and negative for p63, CK5/6, and DOG-1 stains in the literature (Table 1) [12]. By FISH, SC shows *ETV6* gene rearrangements. The most well-known fusion partner gene is *NTRK3* (more than 90%) [13]. When fusions happen, the *ETV6* gene enrolls – transcriptional regulator gene and the other genes (ex: *NTRK3*, *MET*, and *RET*) enroll a membrane receptor kinase. The above finding can lead to well-designed-target therapy in the literature [14]: We may choose Trk inhibitors (e.g., entrectinib and larotrectinib) for patients with a *NTRK3* fusion gene and c-Met inhibitors (e.g., cabozantinib) for patients with a *MET* fusion gene if systemic therapy is needed [15] (Table 2).

The treatment of choice for SC is complete surgical excision. Due to the low incidence of lymph node involvement, elective neck dissection is not suggested for every case. Neither radiotherapy nor

chemotherapy is needed after the operation due to its favorable prognosis [4].

Table 1: Comparison of findings between SC and AciCC

	Cytology	Immunostain				FISH
		Mammaglobin	S100	Vimentin	DOG-1	ETV6 split
SC	Papillary fragment, mucoid background	+	++	++	-	+
AciCC	Cell groups and single cells, granular with nuclei background	-	+	+	+	-

SC= secretory carcinoma, AciCC = acinic cell carcinoma

In our case, the immunohistochemical studies were positive for S-100 and CK7 and negative for p63, CK5/6, and DOG-1. FISH study with a break-apart of *ETV6* gene was noticed. The diagnosis of MASC was confirmed.

Table 2: The gene rearrangements of SC and the suggestive target therapy

Transcriptional regulator gene	Membrane receptor kinase-type	Treatment
<i>ETV6</i>	NTRK3(>90%)	Entrectinib and larotrectinib
	MET	RXDX105, LOXO292
	RET(2-5%)	Cabozantinib

Although MASC is considered a low-grade malignant tumor in most cases, recurrence and disease dissemination still happen sometimes. Long-term follow-up is suggested.

Conclusion

Secretory carcinoma of the salivary gland is a low-grade carcinoma with a favorable prognosis. It has low regional lymph node and distant metastasis potential. Due to the possibility of misdiagnosis, immunohistochemical studies and FISH are suggested. The most effective treatment is complete surgical excision with negative surgical margins. FISH is recommended for SC patients, not only to provide definite result but also to obtain information for further target therapy if needed.

Ethical Statement

The paper had complied with the guidelines for human studies and animal welfare regulations. The authors state that subjects have given their informed consent and that the study protocol has been approved by the Institute's Committee on Human Research (TMU-JIRB number: N201812013). The authors also state that animal experiments conform to institutional standards.

References

- Khalele BA. Systematic review of mammary analog secretory carcinoma of salivary glands at 7 years after description. *Head Neck*. 2017;39(6):1243-8. <https://doi.org/10.1002/hed.24755> PMID:28370824
- McDivitt RW, Stewart FW. Breast carcinoma in children. *JAMA*. 1966;195(5):388-90. <https://doi.org/10.1001/jama.195.5.388> PMID:4285563
- Kravtsov O, Hunt B, Peterson J, Carrillo L, Bonneau P, Giorgadze T. Secretory carcinoma of the parotid with adenoid cystic carcinoma cytological pattern: A cytological-pathological correlation with literature review. *Ann Diagn Pathol*. 2018;33:58-61. <https://doi.org/10.1016/j.anndiagpath.2017.12.006> PMID:29566949
- Boon E, Valstar MH, van der Graaf WT, Bloemena E, Willems SM, Meeuwis CA, *et al*. Clinicopathological characteristics and outcome of 31 patients with ETV6-NTRK3 fusion gene confirmed (mammary analogue) secretory carcinoma of salivary glands. *Oral Oncol*. 2018;82:29-33. <https://doi.org/10.1016/j.oraloncology.2018.04.022> PMID:29909898
- Takahama A Jr., de Almeida OP, Kowalski LP. Parotid neoplasms: Analysis of 600 patients attended at a single institution. *Braz J Otorhinolaryngol*. 2009;75(4):497-501. <https://doi.org/10.1590/s1808-86942009000400005> PMID:19784416
- Hycza MD, Andreasen S, Melchior LC, Tucker T, Heegaard S, White VA. Primary secretory carcinoma of the lacrimal gland: Report of a new entity. *Am J Ophthalmol*. 2018;193:178-83. <https://doi.org/10.1016/j.ajo.2018.06.019> PMID:29963997
- Bao Y, Li J, Zhu Y. Mammary analog secretory carcinoma with ETV6 rearrangement arising in the conjunctiva and eyelid. *Am J Dermatopathol*. 2018;40(7):531-5. <https://doi.org/10.1097/dad.0000000000001062> PMID:29251639
- Moore RF, Cuda JD. Secretory carcinoma of the skin: Case report and review of the literature. *JAAD Case Rep*. 2017;3(6):559-62. <https://doi.org/10.1016/j.jidcr.2017.07.003> PMID:29159250
- Liao H, Khan A, Miron PM, Cornejo KM. Mammary analogue secretory carcinoma of the thyroid mimicking locally advanced papillary thyroid carcinoma: A rare case report. *Int J Surg Pathol*. 2018;26(5):459-63. <https://doi.org/10.1177/1066896917747076> PMID:29228842
- Huang T, McHugh JB, Berry GJ, Myers JL. Primary mammary analogue secretory carcinoma of the lung: A case report. *Hum Pathol*. 2018;74:109-13. <https://doi.org/10.1016/j.humpath.2017.10.027> PMID:29104113
- Kai K, Minesaki A, Suzuki K, Monji M, Nakamura M, Tsugitomi H, *et al*. Difficulty in the cytodiagnosis of mammary analogue secretory carcinoma: Survey of 109 cytologists with a case originating from a minor salivary gland. *Acta Cytol*. 2017;61(6):469-76. <https://doi.org/10.1159/000477390> PMID:28738326
- Badlani J, Gupta R, Balasubramanian D, Smith J, Luk P, Clark J. Primary salivary gland malignancies: A review of clinicopathological evolution, molecular mechanisms and management. *ANZ J Surg*. 2018;88(3):152-7. <https://doi.org/10.1111/ans.14201> PMID:28982213
- Skalova A, Stenman G, Simpson RH, Hellquist H, Slouka D,

-
- Svoboda T, *et al.* The role of molecular testing in the differential diagnosis of salivary gland carcinomas. *Am J Surg Pathol.* 2018;42(2):e11-27. <https://doi.org/10.1097/pas.0000000000000980>
PMid:29076877
14. Rooper LM, Karantanos T, Ning Y, Bishop JA, Gordon SW, Kang H, *et al.* Salivary secretory carcinoma with a novel ETV6-MET fusion: Expanding the molecular spectrum of a recently described entity. *Am J Surg Pathol.* 2018;42(8):1121-6. <https://doi.org/10.1097/pas.0000000000001065>
PMid:29683815
15. Skalova A, Vanecek T, Martinek P, Weinreb I, Stevens TM, Simpson RH, *et al.* Molecular profiling of mammary analog secretory carcinoma revealed a subset of tumors harboring a novel ETV6-RET translocation: Report of 10 cases. *Am J Surg Pathol.* 2018;42(2):234-46. <https://doi.org/10.1097/pas.0000000000000972>
PMid:29076873