



# Serum Granulysin in Differentiation of Stevens-Johnson Syndrome/toxic Epidermal Necrolysis and Erythema Multiforme

Tran Thi Huyen<sup>1,2\*</sup>, Pham Dinh Hoa<sup>1,2</sup>, Trinh Minh Trang<sup>2</sup>, Riichiro Abe<sup>3</sup>, Nguyen Van Thuong<sup>1,2</sup>, Pham Thi Lan<sup>1,2</sup>, Michael Tirant<sup>1</sup>

<sup>1</sup>Department of Dermatology and Venereology, Hanoi Medical University, Hanoi, Vietnam; <sup>2</sup>Department of Skin Diseases of Women and Children, National Hospital of Dermatology and Venereology, Hanoi, Vietnam; <sup>3</sup>Department of Dermatology, Niigata University, Japan

## Abstract

**Edited by:** Ksenija Bogoeva-Kostovska

**Citation:** Huyen TT, Hoa PD, Trang TM, Abe R, Van Thuong N, Lan PT, Tirant M. Serum Granulysin in Differentiation of Stevens-Johnson Syndrome/toxic Epidermal Necrolysis and Erythema Multiforme. Open Access Maced J Med Sci. 2020 Jun 10; 8(B):381-388. https://doi.org/10.3889/oamjms.2020.4905

**Keywords:** Stevens-Johnson syndrome; Toxic epidermal necrolysis; Serum granulysin; Erythema multiforme, re-epithelialization; SCORTEN

**\*Correspondence:** Tran Thi Huyen, Hanoi Medical University, Hanoi, Vietnam, 1A Ton That Tung street, Hanoi, Vietnam. E-mail: drhuyentran@gmail.com

**Received:** 07-May-2020

**Revised:** 24-May-2020

**Accepted:** 29-May-2020

**Copyright:** © 2020 Tran Thi Huyen, Pham Dinh Hoa, Trinh Minh Trang, Riichiro Abe, Nguyen Van Thuong, Pham Thi Lan, Michael Tirant

**Funding:** This research did not receive any financial support

**Competing Interest:** The authors have declared that no competing interest exists

**Open Access:** This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

**BACKGROUND:** Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) are acute, life-threatening drug reactions, which lead to massive epidermal necrolysis. Granulysin plays an important role as a key mediator for keratinocyte apoptosis in these conditions. Erythema multiforme (EM) may have skin manifestation similar to SJS/TEN.

**AIMS:** The aim of the study was to compare serum granulysin levels in patients with SJS/TEN and EM as well as to investigate a possible association between serum granulysin levels and the severity of SJS/TEN.

**METHODS:** In total, 48 patients with SJS/TEN, 43 patients with EM, and 20 health controls (HCs) were enrolled. We measured serum granulysin levels using enzyme-linked immunosorbent assay.

**RESULTS:** The average level of serum granulysin in the SJS/TEN patients was 23.0 ng/ml (range 1.2–144.6 ng/ml), significantly higher than that of EM group (20.1 ng/ml; range 8.5–121 ng/ml,  $p < 0.05$ ) and HCs group (20.8 ng/ml; range 10.1–46.7 ng/ml,  $p < 0.05$ ). Of 48 SJS/TEN patients, the 25 samples collected  $< 6$  days after onset showed higher level of serum granulysin (27.7 ng/ml; range 2.5–144.6 ng/ml) than those collected  $\geq 6$  days after onset (17.9 ng/ml; range 1.2–59 ng/ml;  $p > 0.05$ ). No significant correlation was found between serum granulysin levels and the body surface area affected and the modified-SCORTEN. At the day of re-epithelialization, serum granulysin levels were not different compared with those at the day of hospitalization.

**CONCLUSIONS:** Serum granulysin levels are significantly higher in SJS/TEN group than in EM group. After the onset, serum granulysin levels in patients with SJS/TEN are not a good biomarker to evaluate the severity of the diseases.

## Introduction

Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN) are severe cutaneous adverse drug reactions (SCARs) characterized by extensive epidermal necrolysis, blisters, and skin sloughing [1], [2]. The most common causative drugs of SJS/TEN are carbamazepine, allopurinol, abacavir, phenytoin, and lamotrigine [1], [3], [4], [5]. The period between taking a drug and onset of symptoms ranges from a few days to 2 months [1], [2], [6]. SJS and TEN are categorized based on the percentage of epidermal detachment area: (i) SJS:  $< 10\%$ , (ii) TEN:  $> 30\%$ , and (iii) overlapping SJS/TEN:  $10\text{--}30\%$  [2].

The pathogenesis of SJS/TEN is not fully understood, but there are some immunological and genetic factors which are believed to be involved [4], [5], [7], [8], [9]. There is a strong association between *HLA-B\*15:02* and carbamazepine-induced SJS/TEN [10], [11], [12], *HLA-B\*58:01* and allopurinol-induced SJS/TEN [13], [14], [15], and *HLA-B\*57:01* and abacavir-induced SJS/TEN [15], [16], [17]. CD8+ cytotoxic

T cells (CTLs) and natural killer (NK) cells play an important role in the pathogenesis of SJS/TEN [1], [4], [5], [8], [9]. The immune response may be triggered by binding an antigenic drug to a specific *HLA* on a keratinocyte [4], [5]. Specific T cell receptors recognize the drug-*HLA* complex and upon the activation, CD8+CTLs and NK cells produce cytokines, chemokines, and cytotoxic proteins, particularly granulysin, that cause extensive keratinocytes apoptosis [4], [5]. Chung reported that granulysin concentrations in the blister fluid of SJS/TEN patients were 2–4 times higher than perforin, granzyme B, or soluble Fas ligand concentrations [18]. Granulysin in the blister fluid was 15-kDa secretory form, and injection of it into mouse skin resulted in features mimicking SJS/TEN [5], [18], [19]. Abe showed that serum granulysin levels in patients with SJS/TEN were elevated before skin detachment or mucosal lesions developed [20]. The rapid immunochromatographic test has been developed for serum granulysin to diagnose early SJS/TEN [21].

Erythema multiforme (EM) may have skin manifestation similar to SJS/TEN [2], [22], [23], but they can be distinguished immunopathologically. In cases of SJS/TEN, the inflammatory infiltrates expressing

granulysin, granzyme B, and perforin accumulated predominantly in the lower epidermal and subepidermal bulla, in contrast, they were relatively sparse in EM [24]. However, this test is not clinically rapid. On the other hand, serum granulysin levels may be elevated among SJS/TEN patients [20], [25]. To the best of our knowledge, there is no study compared serum granulysin levels in SJS/TEN with those in EM. We conducted this study to investigate serum granulysin levels in patients with SJS/TEN and EM as well as possible associations between serum granulysin levels and the severity of SJS/TEN.

## Materials and Methods

### Participants

#### *SJS/TEN patients*

In total, 48 patients with SJS/TEN were enrolled from January 2018 to October 2019 at two hospitals in Hanoi, Vietnam (National Hospital of Dermato-venereology and Bach Mai Hospital). The SJS/TEN patients had their vital signs and systemic symptoms and the percentage of body surface area affected (skin detachment) examined. SJS and TEN were classified in accordance with Bastuji-Garin *et al.* [2]. All patients were aged 18 or older. The onset was defined as the day mucocutaneous or ocular lesions were first eroded or ulcerated [20]; the re-epithelialization was defined as complete healing of the skin without any erosion [26]. The severity of SJS/TEN patients was also assessed by the modified-SCORTEN (SCORE of TEN) scale that was composed of six well-defined criteria, including age >40 years, malignancy, tachycardia >120 beats/min, percentage of epidermal detachment >10%, serum urea >10 mmol/L, and serum glucose >14 mmol/L. This scale was based on SCORTEN scale of Bastuji-Garin *et al.* consisting of seven criteria [27].

#### *EM patients and health controls (HCs)*

There were 43 patients with EM recruited in this study. They had the presence of typical or atypical cutaneous target lesions, with or without mucous membrane lesions. The causes of EM were either drugs or unknown. Twenty HCs recruited were healthy medical staff in the National Hospital of Dermatology and Venereology, Ha Noi, Vietnam.

#### *Measuring serum granulysin levels*

In the SJS/TEN group, we took blood samples at 2-time points (1) at the day of hospitalization and (2) at the day of re-epithelialization. In the EM group, blood samples were taken at the day of hospitalization.

All blood samples were left to coagulate at room temperature 10–20 min, then centrifuged in 20 min at a speed of 2000–000 r.p.m, finally serum was taken and stored at  $-20^{\circ}\text{C}$  until proceeding the granulysin measurement.

Using enzyme-linked immunosorbent assay (ELISA) (Human Granulysin ELISA Kit, MELISIN, China), we quantified granulysin in all serum samples. The kit used ELISA-double antibody sandwich principle to assess granulysin levels. The microELISA strip plate provided in this kit has been coated by purified granulysin antibody to make solid-phase antibody, in sequence granulysin is added to wells, combined with granulysin antibody labeled by horseradish peroxidase, which then becomes an antibody-antigen-enzyme-antibody complex. After washing completely to remove the uncombined enzyme, chromogen solution A and chromogen solution B were added, changing the color of the liquid to blue which in turn turns into yellow due to the effect of acid. The color change was measured spectrophotometrically at a wavelength of 450 nm. The concentration of granulysin in the samples was then determined by comparing the O.D. of the samples to the standard curve. The minimum detectable dose of granulysin is typically <0.15 ng/ml (following manufacturer's instruction).

#### *Statistical analysis and ethical clearance*

Data entry and analysis were conducted by using SPSS software version 16.0 (IBM, Armonk, NY, USA). The Mann–Whitney U test and Wilcoxon test were used to compare quantitative variables. Differences were considered to be statistically significant at  $p < 0.05$ .

The study was approved by the Ethical Review Committee on Research Involving Human Subjects, Hanoi Medical University (Number 04NCS17, dated February 08, 2018). Written consent was obtained from all participants.

## Results

There were 19 SJS patients (39.5%) and 29 TEN patients (60.5%) participating in our study. Their characteristics are shown in Table 1. The mean age was 49.3, range 19–77 years (47.9% males; 52.1% females). The most common causative drugs were traditional medicine (29.1%), carbamazepine (12.5%), and allopurinol (12.5%). The time between the onset and the day of hospitalization was  $5.9 \pm 2.7$  days (range 2–18 days). The mean body surface area affected was 43.7%. The mean m-SCORTEN score was 1.6. The mean time of re-epithelialization was 15.9 days (range 9–31 days). All SJS/TEN patients got re-epithelialization and total resolution, no one with in-hospital mortality.

The characteristics of EM patients are shown in Table 2. The mean age was 41.4, range 19–76 (30.2% males; 69.8% females). The medication was responsible for 41.9% of patients with EM.

**Table 1: Characteristics of patients with SJS/TEN**

Characteristics	SJS (n=19)	TEN (n=29)	SJS/TEN (n=48)
Age, years (Range)	44.9±15.3 (19-72)	52.2±14.3 (25-77)	49.3±15.0 (19-77)
Sex, n (%)			
Male	11 (57.9)	12 (41.4)	23 (47.9)
Female	8 (42.1)	17 (58.6)	25 (52.1)
Causative drugs, n (%)			
Carbamazepine	5 (26.2)	1 (3.4)	6 (12.5)
Allopurinol	4 (21.1)	2 (6.9)	6 (12.5)
Traditional medicine	2 (10.5)	12 (41.5)	14 (29.1)
Antibiotics	1 (5.3)	2 (6.9)	3 (6.2)
NSAIDs	1 (5.3)	3 (10.3)	4 (8.4)
Thalidomide	0 (0)	1 (3.4)	1 (2.1)
Unknown	6 (31.6)	8 (27.6)	14 (29.2)
The time between onset and hospitalization, days	5.5 ± 1.6	6.1 ± 3.2	5.9 ± 2.7
Taking corticosteroids before hospitalization, n (%)			
Yes	7 (36.8)	13 (44.8)	20 (41.7)
No	8 (42.1)	13 (44.8)	21 (43.7)
Unknown	4 (21.1)	3 (10.4)	7 (14.6)
Body surface area affected, %	7±7.4	67.8±21.7	43.7±34.7
m-SCORTEN score	1±0.9	2±0.7	1.6±0.9
Fever, n (%)	8 (42.1)	19 (65.5)	27 (56.2)
High levels of liver enzymes, n (%)	11 (57.9)	19 (65.5)	30 (62.5)
Malfunction of kidney	6 (31.6)	14 (48.3)	20 (41.7)
Mucous membrane lesions	18 (94.7)	21 (72.4)	39 (81.2)
Pneumonia	0 (0)	8 (27.6)	8 (16.7)
Treatment given during hospitalization, n (%)			
Systemic corticoid	16 (84.2)	19 (65.5)	35 (72.9)
Ciclosporin A	3 (15.8)	8 (27.6)	11 (22.9)
Only care support	0 (0)	2 (6.9)	2 (4.2)
The time of re-epithelialization, days (Range)	13.8±3.7 (9–23)	17.2±4.6 (10–31)	15.9±4.6 (9–31)

SJS: Stevens-Johnson syndrome, TEN: Toxic epidermal necrolysis, NSAIDs: Nonsteroid anti-inflammatory drugs, SCORTEN: SCORe of toxic epidermal necrolysis


**Table 2: Characteristics of patients with EM and HCs**

Characteristics	EM (n=43)	HCs (n=20)
Age, years (range)	41.4±17.3 (19–76)	28.4±3.5 (25–37)
Sex, n (%)		
Male	13 (30.2)	10 (50)
Female	30 (69.8)	10 (50)
Causes, n (%)		NA
Medication	18 (41.9)	
Traditional medicine	8 (44.4)	
Antibiotics	5 (27.8)	
Allopurinol	3 (16.6)	
Herbal food	2 (11.2)	
Unknown	25 (58.1)	
Cutaneous lesions, n (%)		NA
Typical targets	7 (16.3)	
Atypical targets	30 (69.7)	
Both typical and atypical targets	6 (14)	
Mucous membrane lesion, n (%)	9 (20.9)	NA
Fever, n (%)	10 (23.3)	NA

EM: Erythema multiforme, HCs: Healthy controls, NA: Not applicable.

Serum granulysin levels of patients with SJS/TEN (23.0 ng/ml; range 1.2–144.6 ng/ml) were significantly higher than those of patients with EM (20.1 ng/ml; range 8.5–121 ng/ml;  $p < 0.05$ ) and those of HCs (20.8 ng/ml; range 10.1–46.7 ng/ml;  $p < 0.05$ ) (Figure 1). There was no significant difference with regard to serum granulysin levels among SJS and TEN patients (21.7 ng/ml versus 23.9 ng/ml;  $p > 0.05$ ) (Figure 2a). In the SJS/TEN group, serum granulysin levels were higher in the 25 samples collected <6 days of the onset (27.7 ng/ml; range 2.5–144.6 ng/ml) than in the 23 samples collected ≥6 days after the onset (17.9 ng/ml; range 1.2–59 ng/ml) but it was not significantly different ( $p > 0.05$ ) (Figure 2b). Serum granulysin levels were not significantly different between male and female patients with SJS/TEN (Figure 2c). In patients with SJS/TEN, there was a negative correlation between serum

granulysin levels and the ages with  $r = (-0.25)$ , but it was not statistically significant ( $p = 0.081$ ) (Figure 3a). There was no correlation between serum granulysin levels and the body surface area affected ( $r = 0.12$ ;  $p = 0.417$ ) (Figure 3b); between serum granulysin levels and modified-SCORTEN with  $r = (-0.16)$ ,  $p = 0.29$  (Figure 3c). Serum granulysin levels were not affected by some clinical/paraclinical such as fever, malfunction of kidney, pneumonia, high level of liver enzymes, and mucous membrane lesion (Figure 4). We had a TEN patient whose progress of disease is shown in Figure 5.




**Figure 1: Serum granulysin levels of healthy controls, erythema multiforme group, and Stevens-Johnson syndrome/toxic epidermal necrolysis  $p > 0.05$  group**

## Discussion

This study showed that serum granulysin levels in SJS/TEN patients were statistically higher than those in EM patients. There was no significant difference with regard to serum granulysin levels in SJS and TEN patients. Serum granulysin levels in patients hospitalized <6 days of the onset were higher than after ≥6 days; however, it was not statistically significant.

Granulysin is a cytolytic molecule presenting in human CTL and NK cell granules [28] and is lytic against a variety of tumor cell targets and microbes [28], [29]. In combination with purified perforin, recombinant granulysin breaks up 90% of intracellular *Mycobacterium tuberculosis*, inducing lesions on the mycobacterial cell surface [29]. Previous studies as regards SJS/TEN support the concept that SJS/TEN are SCARs initiated by CTLs [5], [8], [18], [30]. It has been observed that in patients with SJS/TEN, CTLs, and NK cells infiltrate the skin lesions, which point to cutaneous recruitment of antigen-primed CTLs in the pathogenesis of SJS/TEN [8], [18], [31]. Chung demonstrated that SJS/TEN blister cells were, in most part, formed by CD8+CTLs, CD56+NK, and NK cells,



**Figure 2:** Comparison of serum granulysin levels (ng/ml) in SJS/TEN: Between SJS and TEN (a); the 25 samples collected <6 days of the onset and the 23 samples collected ≥6 days after the onset (b); male and female patients (c); At the day of hospitalization and re-epithelialization (d). SJS: Stevens-Johnson syndrome; TEN: toxic epidermal necrolysis


and these effector cells showed cytotoxicity against target cells [18]. These findings are favorable to the key role of granulysin, a product of CTLs and NK cells, in the pathogenesis of SJS/TEN.

EM is a cutaneous reaction characterized by typical or atypical target lesions that mimic cutaneous manifestations of SJS/TEN in the early phase [2], [22], [23]. The causes of EM may be a viral infection or drug reaction [22], [23], [32]. Serum granulysin can be also elevated in viral infected conditions [33] such as virus-associated EM. Nevertheless, our results may imply the use of serum granulysin as a biomarker for distinguishing between EM and SJS/TEN. In fact, serum granulysin levels can be affected by some factors such as infection [33], [34], [35], sepsis [36], cancers [34], [37], age [33], and immunological condition [35], hence, it was a large range among our SJS/TEN patients. However, systemic corticosteroid treatment before hospitalization did not affect serum granulysin levels in patients with SJS/TEN.


In the present study, there was no association between serum granulysin levels and some clinical/paraclinical manifestations of SJS/TEN. At the day of re-epithelialization, serum granulysin levels were not significantly different compared with those at the day of being hospitalized. These findings could be explained by the fact that nearly all the 48 patients with SJS/TEN in this study had their serum granulysin measured after the onset (mean 5.9 days) when serum granulysin levels could be decreased. Abe *et al.* showed that in SJS/TEN, serum granulysin levels were elevated before the onset (when skin detachment or mucosal lesion develop) and higher than those of ordinary drug-induced skin reactions [20]. However, the mean level of serum granulysin in this study was consistent with that in Abe's *et al.* study (24.8 ng/ml) [20]. It could be due to the difference in the size of samples in each study.

By using the sandwich ELISA method, a study in Japan shows the mean level of serum granulysin among





**Figure 3:** Correlation between serum granulysin levels in SJS/TEN patients and the ages (a), the body surface area affected (b), the modified-SCORTEN (c). SJS: Stevens-Johnson syndrome; TEN: Toxic epidermal necrolysis; SCORTEN: SCORE of toxic epidermal necrolysis



**Figure 4:** Serum granulysin levels (ng/ml) and some clinical/paraclinical in patients with SJS/TEN; \* $p < 0.05$ . SJS: Stevens-Johnson syndrome; TEN: Toxic epidermal necrolysis

healthy individuals is 3.7 ng/ml, and it also demonstrates a positive correlation between serum granulysin

levels and age, but no significant difference in serum granulysin levels between genders [33]. Furthermore,




Figure 5: A 52-year-old male patient with allopurinol-induced toxic epidermal necrolysis and serum granulysin level results. (a) At the day of hospitalization: Serum granulysin level was 22.2 ng/ml, (b) ten days after hospitalization, extensive necrolysis of skin, (c) at the day of re-epithelialization: Serum granulysin level was 17.1 ng/ml

a study among health staffs from tuberculosis hospitals in Hanoi, Vietnam, also shows lower levels of serum granulysin as compared with those of the present study [35]. This may be due to the difference in study populations, age, and immunological conditions. On the other hand, HCs in our study were medical staff, who possibly have higher chances of being exposed to bacterial or viral infections in hospital environments.

Cho *et al.* showed that the serum level of granulysin of patients with generalized bullous fixed drug eruption (GBFDE) was significantly lower than that of patients with SJS/TEN overlap or TEN [25]. In GBFDE, granulysin is mainly produced by NK

cells, whilst in SJS/TEN both NK cells and CTLs play an essential role. The presence of granulysin in lesional skin and in serum is a principal factor to distinguish GBFDE from SJS/TEN [25]. Furthermore, serum granulysin levels are not elevated exclusively in SJS/TEN; other conditions have also shown to be responsible for high serum granulysin levels, such as graft-versus-host disease [33], [38], drug reaction with eosinophilia and systemic symptoms, or drug-induced hypersensitivity syndrome [39]. Another study in Taiwan shows that high plasma granulysin levels were associated with renal impairment and mortality rates of patients with allopurinol-induced SJS/TEN [14]. Deceased patients showed higher concentrations of granulysin for extended periods of time when compared to the survivors [14]. Su *et al.* demonstrated that the levels of interleukin-15 and granulysin were significantly correlated with the disease extension in SJS/TEN [36]. These studies implied that serum granulysin levels may be used as a biomarker for not only the diagnosis but also the prognosis of SCARs. However, in our study, after the onset of SJS/TEN, serum granulysin level is not useful for evaluating the severity of SJS/TEN.

This study had some limitations. First, we collected serum samples several days after the onset of SJS/TEN when serum granulysin levels might have been decreased a lot. Second, HCs group were healthy health staffs whose serum granulysin levels might be higher than those of the general population. Third, we only measured granulysin levels in sera and were not able to quantify them in blister fluids and identify the presence of granulysin on skin tissue. Fourth, we did not identify the cells secreting granulysin. In fact, there are other cytotoxic proteins and molecules such as perforin, granzyme B [5], [18], [40], and annexin A1 [41] relating to the pathogenesis of SJS/TEN but we could not investigate all of them. Finally, we used modified-SCORTEN as serum bicarbonate test was not available in the study settings. However, the mean m-SCORTEN in this study was consistent with the mean SCORTEN of a previous study in Vietnam [11].

## Conclusion

Serum granulysin levels are significantly higher in SJS/TEN patients than in EM patients. After the onset of SJS/TEN, serum granulysin is not associated with the severity of the diseases.

## References

1. Schwartz RA, McDonough PH, Lee BW. Toxic epidermal necrolysis: Part I. Introduction, history, classification,

- clinical features, systemic manifestations, etiology, and immunopathogenesis. *J Am Acad Dermatol* 2013;69(2):173.e1-13; quiz 185-6.  
PMid:23866878
2. Bastuji-Garin S, Rzany B, Stern RS, Shear NH, Naldi L, Roujeau JC. Clinical classification of cases of toxic epidermal necrolysis, Stevens-Johnson syndrome, and erythema multiforme. *Arch Dermatol* 1993;129(1):92-6. <https://doi.org/10.1001/archderm.129.1.92>
  3. Sassolas B, Haddad C, Mockenhaupt M. ALDEN, an algorithm for assessment of drug causality in Stevens-Johnson syndrome and toxic epidermal necrolysis: Comparison with case-control analysis. *Clin Pharmacol Ther* 2010;88(01):60-8. <https://doi.org/10.1038/clpt.2009.252>  
PMid:20375998
  4. Chung WH, Wang CW, Dao RL. Severe cutaneous adverse drug reactions. *J Dermatol* 2016;43(7):758-66. <https://doi.org/10.1111/1346-8138.13430>  
PMid:27154258
  5. Su SC, Chung WH. Cytotoxic proteins and therapeutic targets in severe cutaneous adverse reactions. *Toxins* 2014;6(1):194-210. <https://doi.org/10.3390/toxins6010194>  
PMid:24394640
  6. Creamer D, Walsh SA, Dziewulski P, Exton LS, Lee HY, Dart JK, et al. U.K. guidelines for the management of Stevens-Johnson syndrome/toxic epidermal necrolysis in adults 2016. *Br J Dermatol* 2016;174:1194-227. <https://doi.org/10.1016/j.jaci.2010.11.048>  
PMid:27216885
  7. Pichler WJ, Naisbitt DJ, Park BK. Immune pathomechanism of drug hypersensitivity reactions. *J Allergy Clin Immunol* 2011;127(3 Suppl):S74-81.  
PMid:21354503
  8. Nassif A, Bensussan A, Boumsell L, Deniaud A, Moslehi H, Wolkenstein P, et al. Toxic epidermal necrolysis: Effector cells are drug-specific cytotoxic T cells. *J Allergy Clin Immunol* 2004;114(5):1209-15. <https://doi.org/10.1016/j.jaci.2004.07.047>  
PMid:15536433
  9. Nassif A, Bensussan A, Dorothée G, Mami-Chouaib F, Bachot N, Bagot M, et al. Drug specific cytotoxic T-cells in the skin lesions of a patient with toxic epidermal necrolysis. *J Invest Dermatol* 2002;118(4):728-33. <https://doi.org/10.1046/j.1523-1747.2002.01622.x>  
PMid:11918724
  10. Man CB, Kwan P, Baum L, Yu E, Lau KM, Cheng AS, et al. Association between HLA-B\*1502 allele and antiepileptic drug-induced cutaneous reactions in Han Chinese. *Epilepsia* 2007;48(5):1015-8. <https://doi.org/10.1111/j.1528-1167.2007.01022.x>  
PMid:17509004
  11. Nguyen DV, Chu HC, Nguyen DV, Phan MH, Craig T, Baumgart K, et al. HLA-B\*1502 and carbamazepine-induced severe cutaneous adverse drug reactions in Vietnamese. *Asia Pac Allergy* 2015;5:68-77. <https://doi.org/10.5415/apallergy.2015.5.2.68>  
PMid:25938071
  12. Yang CW, Hung SI, Juo CG, Lin YP, Fang WH, Lu IH, et al. HLA-B\*1502-bound peptides: Implications for the pathogenesis of carbamazepine-induced Stevens-Johnson syndrome. *J Allergy Clin Immunol* 2007;120(4):870-7. <https://doi.org/10.1016/j.jaci.2007.06.017>  
PMid:17697703
  13. Hung SI, Chung WH, Liou LB, Chu CC, Lin M, Huang HP, et al. HLA-B\*5801 allele as a genetic marker for severe cutaneous adverse reactions caused by allopurinol. *Proc Natl Acad Sci U S A* 2005;102(11):4134-9. <https://doi.org/10.3410/f.1088014.540977>  
PMid:15743917
  14. Chung WH, Chang WC, Stocker SL, Juo CG, Graham GG, Lee MH, et al. Insights into the poor prognosis of allopurinol-induced severe cutaneous adverse reactions: The impact of renal insufficiency, high plasma levels of oxypurinol and granulysin. *Ann Rheum Dis* 2015;74(12):2157-64. <https://doi.org/10.1136/annrheumdis-2014-205577>  
PMid:25115449
  15. Fricke-Galindo I, LLerena A, López-López M. An update on HLA alleles associated with adverse drug reactions. *Drug Metab Pers Ther* 2017;32(2):73-87. <https://doi.org/10.1515/dmpt-2016-0025>  
PMid:28315856
  16. Mounzer K, Hsu R, Fusco JS, Brunet L, Henegar CE, Vannappagari V, et al. HLA-B\*57:01 screening and hypersensitivity reaction to abacavir between 1999 and 2016 in the OPERA® observational database: A cohort study. *AIDS Res Ther* 2019;16(1):1. <https://doi.org/10.1186/s12981-019-0217-3>  
PMid:30651100
  17. Martínez Buitrago E, Oñate JM, García-Goez JF, Álvarez J, Lenis W, Sañudo LM, et al. HLA-B\*57:01 allele prevalence in treatment-Naïve HIV-infected patients from Colombia. *BMC Infect Dis* 2019;19(1):793. <https://doi.org/10.1186/s12879-019-4415-3>  
PMid:31500584
  18. Chung WH, Hung SI, Yang JY, Su SC, Huang SP, Wei CY, et al. Granulysin is a key mediator for disseminated keratinocyte death in Stevens-Johnson syndrome and toxic epidermal necrolysis. *Nat Med* 2008;14(12):1343-50. <https://doi.org/10.1038/nm.1884>  
PMid:19029983
  19. Krensky AM, Clayberger C. Biology and clinical relevance of granulysin. *Tissue Antigens* 2009;73(3):193-8. <https://doi.org/10.1111/j.1399-0039.2008.01218.x>  
PMid:19254247
  20. Abe R, Yoshioka N, Murata J, Fujita Y, Shimizu H. Granulysin as a marker for early diagnosis of the Stevens-Johnson syndrome. *Ann Intern Med* 2009;151(7):514-5. <https://doi.org/10.7326/0003-4819-151-7-200910060-00016>  
PMid:19805776
  21. Fujita Y, Yoshioka N, Abe R, Murata J, Hoshina D, Mae H, et al. Rapid immunochromatographic test for serum granulysin is useful for the prediction of Stevens-Johnson syndrome and toxic epidermal necrolysis. *J Am Acad Dermatol* 2011;65(1):65-8. <https://doi.org/10.1016/j.jaad.2010.04.042>  
PMid:21507517
  22. Auquier-Dunant A, Mockenhaupt M, Naldi L. Correlations between clinical patterns and causes of erythema multiforme major, Stevens-Johnson syndrome, and toxic epidermal necrolysis: Result of an international prospective study. *Arch Dermatol* 2002;138:1019-24. <https://doi.org/10.1001/archderm.138.8.1019>  
PMid:12164739
  23. Morsy H, Taha EA, Nigm DA, Shahin R, Youssef EM. Serum IL-17 in patients with erythema multiforme or Stevens-Johnson syndrome/toxic epidermal necrolysis drug reaction, and correlation with disease severity. *Clin Exp Dermatol* 2017;42(8):868-73. <https://doi.org/10.1111/ced.13213>  
PMid:28940568
  24. Iwai S, Sueki H, Watanabe H, Sasaki Y, Suzuki T, Iijima M. Distinguishing between erythema multiforme major and Stevens-Johnson syndrome/toxic epidermal necrolysis immunopathologically. *J Dermatol* 2012;39(9):781-6. <https://doi.org/10.1111/j.1346-8138.2012.01532.x>



- PMid:22458564
25. Cho YT, Lin JW, Chen YC, Chang CY, Hsiao CH, Chung WH, *et al.* Generalized bullous fixed drug eruption is distinct from Stevens-Johnson syndrome/toxic epidermal necrolysis by immunohistopathological features. *J Am Acad Dermatol* 2014;70(3):539-48. <https://doi.org/10.1016/j.jaad.2013.11.015>
26. Valeyrie-Allanore L, Wolkenstein P, Brochard L, Ortonne N, Maître B, Revuz J, *et al.* Open trial of ciclosporin treatment for Stevens-Johnson syndrome and toxic epidermal necrolysis. *Br J Dermatol* 2010;163(4):847-53. <https://doi.org/10.1111/j.1365-2133.2010.09863.x>  
PMid:20500799
27. Bastuji-Garin S, Fouchard N, Bertocchi M, Roujeau JC, Revuz J, Wolkenstein P. SCORTEN: A severity-of-illness score for toxic epidermal necrolysis. *J Invest Dermatol* 2000;115(2):149-53. <https://doi.org/10.1046/j.1523-1747.2000.00061>  
PMid:10951229
28. Peña SV, Hanson DA, Carr BA, Goralski TJ, Krensky AM. Processing, subcellular localization, and function of 519 (granulysin), a human late T cell activation molecule with homology to small, lytic, granule proteins. *J Immunol* 1997;158(6):2680-8.  
PMid:9058801
29. Stenger S, Hanson DA, Teitelbaum R, Dewan P, Niazi KR, Froelich CJ, *et al.* An antimicrobial activity of cytolytic T cells mediated by granulysin. *Science* 1998;282(5386):121-5. <https://doi.org/10.1126/science.282.5386.121>  
PMid:9756476
30. Chung WH, Hung SI. Genetic markers and danger signals in Stevens-Johnson syndrome and toxic epidermal necrolysis. *Allergol Int J Jpn Soc Allergol* 2010;59(4):325-32. <https://doi.org/10.2332/allergolint.10-rai-0261>  
PMid:20962567
31. Chung WH, Hung SI, Hong HS, Hsieh MS, Yang LS, Ho HC, *et al.* Medical genetics: A marker for Stevens-Johnson syndrome. *Nature* 2004;428(6982):486.  
PMid:15057820
32. Akkurt ZM, Uçmak D, Türkcü G, Yüksel H, Yıldız K, Arıca M. Expression of interleukin-17 in lesions of erythema multiforme may indicate a role for T helper 17 cells. *Cent-Eur J Immunol* 2014;39(3):370-6. <https://doi.org/10.5114/ceji.2014.45950>  
PMid:26155150
33. Ogawa K, Takamori Y, Suzuki K, Nagasawa M, Takano S, Kasahara Y, *et al.* Granulysin in human serum as a marker of cell-mediated immunity. *Eur J Immunol* 2003;33(7):1925-33. <https://doi.org/10.1002/eji.200323977>
- PMid:12884856
34. Lin J, Huang Y, Zhang L, Tang W, Li X, Wang X, *et al.* Evaluation of serum granulysin as a potential biomarker for nasopharyngeal carcinoma. *Clin Chim Acta Int J Clin Chem* 2016;454:72-6. <https://doi.org/10.1016/j.cca.2015.12.035>  
PMid:26751807
35. Thuong PH, Tam DB, Sakurada S, Le Hang NT, Hijikata M, Hong LT, *et al.* Circulating granulysin levels in healthcare workers and latent tuberculosis infection estimated using interferon-gamma release assays. *BMC Infect Dis* 2016;16:580. <https://doi.org/10.1186/s12879-016-1911-6>  
PMid:27756230
36. Su SC, Mockenhaupt M, Wolkenstein P, Dunant A, Le Gouvello S, Chen CB, *et al.* Interleukin-15 is associated with severity and mortality in Stevens-Johnson syndrome/toxic epidermal necrolysis. *J Invest Dermatol* 2017;137(5):1065-73. <https://doi.org/10.1016/j.jid.2016.11.034>  
PMid:28011147
37. Saigusa S, Ichikura T, Tsujimoto H, Sugawara H, Majima T, Kawarabayashi N, *et al.* Serum granulysin level as a novel prognostic marker in patients with gastric carcinoma. *J Gastroenterol Hepatol* 2007;22(8):1322-7. <https://doi.org/10.1111/j.1440-1746.2006.04796.x>  
PMid:17688669
38. Nagasawa M, Isoda T, Itoh S, Kajiwara M, Morio T, Shimizu N, *et al.* Analysis of serum granulysin in patients with hematopoietic stem-cell transplantation: Its usefulness as a marker of graft-versus-host reaction. *Am J Hematol* 2006;81(5):340-8. <https://doi.org/10.1002/ajh.20570>  
PMid:16628730
39. Saito N, Abe R, Yoshioka N, Murata J, Fujita Y, Shimizu H. Prolonged elevation of serum granulysin in drug-induced hypersensitivity syndrome. *Br J Dermatol* 2012;167(2):452-3. <https://doi.org/10.1111/j.1365-2133.2012.10921.x>  
PMid:22384988
40. Posadas SJ, Padial A, Torres MJ, Mayorga C, Leyva L, Sanchez E, *et al.* Delayed reactions to drugs show levels of perforin, granzyme B, and Fas-L to be related to disease severity. *J Allergy Clin Immunol* 2002;109(1):155-61. <https://doi.org/10.1067/mai.2002.120563>  
PMid:11799383
41. Saito N, Qiao H, Yanagi T, Shinkuma S, Nishimura K, Suto A, *et al.* An annexin A1-FPR1 interaction contributes to necroptosis of keratinocytes in severe cutaneous adverse drug reactions. *Sci Transl Med* 2014;6(245):245ra95. <https://doi.org/10.1126/scitranslmed.3008227>  
PMid:25031270