



# A Review of the Efficacy of the Dietary Intervention in Autism Spectrum Disorder

Zainab Taha<sup>1\*</sup>, Khalid A. Abdalhai<sup>2</sup>

<sup>1</sup>Department of Health Sciences, Zayed University, Abu Dhabi, United Arab Emirates; <sup>2</sup>Department of Psychiatry, International University of Africa, Sudan

## Abstract

**Edited by:** Ksenija Bogoeva-Kostovska  
**Citation:** Taha Z, Abdalhai KA. A review of the efficacy of the dietary intervention in autism spectrum disorder. Open Access Maced J Med Sci. 2021 Mar 18; 9(F):88-94. <https://doi.org/10.3889/oamjms.2021.5743>  
**Keywords:** Autism spectrum disorder; Vitamin supplementation; Dietary intervention  
**\*Correspondence:** Zainab Taha, Department of Health Sciences, Zayed University, Abu Dhabi, UAE. E-mail: [zainab.taha@zu.ac.ae](mailto:zainab.taha@zu.ac.ae)  
**Received:** 19-Jan-2021  
**Revised:** 07-Mar-2021  
**Accepted:** 08-Mar-2021  
**Copyright:** © 2021 Zainab Taha, Khalid A. Abdalhai  
**Funding:** This research did not receive any financial support  
**Competing Interest:** The authors have declared that no competing interest exists  
**Open Access:** This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

**BACKGROUND:** Autism spectrum disorder (ASD) is an early-onset neurodevelopmental disorder with 1 in 68 children prevalence. The key symptoms of ASD include social deficits, verbal and non-verbal communication deficits, and restricted, repetitive patterns of behavior, interests, or activities. For unknown reasons, gastrointestinal symptoms are related to ASD.

**AIM:** The aim of the current paper is to review the available literature on dietary interventions in children with ASD and provide up-to-date evidence.

**METHODS:** Searches in online databases, using the search terms 'dietary intervention', 'autism spectrum disorders', and 'autism' were conducted. The search targeted publications ranging from 2010 to 2020.

**RESULTS:** Twenty-six studies with different study designs which evaluated the effectiveness of specific dietary interventions in children with ASD were included in our review. Several dietary interventions were of popular practice in the management of children with ASD, including elimination (gluten-/casein-free diet and oligoantigenic diet), modification (modified ketogenic diet), supplementation (minerals, vitamins, omega 3, and omega 6), and exclusion (food additives). Studies showed different results with varying degrees of evidence. Overall, data were inadequate to make accurate conclusions.

**CONCLUSION:** There is little evidence to support the use of dietary interventions for children with ASD. Additional high-quality studies and further research are, therefore, needed.

## Background

Autism spectrum disorder (ASD) is a neurodevelopmental disorder that starts early in childhood and is characterized by impairment in reciprocal social communication and a tendency to engage in a repetitive, stereotyped pattern of behavior, interest, and activities [1]. The prevalence of ASD is increasing worldwide, with boys more affected than girls [2]. Recent studies estimated the prevalence of ASD to be 1 in 68 children, with as many as 50% of children showing behavioral regression around the age of three [3]. The etiological factors are heterogeneous and include genetic, epigenetic, and environmental factors; however, its pathogenesis remains unknown [1]. About 5%–15% of individuals with ASD have either a single gene disorder or chromosomal abnormality, and despite the link between ASD and some genetic disorders, for example, Fragile X syndrome and tuberous sclerosis, no specific gene has been identified [4]. Genetic factors are modulated by prenatal (e.g., exposed to antiepileptic drugs and advanced paternal age) [5], perinatal (e.g., infections during pregnancy and maternal autoimmune diseases) [6], and postnatal (e.g., prematurity and

obstetrical complications) [7] environmental factors in some patients.

It is apparent that, for unknown reasons, systemic clusters of symptoms are related to ASD, including gastrointestinal [GI] problems, mitochondrial dysfunction, and "allergic" symptoms, especially food intolerance and eczema [8]. Parents of children with ASD reported selective eating behavior in 58–67% of the cases [9].

With no known cure for ASD, most treatments have focused on improving the social, behavioral, and communicative symptoms. As a general principle, individuals with ASD require a comprehensive management approach that is individualized according to the child's age and specific needs. The variation and complexity of the presentation of children with ASD can lead to difficulties in determining the most effective interventions. At present, no medication treatments are approved for the core symptoms of ASD; however, some medications are approved by the Food Drug Administration for the associated irritability and aggression in children with ASD (i.e., risperidone and aripiprazole, [10]). A comprehensive multidisciplinary approach is recommended in the management of ASD in the practical clinical guidelines [11]. One of

the suggested elements of this approach is a dietary intervention, an available approach used with and without clinical supervision to alleviate GI and behavioral symptoms. Some authors have suggested that support for the nutritional and metabolic problems may be ameliorated by increasing nutrient intake, which may reduce the symptoms and comorbidities associated with ASD [12].

Accordingly, some therapists in clinical practice advise parents of children with ASD to follow certain dietary interventions, to improve some core or GI-related symptoms of ASD [13]. Several studies in the literature have investigated the effects of dietary interventions on ASD symptoms [14], [15], [16]. It would be beneficial for both clinicians and parents of children with ASD to receive evidence-based dietary interventions that address food preference and food selectivity in children with ASD.

The purpose of this paper is to provide up-to-date evidence for dietary interventions in children with ASD, including gluten-free and casein-free, ketogenic, and specific carbohydrate diets, as well as probiotics, polyunsaturated fatty acids, and dietary supplements (Vitamins A, C, B6, and B12; magnesium; and folate) by reviewing the available data. Focusing on medications is out of the scope of this paper.

## Methodology

### Methods

The researchers independently conducted a systematic search and review in two databases: PubMed and Google Scholar, using the search key terms “dietary intervention,” “autism spectrum disorders,” and “autism.” The reference lists of identified articles were also assessed to identify additional relevant articles. The search also included psychiatry textbooks and published and unpublished treatment guidelines. The targeted publications ranged from 2010 to 2020. A total of 26 articles were included and reviewed. The analyzed data were categorized into clinical and nutrition groups. Finally, the results were combined to ascertain information about the correlation between diet, dietary interventions, and ASD.

We developed inclusion criteria in consultation with an expert panel of clinicians, dietitians, and researchers. We included studies published in English of any design that examined relationships between diet, dietary interventions, and ASD.

The studies reviewed in this article used standardized criteria from DSM-IV, DSM-V, or ICD-10 (American Psychiatric Association 1994, 2013; World Health Organization 1993) to diagnose individuals with ASD.

## Results

In some individuals with ASD, early diagnosis of nutritional deficiencies and metabolic disorders and implementing appropriate therapeutic interventions have significantly improved cognitive abilities and behavioral difficulties [17], [18], [19]. Despite the scarcity of sufficient scientific and clinical evidence that supports the role of dietary interventions as part of ASD management [20]; it is still commonly recommended as a part of ASD management by some health providers [21]. These dietary interventions are based on either addition or elimination of certain food elements and nutrients associated with GI symptoms in children with ASD, such as inflammatory bowel disease, food intolerance, and allergies. Children with ASD tend to have food selectivity in comparison to other kids without neurodevelopmental disorders. Factors that may affect food selectivity in children with ASD include GI problems and “allergic” symptoms, especially food intolerance and eczema. Other factors, as reviewed by Whiteley *et al.*, include food texture (69%), food appearance (58%), food taste (45%), smell (36%), and food temperature (22%) [22]. This food selectivity can result in nutritional imbalance which may eventually be detrimental on brain function [23]. Hence, the problem of food selectivity needs to be carefully addressed through appropriate dietary interventions.

### Types of dietary interventions

In an effort to identify the role of dietary intervention in the management of ASD, researchers have investigated the GI tract (GIT)-related ASD symptoms. They found that protein digestive capacity may be impaired in some children with ASD, contributing to the abnormal immune activation and GI symptoms observed in these populations [24]. Studies have revealed decreased activity of several digestive enzymes as well as GI symptoms (loose stools and/or gaseousness), in children with ASD [25]. Other researchers have reported elevated levels of urinary peptides of dietary origin in children with ASD, providing further evidence that this population experiences impaired protein digestion, coupled with increased intestinal permeability [26]. Based on the results of the above-mentioned studies, researchers have found promising evidence to support the role of dietary interventions in children with ASD [24].

### Gluten-free/Casein-free (GFCF) diets

Despite little evidence to support positive effects, GFCF diets are considered the most common types of dietary interventions used in ASD [27]. The focus on GFCF diets, as an important dietary approach for children with ASD, originates from the evidence of

comorbid GIT problems related to consumption of foods containing casein and gluten [28], [29]. However, the explanation for increased GI problems in ASD remains unclear. One possible explanation that received attention could be related to increased intestinal permeability, the so-called leaky gut [30], [31]. Based on the “opioid excess theory,” digestion of gluten and casein results in the production of peptides with an opioid activity that can enter the bloodstream when the gut permeability is high [32], [33]. Therefore, these neuroactive peptides would, in turn, bind to opioid receptors and are ultimately speculated to affect processes in the central nervous system [34].

The use of a gluten-free diet necessitates the examination of the effects of removing all the food items containing gluten (the protein found in starch foods such as wheat, oats, barley, or rye). This dietary intervention is based on the exclusion of these food items from the diet and replaced by special gluten-free food items. Another dietary approach used for individuals with ASD, is the gluten-free diet combined with a casein-free diet. Casein is a peptide commonly found in milk and other dairy products. Consequently, a casein-free diet involves eliminating the intake of milk and dairy products. The main idea is to follow this elimination for a certain period and then reintroduce the products containing gluten/casein gradually to examine its contribution to ASD symptoms [35]. Studies that evaluated gluten-/casein-free diets reported some improvements in communication and challenging behaviors; however, the evidence was inconclusive [36], [37], [38].

### **Modified ketogenic, gluten-free diet regimen**

The ketogenic diet (KD) is a low-carbohydrate, moderate protein, high-fat diet that has emerged as a potential dietary intervention for ASD [39]. Some studies have supported the effectiveness of a modified ketogenic, gluten-free diet regimen with supplemental medium-chain triglyceride (MCT) as an effective dietary intervention for children with ASD. These studies support the notion that the KD alters neural cellular metabolism by utilizing ketone bodies as an alternative fuel for the brain [40].

Lee *et al.* reported moderate and substantial improvement in Autism Diagnostic Observation Scale-2 (approximately 20–50% improvement in the social affect domain) following the use of modified KD/GF/MCT for 3 months in children with ASD [39]. Accordingly, the authors have concluded that their results could support the suggestion that the KD components may offer an effective and safe treatment for children with ASD, especially for their social impairment. On the contrary, other researchers did not observe a significant correlation between beta-hydroxybutyrate “BHB” levels (a ketone synthesized in the liver from fatty acids when

the supply of glucose is too low for the body’s energetic needs, such as during periods of prolonged exercise, starvation, or absence of dietary sources) and the degree of improvement in ASD symptoms [41].

### **Role of vitamins and minerals in ASD**

Some health-care professionals consider vitamins, mineral supplements, and other non-pharmacological interventions to play a role in improving some ASD symptoms. There is some evidence that vitamins and mineral supplementations support the body’s basic physiologic processes and impact various metabolic and nutritional abnormalities (e.g., sulfation, methylation, glutathione redox imbalances, oxidative stress, and mitochondrial dysfunction). Nevertheless, there is no substantial evidence to support the efficacy of vitamin and mineral supplementations in improving children with ASD symptoms [42]. A recent study investigated the effect of Vitamin A deficiency on children with ASD and concluded Vitamin A deficiency exacerbates the core symptoms in children with ASD, especially when there are comorbid GI symptoms [43]. Vitamin C may have an effect on children with ASD, based on its vital role in brain neurons development, functional maturation, and antioxidant effect [44].

Hatice *et al.* studied the relationship between Vitamin D and its receptors, homocysteine, Vitamins B6, B12, and folate levels in 60 children with ASD and 45 children as the control group. The study observed that the total score of the Childhood Autism Rating Scale was positively linked to low serum levels of Vitamins D, B6, B12, folate, and Vitamin D receptors and high homocysteine levels [45]. This implies that these nutrients may be involved in the etiopathogenesis of ASD [46]. These findings may be of fundamental importance to care providers of children with ASD.

### **$\omega$ -3 and $\omega$ -6 fatty acid supplementation**

Researchers have also investigated the effect of supplementation with n–3 ( $\omega$ -3) including docosahexaenoic acid and n–6 ( $\omega$ -6) fatty acids including  $\gamma$ -linolenic acid on children born preterm who display early signs of ASD. They found that they helped promote cognitive development [47], [48]. They observed that children assigned to treatment with n–3 ( $\omega$ -3) and n–6 ( $\omega$ -6) fatty acids supplementation exhibited a significant reduction in ASD symptoms as per the Brief Infant-Toddler Social and Emotional Assessment scale, in comparison to the placebo group. However, the number of studies investigating the effectiveness of omega 3 supplements on alleviating the core symptoms of ASD was limited, and the overall effects were small, precluding definitive conclusions. Future large-scale randomized clinical trials are needed to confirm or refute the efficacy of omega 3 and 6 [49], [50].

### Food additives exclusion diet

Feingold, an American pediatrician and allergist, was the first to suggest that allergy to food additives, such as artificial flavors and colors, as well as naturally occurring salicylates, could contribute to symptoms of neurodevelopmental disorders [51]. It is believed that people with ASD cannot tolerate a range of additives, including aspartame, monosodium glutamate, and artificial colors (e.g., sunset yellow {E110}, tartrazine {E102}, carmoisine {E122}, Ponceau 4R {E124}), and sodium benzoate {E211}), evidenced by adverse effects on their behavior. While the avoidance of particular additives is very common, there has been little good quality research on the effect of food additives on patients with ASD [52].

### Oligoantigenic diet in ASD

Another dietary approach that has received considerable attention in children with ASD is the oligoantigenic diets. The basis for this diet is to mainly focus on eliminating suspected high allergenic food products rather than eliminating artificial colors, flavors, and preservatives precisely [35]. The oligoantigenic diet intervention, also known as the restricted elimination diet or hypoallergenic diet, follows a process that focuses on eliminating foods that are often highly allergenic, such as cow's milk, cheese, eggs, chocolate, and nuts. The oligoantigenic diet approach involves an elimination phase, typically 2–5 weeks, where specific food items would be completely excluded. The food items allowed in the elimination phase could consist of only a few hypoallergenic foods such as rice, turkey, lettuce, pears, and water [53]. The patient would then be monitored for a substantial decrease in ASD symptoms, which would indicate "food sensitivity." This would be followed by the reintroduction phase, which involves the introduction of specific food items and observation for triggering of symptoms. In total, the assessment period could take up to 18 months. Therefore, the elimination/reintroduction of diet serves as a "diagnostic" tool for identifying the specific food items responsible for the physical and/or behavioral disorders in children with ASD.

Jose *et al.* reviewed the most relevant available data for the dietary and nutritional interventions in children diagnosed with ASD. Table 1 summarizes the level of evidence to each dietary intervention that has been addressed in the current review.

acid supplementations, food additives, exclusion diet, and the oligoantigenic diet. Despite the variation in the results of the studies conducted on dietary interventions and supplements in children with ASD, the interventions long-term effects are not well addressed in the research. The Center for Disease Control and Prevention stresses that "while remedies that work for one child may not work for another, dietary changes or supplements are a worthwhile option to explore [55]." Many families of children with ASD use dietary interventions and supplements, even without clear evidence on safety and efficacy. Therefore, a comprehensive multidisciplinary team is required and should include families of autistic children in the management plan with particular advice on the risk and benefits of using dietary intervention and supplements. Child and adolescents' psychiatrists as well as other mental health-care professionals must receive training in nutritional principles for patients with ASD and other neurodevelopmental disorders.

**Table 1: Levels of evidence of complementary and alternative treatments in children diagnosed with ASD\***

Treatment	Evidence	Comments	Grade of evidence
Gluten-free diet/casein-free diet	Single-blind trials suggested a potential benefit in 5–7-year-old children with GI symptoms A double-blind trial does not show benefit	Need for studies to clarify the utility Risk of nutritional involvement Supervision by a specialist in nutrition	B
Omega-3 fatty acids	Several systematic reviews Studies report benefits but with many methodological problems	Absence of high-quality evidence that shows effectiveness on the core and associated symptoms ASD Need for more studies based on promising effects in other populations	C
Vitamins/minerals/supplements Vitamin A	A double-blind, placebo-controlled randomized trials Theories	Significant Significant possibility of harm	C
Vitamin C	No evidence of effectiveness Two double-blind, placebo-controlled trials	Some preliminary evidence Non-significant toxicity	B
Vitamin D	Circumstantial evidence-based treatment 2005 Cochrane review: No recommendation	Theories Methodological problems	D
Vitamins B6 and magnesium	Update 2010: Same conclusion	Poor quality of the studies No recommendation Potential neurotoxicity of B6 and/or magnesium	D
Vitamin B12	A small open pilot study No additional studies since 2008 review Twelve-week double-blind, placebo-controlled study Improvement in a subgroup	Need for more studies to define responders' group	C+

A: More than 1 high-quality study with consistent results or an extensive multicenter study. B: A high-quality study or several studies with mild limitations. C: A study with important limitations. D: No evidence/theories/multiple studies with very serious limitations. \*Adapted from Levy SE, Hyman SL [54]. ASD: Autism spectrum disorder.

## Discussion

This review summarizes different dietary interventions for ASD, including gluten-free and casein-free, KDs, vitamins and minerals, omega 3 and 6 fatty

## Conclusion

Despite many research efforts investigating the effects of dietary interventions in ASD, our review shows no consensus exists to support specific dietary



interventions and supplementations in this patient population. More high-quality, longitudinal studies are needed to investigate the effects of dietary interventions and supplements in children with ASD. Future guidelines should be more explicit concerning dietary interventions and supplementations for this complex and impairing condition.

## Review Limitations

The number of studies investigating dietary interventions and supplementation in children with ASD is variable in their designs and results. ASD symptoms were not clearly described in some of these studies.

## Acknowledgments

We appreciate the time and work undertaken by Dr. Teresa Arora for revising and editing the manuscript.

## References

- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders: DSM-5™. 5<sup>th</sup> ed. Arlington, VA: American Psychiatric Association; 2013. [https://doi.org/10.1007/springerreference\\_179660](https://doi.org/10.1007/springerreference_179660)
- Al-Mamri W, Idris AB, Dakak S, Al-Shekaili M, Al-Harathi Z, Alnaamani AM, *et al.* Revisiting the prevalence of autism spectrum disorder among Omani children: A multicentre study. *Sultan Qaboos Univ Med J.* 2019;19:e305. <https://doi.org/10.18295/squmj.2019.19.04.005>  
PMid:31897313
- Zablotsky B, Black LI, Maenner MJ, Schieve LA, Blumberg SJ, *et al.* Estimated prevalence of autism and other developmental disabilities following questionnaire changes in the 2014 National Health Interview Survey. *Natl Health Stat Report.* 2015;87:1-20. PMid:26632847
- Devlin B, Scherer SW. Genetic architecture in autism spectrum disorder. *Curr Opin Genet Dev.* 2012;22(3):229-37. PMid:22463983
- Rasalam A, Hailey H, Williams JH, Moore SJ, Turnpenny PD, Lloyd DJ, *et al.* Characteristics of fetal anticonvulsant syndrome associated autistic disorder. *Dev Med Child Neurol.* 2005;47(8):551-5. <https://doi.org/10.1017/s0012162205001076>  
PMid:16108456
- Xiang AH, Wang X, Martinez MP, Page K, Buchanan TA, Feldman RK, *et al.* Maternal Type 1 diabetes and risk of autism in offspring. *JAMA.* 2018;320(1):89-91. <https://doi.org/10.1001/jama.2018.7614>  
PMid:29936530
- Newschaffer CJ, Croen LA, Daniels J, Giarelli E, Grether JK, Levy SE, *et al.* The epidemiology of autism spectrum disorders. *Annu Rev Public Health* 2007;28:235-58. <https://doi.org/10.1146/annurev.publhealth.28.021406.144007>  
PMid:17367287
- Trajkovski V. Health Condition in Persons with Autism Spectrum Disorders. *J ReAttach Ther Dev Divers.* 2019;1(2):112-24.
- Cermak SA, Curtin C, Bandini LG. Food selectivity and sensory sensitivity in children with autism spectrum disorders. *J Am Dietetic Assoc.* 2010;110(2):238-46. <https://doi.org/10.1016/j.jada.2009.10.032>  
PMid:20102851
- Ghaleiha A, Asadabadi M, Mohammadi MR, Shahei M, Tabrizi M, Hajiaghvae R, *et al.* Memantine as adjunctive treatment to risperidone in children with autistic disorder: A randomized, double-blind, placebo-controlled trial. *Int J Neuropsychopharmacol.* 2013;16(4):783-9. <https://doi.org/10.1017/s1461145712000880>  
PMid:22999292
- Volkmar F, Siegel M, Woodbury-Smith M, King B, McCracken J, State M, *et al.* Practice parameter for the assessment and treatment of children and adolescents with autism spectrum disorder. *J Am Acad Child Adolesc Psychiatry* 2014;53:237-57. <https://doi.org/10.1016/j.jaac.2013.10.013>  
PMid:24472258
- Kaluźna-Czaplińska J, Jóźwik-Pruska J. Nutritional strategies and personalized diet in autism-choice or necessity? *Trends Food Sci Technol* 2016;49:45-50. <https://doi.org/10.1016/j.tifs.2016.01.005>
- Rubenstein E, Schieve L, Bradley C, DiGiuseppi C, Moody E, Thomas K, *et al.* The prevalence of gluten free diet use among preschool children with autism spectrum disorder. *Autism Res* 2018;11:185-93. <https://doi.org/10.1002/aur.1896>  
PMid:29155492
- Doenyaş C. Dietary interventions for autism spectrum disorder: New perspectives from the gut-brain axis. *Physiol Behav* 2018;194:577-82. <https://doi.org/10.1016/j.physbeh.2018.07.014>  
PMid:30036560
- Ristori MV, Quagliarello A, Reddel S, Ianiro G, Vicari S, Gasbarrini A, *et al.* Autism, gastrointestinal symptoms and modulation of gut microbiota by nutritional interventions. *Nutrients.* 2019;11(11):2812. <https://doi.org/10.3390/nu11112812>  
PMid:31752095
- Siafis S, Çıray O, Schneider-Thoma J, Bighelli I, Krause M, Rodolico A, *et al.* Placebo response in pharmacological and dietary supplement trials of autism spectrum disorder (ASD): Systematic review and meta-regression analysis. *Mol Autism.* 2020;11(1):1-19. <https://doi.org/10.1186/s13229-020-00372-z>  
PMid:32847616
- Karhu E, Zukerman R, Eshraghi RS, Mittal J, Deth RC, Castejon AM, *et al.* Nutritional interventions for autism spectrum disorder. *Nutr Rev.* 2020;78(7):515-31. <https://doi.org/10.1093/nutrit/nuz092>  
PMid:31876938
- Gogou M, Kolios G. Are therapeutic diets an emerging additional choice in autism spectrum disorder management? *World J Pediatr.* 2018;14(3):215-23. <https://doi.org/10.1007/s12519-018-0164-4>  
PMid:29846886
- Cekici H, Sanlier N. Current nutritional approaches in managing autism spectrum disorder: A review. *Nutr Neurosci* 2019;22(3):145-55. <https://doi.org/10.1080/1028415x.2017.1358481>

- PMid:28762296
20. Faras H, Al Ateeqi N, Tidmarsh L. Autism spectrum disorders. *Ann Saudi Med.* 2010;30(4):295-300. <https://doi.org/10.4103/0256-4947.65261>  
PMid:20622347
  21. Marti LF. Dietary interventions in children with autism spectrum disorders-an updated review of the research evidence. *Curr Clin Pharmacol.* 2014;9(4):335-49. <https://doi.org/10.2174/15748847113086660074>  
PMid:24050740
  22. Whiteley P, Rodgers J, Shattock P. Feeding Patterns in Autism. *Autism.* 2000;4:207-11. <https://doi.org/10.1177/1362361300004002008>
  23. Emond A, Emmett P, Steer C, Golding J. Feeding symptoms, dietary patterns, and growth in young children with autism spectrum disorders. *Pediatrics.* 2010;126(2):e337-42. <https://doi.org/10.1542/peds.2009-2391>  
PMid:20643716
  24. Sanctuary MR, Kain JN, Angkustsiri K, German JB. Dietary considerations in autism spectrum disorders: The potential role of protein digestion and microbial putrefaction in the gut-brain axis. *Front Nutr.* 2018;5:40. <https://doi.org/10.3389/fnut.2018.00040>  
PMid:29868601
  25. Horvath K, Papadimitriou JC, Rabsztyl A, Drachenberg C, Tildon JT. Gastrointestinal abnormalities in children with autistic disorder. *J Pediatr.* 1999;135(5):559-63. [https://doi.org/10.1016/s0022-3476\(99\)70052-1](https://doi.org/10.1016/s0022-3476(99)70052-1)  
PMid:10547242
  26. Reichelt KL, Knivsberg AM. Can the pathophysiology of autism be explained by the nature of the discovered urine peptides? *Nutr Neurosci.* 2003;6(1):19-28. <https://doi.org/10.1080/1028415021000042839>  
PMid:12608733
  27. Marsden RE, Francis J, Garner I. Use of GFCF diets in children with ASD. An investigation into parents' beliefs using the theory of planned behaviour. *J Autism Dev Disord.* 2019;49(9):3716-31. <https://doi.org/10.1007/s10803-019-04035-8>  
PMid:31165958
  28. Moon K, Rao SC, Schulzke SM, Patole SK, Simmer K, *et al.* Longchain polyunsaturated fatty acid supplementation in preterm infants. *Cochrane Database Syst Rev.* 2016;12:Cd000375. <https://doi.org/10.1002/14651858.cd000375.pub3>  
PMid:27995607
  29. Coury DL, Ashwood P, Fasano A, Fuchs G, Geraghty M, Kaul A, Mawe G, *et al.* Gastrointestinal conditions in children with autism spectrum disorder: Developing a research agenda. *Pediatrics.* 2012;130(Suppl 2):S160-8. <https://doi.org/10.1542/peds.2012-0900n>  
PMid:23118247
  30. de Magistris L, Familiari V, Pascotto A, Sapone A, Froli A, Iardino P, *et al.* Alterations of the intestinal barrier in patients with autism spectrum disorders and in their first-degree relatives. *J Pediatr Gastroenterol Nutr.* 2010;51(4):418-24. <https://doi.org/10.1097/mpg.0b013e3181dccc4a5>  
PMid:20683204
  31. Strati F, Cavalieri D, Albanese D, De Felice C, Donati C, Hayek J, *et al.* New evidences on the altered gut microbiota in autism spectrum disorders. *Microbiome.* 2017;5(1):24. <https://doi.org/10.1186/s40168-017-0242-1>  
PMid:20683204
  32. Lazaro CP, Ponde MP, Rodrigues LE. Opioid peptides and gastrointestinal symptoms in autism spectrum disorders. *Braz J Psychiatry.* 2016;38(3):243-6. <https://doi.org/10.1590/1516-4446-2015-1777>  
PMid:27304256
  33. Shattock P, Kennedy A, Rowell F, Berney T. Role of neuropeptides in autism and their relationships with classical neurotransmitters. *Brain Dysfunct.* 1990;3((5-6)):328-45.
  34. Shattock P, Whiteley P. Biochemical aspects in autism spectrum disorders: Updating the opioid-excess theory and presenting new opportunities for biomedical intervention. *Expert Opin Ther Targets.* 2002;6(2):175-83. <https://doi.org/10.1517/14728222.6.2.175>  
PMid:12223079
  35. Ly V, Bottelier M, Hoekstra PJ, Vasquez AA, Buitelaar JK, Rommelse NN, *et al.* Elimination diets' efficacy and mechanisms in attention deficit hyperactivity disorder and autism spectrum disorder. *Eur Child Adolesc Psychiatry.* 2017;26(9):1067-79. <https://doi.org/10.1007/s00787-017-0959-1>  
PMid:28190137
  36. Çöpür M, Çöpür S. Autism spectrum disorders and gluten/casein free diet treatment: A systematic review (1990-2016). *Int J Nutr.* 2017;2(4):7. <https://doi.org/10.14302/issn.2379-7835.ijn-17-1710>
  37. Sathe N, Andrews JC, McPheeters ML, Warren ZE. Nutritional and dietary interventions for autism spectrum disorder: A systematic review. *Pediatrics.* 2017;139:e20170346. <https://doi.org/10.1542/peds.2017-0346>  
PMid:28562286
  38. Tye C, Runicles AK, Whitehouse AJ, Alvares GA, *et al.* Characterizing the interplay between autism spectrum disorder and comorbid medical conditions: An integrative review. *Front Psychiatry.* 2019;9:751. <https://doi.org/10.3389/fpsy.2018.00751>  
PMid:30733689
  39. Lee RW, Corley MJ, Pang A, Arakaki G, Abbott L, Nishimoto M, *et al.* A modified ketogenic gluten-free diet with MCT improves behavior in children with autism spectrum disorder. *Physiol Behav.* 2018;188:205-11. <https://doi.org/10.1016/j.physbeh.2018.02.006>  
PMid:29421589
  40. Stafstrom CE, Rho JM. The ketogenic diet as a treatment paradigm for diverse neurological disorders. *Front Pharmacol.* 2012;3:59. <https://doi.org/10.3389/fphar.2012.00059>  
PMid:22509165
  41. Newman JC, Verdin E. Beta-hydroxybutyrate: A signaling metabolite. *Annu Rev Nutr.* 2017;37:51-76. <https://doi.org/10.1146/annurev-nutr-071816-064916>  
PMid:28826372
  42. Bjorklund G, Waly MI, Al-Farsi Y, Saad K, Dadar M, Rahman M, *et al.* The role of vitamins in autism spectrum disorder: What do we know? *J Mol Neurosci.* 2019;67(3):373-87. <https://doi.org/10.1007/s12031-018-1237-5>  
PMid:30607900
  43. Cheng B, Zhu J, Yang T, Guo M, Lai X, Li Q, *et al.* Vitamin A deficiency increases the risk of gastrointestinal comorbidity and exacerbates core symptoms in children with autism spectrum disorder. *Pediatr Res.* 2021;89(1):211-6. <https://doi.org/10.1038/s41390-020-0865-y>  
PMid:32225174
  44. Qiu S, Li L, Weeber EJ, May JM. Ascorbate transport by primary cultured neurons and its role in neuronal function and protection against excitotoxicity. *J Neurosci Res.* 2007;85(5):1046-56. <https://doi.org/10.1002/jnr.21204>  
PMid:17304569
  45. Altun H, Kurutaş EB, Şahin N, Güngör O, Fındıklı E. The levels of Vitamin D, Vitamin D receptor, homocysteine and complex B Vitamin in children with autism spectrum disorders. *Clin Psychopharmacol Neurosci.* 2018;16(4):383. <https://doi.org/10.9758/cpn.2018.16.1.383>

- PMid:30466210
46. Altun H, Kurutaş EB, Şahin N, Güngör O, Fındıklı E. The levels of Vitamin D, Vitamin D receptor, homocysteine and complex B Vitamin in children with autism spectrum disorders. *Clin Psychopharmacol Neurosci*. 2018;16(4):383. <https://doi.org/10.9758/cpn.2018.16.1.383>  
PMid:30466210
47. Keim SA, Gracious B, Boone KM, Klebanoff MA, Rogers LK, Rausch J, *et al*. Omega-3 and omega-6 fatty acid supplementation may reduce autism symptoms based on parent report in preterm toddlers. *J Nutr*. 2018;148(2):227-35. <https://doi.org/10.1093/jn/nxx047>  
PMid:29490101
48. Keim SA, Boone KM, Klebanoff MA, Turner AN, Rausch J, Nelin MA, *et al*. Effect of docosahexaenoic acid supplementation vs placebo on developmental outcomes of toddlers born preterm: A randomized clinical trial. *JAMA Pediatr*. 2018;172(12):1126-34. <https://doi.org/10.1001/jamapediatrics.2018.3082>  
PMid:30357263
49. Cheng YS, Tseng PT, Chen YW, Stubbs B, Yang WC, Chen TY, *et al*. Supplementation of omega 3 fatty acids may improve hyperactivity, lethargy, and stereotypy in children with autism spectrum disorders: A meta-analysis of randomized controlled trials. *Neuropsychiatr Dis Treat*. 2017;13:2531-43. <https://doi.org/10.2147/ndt.s147305>  
PMid:29042783
50. James S, Montgomery P, Williams K. Omega-3 fatty acids supplementation for autism spectrum disorders (ASD). *Cochrane Database Syst Rev*. 2011;9(11):Cd007992. <https://doi.org/10.1002/14651858.cd007992.pub2>  
PMid:22071839
51. Feingold BF. Hyperkinesis and learning disabilities linked to artificial food flavors and colors. *Am J Nurs*. 1975;75(5):797-803. <https://doi.org/10.1097/00000446-197505000-00021>  
PMid:1039267
52. INDI-Irish Nutrition and Dietetic Institute. *Autism Spectrum Disorders and Diet in Children*. Ashgrove House, Kill Avenue: Dun Laoghaire, Co., Dublin; 2010. Available from: <https://www.indi.ie/diseases,-allergies-and-medical-conditions/disability/361-diet-and-autism.html>.
53. PelslerLM, FrankenaK, ToormanJ, SavelkoulHF, DuboisAE, Pereira RR, *et al*. Effects of a restricted elimination diet on the behaviour of children with attention-deficit hyperactivity disorder (INCA study): A randomised controlled trial. *Lancet*. 2011;377(9764):494-503. [https://doi.org/10.1016/s0140-6736\(10\)62227-1](https://doi.org/10.1016/s0140-6736(10)62227-1)  
PMid:21296237
54. Levy SE, Hyman AL. Complementary and alternative medicine treatments for children with autism spectrum disorders. *Child Adolesc Psychiatr Clin*. 2015;24(1):117-43.  
PMid:25455579
55. Centers for Disease Control and Prevention. *Treatment and Intervention Services for Autism Spectrum Disorder*. Atlanta, Georgia: Centers for Disease Control and Prevention; 2015.