



Morphofunctional and Hematological Characteristics of Health in Students from the Northern and Southern Regions of Kazakhstan

Saule Bazarbaeva^{1,2}, Aigul Dinmukhamedova^{2*}, Gulnara Tleubergenova¹, Zhanar Rakhimzhanova³, Kamila Sembekova³, Sholpan Karbayeva⁴, Elmira Kuandykova⁴

¹Department of Biology, M. Kozybayev North-Kazakhstan State University, Petropavlovsk, Kazakhstan; ²Department of General Biology and Genomics, L.N. Gumilyov Eurasian National University, Nur-Sultan, Kazakhstan; ³Department of Normal Physiology, Astana Medical University, Nur-Sultan, Kazakhstan; ⁴Department of Geography, Ecology and Tourism, Abai Kazakh National Pedagogical University, Almaty, Kazakhstan

Abstract

BACKGROUND: Over the past decade, the Republic of Kazakhstan has undergone various socio-economic transformations, which have revealed many negative factors affecting the health and quality of life of the population, especially the younger generation.

AIM: The study aim is to study the morphofunctional and hematological parameters of the 1st year university students.

METHODS: The health status for a total of 400 1st-year students, who arrived for studying from the northern regions (North Kazakhstan, Petropavlovsk) and the southern regions (Turkistan, Kyzylorda), was evaluated during the study. The study investigated some morphological and functional parameters of the organism (body length, body mass index, chest circumference, vital lung capacity, vital index, blood pressure, heart rate, etc.), and hematological indicators (the total number of red blood cells, white blood cells, the concentration of hemoglobin in the blood, etc.). The results were processed by means of variational and differential statistics.

RESULTS: Comparing the physical health of students from both regions, a significant difference in main morphofunctional and hematological indices has been established. The functional indicators of the cardiovascular and respiratory systems were higher in students from the southern regions, which reflected a good adaptive reaction to studying at the university, as well as new climatic and geographical conditions. Students from the southern regions were reported to have lower red blood cell amounts and mild anemia compared to those from the northern regions.

CONCLUSION: Therefore, constant monitoring of the health state among youth is required to develop effective tactics for the students' smooth adjusting to living in new social and climatic conditions.

Edited by: Sinisa Stojanoski
Citation: Bazarbaeva S, Dinmukhamedova A, Tleubergenova G, Rakhimzhanova Z, Sembekova K, Karbayeva S, Kuandykova E. Morphofunctional and hematological characteristics of health in students from the Northern and Southern regions of Kazakhstan. Open Access Maced J Med Sci. 2021 Sep 11; 9(A):753-759. <https://doi.org/10.3889/oamjms.2021.6434>
Keywords: Adaptation; Cardio-respiratory system; Functional characteristics; Morphological characteristics; Physical development
***Correspondence:** Aigul Dinmukhamedova, L.N. Gumilyov Eurasian National University, 13 Kazhymukan Munaipasov Street, 010000 Nur-Sultan, Kazakhstan. Phone: +77019181131. E-mail: dinmukhamedova2020@rambler.ru
Received: 12-May-2021
Revised: 23-Jun-2021
Accepted: 01-Sep-2021
Copyright: © 2021 Saule Bazarbaeva, Aigul Dinmukhamedova, Gulnara Tleubergenova, Zhanar Rakhimzhanova, Kamila Sembekova, Sholpan Karbayeva, Elmira Kuandykova
Funding: This research did not receive any financial support
Competing Interest: The authors have declared that no competing interest exists
Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Introduction

In recent decades, different socio-economic transformations that occurred in the Republic of Kazakhstan revealed many negative phenomena conceivably affecting the state of health and quality of life of the population. This particularly concerns the deterioration of health among the younger generation [1], [2]. It has been shown that the pace of morphological and functional development in youth and their health state depend not only on the gene pool but to a large extent on environmental factors [3], [4].

Residents of various climatic-geographic zones differ in morphological parameters, as well as by the specifics of metabolic processes in each region [5], [6].

The Government of the Republic of Kazakhstan has adopted and is implementing the Serpin-2050 social project, according to which preferential state educational grants are provided for applicants from the

southern regions at universities in Northern Kazakhstan. This project can allow equalizing the regional imbalance in the employment market, which is caused by low birth rates and external migration in the northern part of the country [7].

At the same time, some researchers note that the process of adaptation of an average healthy person to the climatic and social conditions is accompanied by fairly pronounced changes in all systems involved in the oxygen supply of tissues. Initial changes at the biochemical, cellular, and tissue levels are manifested afterward in the functional changes of the respiratory system, blood circulation, and red blood cells. The blood circulation system plays a dominant role in maintaining homeostasis in constantly changing environmental conditions and is one of the first to be included in the adaptation process [8], [9], [10], [11], [12].

Based on the foregoing, research was undertaken to investigate the morphofunctional and hematological indicators of the health state among

1st-year students, who came to study from the northern and southern regions of the Republic.

Methods

The study was conducted on the basis of M. Kozybaev North Kazakhstan State University in September of 2018–2019 academic year. The study included 17–18 years old 1st-year students (200 girls and 200 boys of Kazakh nationality). The participants were selected by age, gender, and place of residence (north, south); residence in the region from birth to the present time was taken into account. All participants belonged to health Groups 1 and 2; in other cases, participants were excluded from the study. The study design included the assessment of morphological, physiological, and hematological characteristics of 1st-year students. The morphological and functional indicators of the students' health were studied in the medical office of the university using standard equipment (height meter, scales, portable dry spirometer SPP, tonometer, and electrocardiograph "Axion EK 1T-07").

Anthropometric parameters were evaluated by BL, BM, and chest circumference (CHC). According to obtained BL, BM, and CHC values, the body mass index (BMI), stenium (SI), and Pinier (PI) indices were calculated [13]. The type of constitution was determined according to Chernorutsky [14].

The indexes of stenium and Pinier were used to assess individual constitutional type.

$$SI = BL, \text{ cm} / (2 \times BM, \text{ kg} + CHC, \text{ cm}) \quad (1)$$

The SI value above 1.25 indicates the predominance of the dolichomorphic body type, SI within the range of 0.85–1.25 expresses the mesomorphic type, and below 0.85 assumes the brachymorphic type [15].

$$PI = BL, \text{ cm} - (BM, \text{ kg} + CHC, \text{ cm}) \quad (2)$$

When assessing the Pinier index, the value above 30 is typical for asthenic body type, within the range from 10 to 30 characterizes the normostenic body type, and below ten assumes that of hypersthenic [14].

For the experiment, the following functional parameters were determined:

1. The strength of the hands (SH) with the help of dynamometer and hand force index (HFI) by the formula:

$$HFI = SH \text{ (kg)} / BM \text{ (kg)} \quad (3)$$
2. Vital lung capacity (VC) with the help of air spirometer and vital index (VI) by the formula:

$$VI = VC \text{ (ml)} / BM \text{ (kg)} \quad [15].$$

To assess the individual value of VC, it was compared with VC (ShVC) based on indicators of body

length (BL) and body mass (BM). The ShVC (ml) was calculated by the following formulas:

$$\text{for males: } ShVC = 40 \times BL \text{ (cm)} + 30 \times BM \text{ (kg)} - 4400 \quad (4)$$

$$\text{for females: } ShVC = 40 \times BL \text{ (cm)} + 10 \times BM \text{ (kg)} - 3800 \quad (5)$$

The functional efficacy of the cardiovascular system in conditions of relative rest was evaluated by the double product (DP):

$$DP = (SBP \times HR) / 100 \quad (6)$$

where SBP is systolic blood pressure (BP) in mm Hg, and HR is the heart rate in bpm [14], [15].

To assess the functioning level of the blood circulatory system, the adaptive potential (AP) was applied [14].

Calculation of AP (in points) was carried out according to the formula:

$$AP = 0.011 (HR) + 0.014 (SBP) + 0.008 (DBP) + 0.014 (A) + 0.009 (BM) - 0.009 (BL) + 0.004 (G) - 0.273 \quad (7)$$

where HR is the heart rate (HR), bpm; SBP; and DBP are systolic and diastolic BP, respectively, mmHg; A is the age, years; BM is the body mass, kg; BL is the body length, cm; and G is the gender (male –1 and female –2). The results were evaluated by the following parameters: satisfactory adaptation - no more than 2.10 points, adaptation stress - from 2.11 to 3.2, unsatisfactory adaptation - from 3.21 to 4.3, and failure of adaptation - from 4.31 and higher [16].

Assessment of the main characteristics of peripheral blood was carried out based on the Regional Blood Center of Petropavlovsk. Counting and differentiation of the formed elements of peripheral blood were performed by an automatic hematological analyzer "Sysmex XS-1000i" (Sysmex Corporation, Japan). Blood sampling was taken from all study participants in the morning on an empty stomach. The analysis of peripheral blood parameters included the total number of red blood cells, white blood cells, and the concentration of hemoglobin in the blood. Erythrocyte indices were examined as well, namely, the average corpuscular volume of an erythrocyte and the average content and concentration of hemoglobin in an erythrocyte.

The results were processed by means of variational and differential statistics applying non-parametric Student's criteria for independent samples at a significance level of $p \leq 0.05$.

All students gave their voluntary consent to take part in the study. The study design was adopted by Ethical Committee of M. Kozybayev North Kazakhstan State University (protocol No. 6 of 27 Jun 2018).

Results

Anthropometric indicators largely depended on the place of previous residence. Students of both sexes from the southern regions had higher body mass and CHC indices, while the BL value was higher for northern residents in all studied groups (Table 1).

Table 1: Anthropometric parameters of students from the northern and southern regions

Indicators	Women of the northern regions (n = 100)	Women of the southern regions (n = 100)	Men of the northern regions (n = 100)	Men of the southern regions (n = 100)
Height, cm	162.3 ± 0.4	160.1 ± 0.3 [#]	172.1 ± 0.4*	171.8 ± 0.3*
Weight, kg	56.8 ± 0.6	60.2 ± 0.8 [#]	71.1 ± 0.4*	73.8 ± 0.4* [#]
CHC, cm	82.9 ± 0.5	88.06 ± 0.6 [#]	91.8 ± 0.4*	95.04 ± 0.4* [#]

*0.05, significant differences between the students of different sex. [#]Significant differences between the students of different regions. CHC: Chest circumference.

A qualitative assessment of anthropometric parameters was given based on the BMI. An analysis of the BMI showed that male and female students from the southern regions had a higher rate than those from northern parts of the country.

Moreover, the BMI in students from the southern regions corresponded to "increased fatness," and in the inhabitants of the northern regions, this indicator had the normal value (Table 2).

Table 2: Indices of students' physical development

Indicators	Women of the northern regions (n = 100)	Women of the southern regions (n = 100)	Men of the northern regions (n = 100)	Men of the southern regions (n = 100)
Body mass index, kg/m ²	21.6 ± 0.2	23.5 ± 0.3 [#]	24.02 ± 0.1*	25.02 ± 0.1* [#]
PI, c.u	22.5 ± 1.01	11.8 ± 1.3 [#]	9.1 ± 0.8*	2.9 ± 0.7* [#]

*0.05 Significant differences between the students of different sex. [#]Significant differences between the students from different regions. PI: Pinier index.

The Pinier index, which characterizes the type of physique of a person, was higher in the group of students from the northern regions compared to representatives of the southern regions. Young men in both groups had a hypersthenic type of constitution, while the women of the northern regions had the Pinier index values consistent with the normosthenic type, while the students from the southern regions tended to have hypersthenia (Table 3).

Table 3: Distribution of students by body weight (%)

Group	Weight loss		Standard		Weight excess	
	Women	Men	Women	Men	Women	Men
North	10	0	77	66	13	24
South	8	0	58	71	34	29

It is shown in Table 3, among the women of the southern regions, persons with excess body weight prevailed, while the majority of women from the northern regions had the normal body weight. In young men, a lack of body weight was not observed, and an excessive body weight was observed in 24% of young men of northern and 29% of the southern regions.

An assessment of the wrist dynamometry among students depending on the place of residence showed that men and women from the southern regions of the Republic had higher indicators of the strength of their right and left hands (Table 4).

Table 4: Functional parameters of body

Indicators	Women of the northern regions (n = 100)	Women of the southern regions (n = 100)	Men of the northern regions (n = 100)	Men of the southern regions (n = 100)
HFI, kg/kg	55.1 ± 1.07	64.03 ± 1.1 [#]	70.6 ± 0.9*	72.7 ± 0.7*
VI, ml/kg	52.6 ± 0.8	57.1 ± 1.1 [#]	59.1 ± 0.9*	62.7 ± 1.04* [#]
Adaptive potential, c.u	1.8 ± 0.01	2.1 ± 0.02 [#]	2.04 ± 0.01*	2.2 ± 0.01* [#]

*0.05 Significant differences between the students examined by gender. [#]Significant differences between the students examined by regions. HFI: Hand force index, VI: Vital index.

Young men who came to study from the southern regions had higher absolute values of hand dynamometry that indicated the higher functional capabilities of the muscular system. Since the absolute values of wrist dynamometry cannot give a complete understanding of functional differences, the calculated wrist index was applied, which showed the relation between the strength of the hands and the body weight. According to the carpal index, students of the southern regions also had higher rates compared to those from the northern regions.

When examining the indicators of the respiratory system functional state, the value of vital capacity for students in the northern regions was noted to be generally lower than the age-appropriate due values, while for the students from the southern regions, the VC indicators were normal (Figure 1).




Figure 1: Functional state of the respiratory system of students from the northern and southern regions. *≤0.05 significant differences between students from the northern and southern regions; (a) Women; (b) Men; VC: Vital capacity, ShVC: Should vital capacity

During the study, it has been established that in students from the southern regions, the HR at rest, as well as SBP and DBP values were higher than similar indicators of people living in the north of the Republic. Thus, BP indicators express the regional differences, which can be attributed to a higher stress level of the functional state of the cardiovascular system in the process of students' adaptation to new climatic and geographical conditions (Figure 2).

Given the indicators of physical development, the level of physical health, and functional reserves of the body were estimated. More than half of the 1st-year students from both regions had an average level of physical health, namely, for the northern regions - 59% female and 47% male students, and for the southern regions - 44% female and 49% male students. At that, the level of health in about 30% of women and 19%




Figure 2: Indicators of the cardiovascular system functioning in students from the northern and southern regions. * ≤ 0.05 Significant differences between students from the northern and southern regions; (a) Women; (b) men; HR: Heart rate, SBP: Systolic blood pressure, DBP: Diastolic blood pressure

of men from the southern regions was stated as below average.

Among the examined students, low health prevailed among young men of the southern regions and amounted to 27%. In general, when evaluating the level of physical health of the 1st-year students according to the method of Apanasenko, students of the northern regions was determined to have an average level of physical health, and for those from the southern regions of the republic, a lower level of health was noted.

The results of studies of blood parameters in students from different climatic and geographical regions are shown in Table 5. It has been established that all the obtained indicators were within referential ranges. However, when comparing the obtained results, some significant differences were noticed. Thus, in residents of the northern regions, the red blood cell counts were higher in comparison with representatives of the south. The hemoglobin content in the blood of students was reported as normal. Although, among women from the southern regions, a deviation from reference values was observed and the hemoglobin level was 108.2 ± 2.7 g/l. Studies conducted in the Kyzylorda region among women between 18 and 45-years-old revealed iron deficiency anemia in 40.2% of the examined females. This value was significantly higher than that found among women in Europe (14%), but below-average values among African (48%) and South Asian women (57%) [17], [18].

The prevalence of anemia in Kazakhstan among women of reproductive age was about 40% in 2008 [17].

An important physiological indicator of the body condition is the erythrocyte sedimentation rate (ESR). All obtained ESR values fit into the physiological norm. However, these indicators were lower for young men compared to women. In terms of geographical division, for students from the northern regions, the ESR parameters were 2.7 ± 0.2 , and for representatives from the southern regions – 9.6 ± 2.4 mm/h. Among women, the differences in this parameter were also reported as significant, but less evident.

Table 5: Blood counts for students from the northern and southern regions

Indicators	Women of the northern regions (n = 100)	Men of the northern regions (n = 100)	Women of the southern regions (n = 100)	Men of the southern regions (n = 100)
The total number of leukocytes, $\times 10^9/l$	6.3 ± 0.4	6.29 ± 0.19	5.72 ± 0.1	$5.5 \pm 0.2^{\#}$
The total number of red blood cells, $\times 10^{12}/l$	4.5 ± 0.09	$4.75 \pm 0.08^*$	$4.11 \pm 0.06^{\#}$	$4.5 \pm 0.1^*$
Hemoglobin concentration, g/l	114 ± 4.5	$139.8 \pm 2.9^*$	108.2 ± 2.7	$122.9 \pm 4.8^{*\#}$
Hematocrit, %	38.6 ± 1.3	$45.3 \pm 0.9^*$	36.9 ± 0.7	$41 \pm 0.2^{*\#}$
The average volume of the red blood cell in the total sample volume, fl (MCV)	85.6 ± 2.5	$95.3 \pm 0.7^*$	90.2 ± 1.8	$90.6 \pm 1.9^{\#}$
The average volume of hemoglobin in red blood cells, pg (MCH)	28.9 ± 0.4	$30.8 \pm 0.1^*$	29.1 ± 0.2	$29.8 \pm 0.3^{\#}$
The average concentration of hemoglobin in red blood cells, gm/dl (MCHC)	273.08 ± 8.2	$222.7 \pm 10.6^*$	$322.8 \pm 21.2^{\#}$	$277.2 \pm 13.2^{\#}$
Total platelet count, $\times 10^9/l$	3.7 ± 0.3	3.6 ± 0.4	3.26 ± 0.1	3 ± 0.1
Absolute number of neutrophils, c.u	2.01 ± 0.1	1.8 ± 0.1	1.82 ± 0.07	1.9 ± 0.1
Absolute number of lymphocytes, c.u	0.49 ± 0.03	$0.6 \pm 0.04^*$	0.52 ± 0.02	$0.4 \pm 0.003^{\#}$
The absolute number of monocytes, c.u	0.11 ± 0.01	$0.14 \pm 0.01^*$	0.14 ± 0.02	0.14 ± 0.01
The absolute number of eosinophils, c.u	0.03 ± 0.005	$0.01 \pm 0.001^*$	0.02 ± 0.001	0.02 ± 0.006
The absolute number of basophils, c.u	13.3 ± 2.5	$2.7 \pm 0.2^*$	11.0 ± 1.6	$9.6 \pm 2.4^{\#}$
ESR, mm/h	2.7 ± 0.2	$2.7 \pm 0.2^*$	11.0 ± 1.6	$9.6 \pm 2.4^{\#}$

* ≤ 0.05 , significant differences between the students of different sex. $^{\#}$ Significant differences between the students from different regions. MCV: Mean corpuscular volume, MCH: Mean corpuscular hemoglobin, MCHC: Mean corpuscular hemoglobin concentration, ESR: Erythrocyte sedimentation rate.

The hematocrit parameters corresponded to standard values. In both gender groups from the northern regions, the obtained rates were higher than in students who came to study from the southern regions. Since the hematocrit reflects a fraction of the blood cell volume, it can be assumed that its increase was associated with an increase in the number of red blood cells in the circulating blood, which in this case was probably due to the stress of the body's adaptive mechanisms.

Changes in the general level of leukocyte counts are of great importance since they illustrate the body's response to the action of various exogenous and endogenous factors. A significant decrease in this parameter was revealed among the 1st-year students from the southern regions compared to those from the northern part.

As to platelet count among students of both sexes, a significant increase was observed in representatives of the southern regions, which may indicate the sensitivity of hemostasis cells to stressful effects on the body.

When studying erythrocyte indices in students, it has been established that the average erythrocyte

volume in the total sample volume among women from the southern regions was higher, while among young men from the northern regions of the Republic this parameter was significantly higher.

Changes in blood counts among students from the southern regions of the Republic compared to those from the northern regions indicated the presence of a stress reaction of the blood system when adapting to new climatic, geographical, and social conditions.

Discussion

The obtained results coincide with the data of Fedorov, who reported the lower values for the functional parameters of external respiration in the male and female population of the northern regions. The level of functioning of the respiratory system largely depends on the students' motor activity and a set of extreme factors (strong winds, low air temperature, daily changes in temperature, atmospheric pressure, etc.) [19].

Satisfactory adaptation, or the first health level, is typical for the most male and female students from the northern regions. Representatives of the southern regions have a lower adaptation potential. The group with functional tension included men and women in the southern regions. These indicators correspond to the stress state of adaptive mechanisms, in which functionality is provided by mobilizing internal reserves. The level of adaptation in males is lower compared to that in females. This can be attributed to their greater sensitivity to environmental changes, psycho-emotional condition, and gender characteristics of the cardiovascular system functioning [12], [20], [21].

The study of healthy female students at Taiba University showed that 64% of students have anemia, and 81% have a microcytic (MCV<80 fL) variety [22]. In a study of university students in the province of Hodeida, Yemen, hypochromic anemia was observed in 30.4% (of which 54% were women, 46% were men); MCV was 72.40 fL for male and 72.30 fL for female [18]. Anemia was observed in 74 students (33.4%) out of total 221 students at the University of Faisalabad. Mean \pm SD of MCV of the anemic patients were 78.36 ± 11.7 and mean \pm SD of MCV of the normal patients were 81 ± 13.10 [23]. In studies of Wu *et al.*, it was found that anemia prevails among female sex and is common for the age of 12–17 years. An increased risk of anemia was observed in underweight children from ordinary and poor rural areas. Differences in the risk of anemia depending on latitude were not observed [24]. In the research performed by female medical students in Karad, Maharashtra (India), it has been established that the prevalence of anemia was 28.6% and was reported as common in both underweight and overweight students [25].

According to Balgimbekov and Tashenova, an increase in the average erythrocyte volume is associated with a positive shift in the prooxidant-oxidative balance in cell membranes, which contributes to an increase in the functional properties of red blood cells [26].

Changes in blood counts among students from the southern regions of the Republic compared to those from the northern regions indicated the presence of a stress reaction of the blood system when adapting to new climatic, geographical, and social conditions.

Thus, a comparative assessment of the physical health of 1st-year students who came to study from different regions of the Republic showed significant differences between morphofunctional and hematological parameters. With high probability, this can be attributed to initial climatic and geographical conditions of residence before entering the university, as well as stress response and adaptation of the body to new living conditions in different regions of the Republic.

The obtained materials broaden the understanding of the influence of the complex of morphofunctional and hematological characteristics on the process of adaptation of students to education at the university. The established patterns in the adaptation of students and the data obtained are prerequisites for the creation of regional standards of physical development and functional indicators of students.

The results of the study make a certain contribution to the fundamental provisions of age physiology about the problems of human adaptation and require further research for the optimal organization of educational activities in higher educational institutions.

Conclusion

BL indicators of students from the northern regions were significantly higher than those of students from the southern regions, while higher body mass and CHC indices were recorded for students from the southern regions. The Kettle index was higher among representatives of the southern regions. By the type of physique, hypersthenics prevailed among students from the southern regions, and normostenics commonly represent the northern regions.

For the students from the southern regions, the values of the hand index, VI, HR, and BP were higher than among the northerners, which indicate good muscular development but a higher stress level of the CVS functional state in the process of adaptation.

Most young men and women of the northern regions are characterized by satisfactory adaptation. Representatives of the southern regions have though a lower adaptation potential. An integral indicator

characterizing the level of physical health showed that a larger percentage of students in the northern regions have an average health condition, while for the students in the southern regions of the Republic, it was reported to be lower.

Blood counts correspond to age and gender standards; however, for the students of the southern regions, a slight degree of anemia was noted. In students from the southern regions, an increase in the indices of platelet link and hematocrit was revealed, which is typical for the adaptation processes of the blood circulation system.

Ethical approval

The study was conducted in accordance with the ethical principles approved by the Human Experiments Ethics Committee of L.N. Gumilyov Eurasian National University (Protocol № 4 of 22.03.2019).

References

- Adilbekova AA, Smagulov NK, Sabiden GS. A comparative assessment of the features of heart rate variability in Kazakhstan and foreign medical students. *Mod Probl Sci Educ.* 2016;4:83-90.
- Mukataeva ZM, Kabieva SZ. Monitoring of physical development and health of students in Pavlodar region. *Bul NGPU.* 2014;1:51-73.
- Fedewa MV, Das BM, Evans EM, Dishman RK. Change in weight and adiposity in college students a systematic review and meta-analysis. *Am J Prev Med.* 2014;47(5):641-52. <https://doi.org/10.1016/j.amepre.2014.07.035>
PMid:25241201
- Silventomen K, Jelenkovic A, Sund R. Genetic and environmental effects on body mass index from infancy to the onset of adulthood: An individual-based pooled analysis of 45 twin cohorts participating in the Collaborative. *Am J Clin Nutr.* 2016;104:371-79. <https://doi.org/10.3410/f.726505636.793547677>
- Allison KR, Irving HM, Adlaf EM, Faulkner GE, Boak A, Manson HE, *et al.* Ten-year trends in overweight/obesity among Ontario middle and high school students and their use in establishing baseline measures for government reduction targets. *Can J Public Health.* 2015;106(8):514-19. <https://doi.org/10.17269/cjph.106.5175>
PMid:26986913
- Liu Q, Yu C, Gao W, Cao W, Lyu J, Wang S, *et al.* Genetic and environmental effects on weight, height, and BMI under 18 years in a Chinese population-based twin sample. *Twin Res Hum.* 2015;5:571-80. <https://doi.org/10.1017/thg.2015.63>
- Minister of Education and Science of the Republic of Kazakhstan. Order of the Minister of Education and Science of the Republic of Kazakhstan dated November 26, 2018 No. 646. On the Approval of the Rules for the Organization and Financing of Personnel Training with Technical and Vocational Education, Short-term Vocational Training, According to the Project "Mangilik el Zhastary-industry? ("Serpin"), Retraining of Labor Resources and Redundant Workers, Training of Qualified Personnel in Demanded Specialties in the Labor Market within the Framework of the "Zhas Maman" Project According to the "100/200" Principle, as Well as Training in the Basics of Entrepreneurship in Organizations of Technical and Professional, Post-secondary Education, Higher and (or) Postgraduate Education. Available form: https://www.online.zakon.kz/document/?doc_id=38280670
- Zueva OM, Dalenov ED, Slivkina NV. On the formation of the basics of proper nutrition and healthy lifestyles among adolescents and youth in the northern regions of Kazakhstan. *Omsk Sci Bull.* 2014;(134):115-7.
- Chandra-Mouli V, Lane C, Wong S. What does not work in adolescent sexual and reproductive health: A review of evidence on interventions commonly accepted as best practices. *Glob Health Sci Pract.* 2015;3(3):333-40. <https://doi.org/10.9745/ghsp-d-15-00126>
- Min J, Chiu DT, Wang Y. Variation in the heritability of body mass index based on diverse twin studies: A systematic review. *Obes Rev.* 2013;11:871-82. <https://doi.org/10.1111/obr.12065>
- Ndayisaba JP, Fanciulli A, Granata R, Duerr S, Hintringer F, Goebel G, *et al.* Sex and age effects on cardiovascular autonomic function in healthy adults. *Clin Auton Res.* 2016;2:169-70. <https://doi.org/10.1007/s10286-015-0335-5>
- Hulka OV. Dynamics of spectral indexes of Heart variability rate of the students with different character of the educational loading. *Fiziol Zh.* 2015;61(4):98-104. <https://doi.org/10.15407/fz61.04.098>
PMid:26552312
- Aizman RI, Lebedev AV, Aizman NI, Rubanovich VB. Methodology and practice of monitoring the health of student youth. *Health Educ.* 2017;19(5):73-8. <https://doi.org/10.26787/nydha-2226-7425-2017-19-5-73-78>
- Chernorutsky MV. The doctrine of the constitution in the clinic of internal diseases. *Proc 7th Cong Russian Ther.* 1925;304-12.
- Aizman RI, Lebedev AV, Aizman NI, Rubanovich VB. Comprehensive Assessment of the Health of Participants in the Educational Process. Moscow: Yurait; 2020.
- Baevsky RM. The problem of estimation and prognosis of functional state of the organism and its development in space medicine. *Success Physiol Sci.* 2006;37(3):42-57.
- Dangour AD, Hill HL, Ismail SJ. Original communication haemoglobin status of adult non-pregnant Kazakh women living in Kzyl-Orda region, Kazakhstan. *Eur J Clin Nutr.* 2001;5:1068-75. <https://doi.org/10.1038/sj.ejcn.1601267>
- Al-Alimi AA, Bashanfer S, Morish MA. Prevalence of iron deficiency anemia among university students in Hodeida Province, Yemen. *Anemia.* 2018;1:4157876. <https://doi.org/10.1155/2018/4157876>
- Fedorov VN. Features of Hemodynamics, the Functional State of the Myocardium and the Autonomic Regulation of Cardiac Rhythm in Young People Studying at the University, PhD Dissertation. Kazakhstan: North Kazakhstan State University; 2007.
- Schrimpf A, Kube J, Neumann J, Horstmann A, Villringer A, Gaebler M. Parasympathetic cardio-regulation during social interactions in individuals with obesity. The influence of negative body image. *Cogn Affect Behav Neurosci.* 2017;17(2):330-47. <https://doi.org/10.3758/s13415-016-0482-8>
PMid:27905081
- Tsitsiashvili KS, Kavtaradze GV, Gvaliia MV, Zhorzholiani IR, Makhadarze TG. Parameters of circadian heart rate variations, night sleep profile and structure during the Holter monitoring procedure in the groups of students with different adaptation potential. *Georgian Med News.* 2005;119:52-4.
PMid:15834182
- Al Hassan NN. The prevalence of iron deficiency anemia in

-
- a Saudi University female students. *J Microsc Ultrastruct.* 2015;3(1):25-8. <https://doi.org/10.1016/j.jmau.2014.11.003>
PMid:30023178
23. Jawed S, Tariq S, Tariq S, Kamal A. Frequency of nutritional anemia among female medical students of Faisalabad. *Pak J Med Sci.* 2017;33(2):398-403. <https://doi.org/10.12669/pjms.332.11854>
PMid:28523045
24. Wu J, Hu Y, Li M, Chen J, Mao D, Li W, *et al.* Prevalence of anemia in Chinese children and adolescents and its associated factors. *Int J Environ Res Public Health.* 2019;16(8):1416. PMid:31010238
25. Vibhute NA, Shah U, Belgaumi U, Kadashetti V, Bommanavar S, Kamate W. Prevalence and awareness of nutritional anemia among female medical students in Karad, Maharashtra, India: A cross-sectional study. *J Fam Med Prim Care.* 2019;8(7):2369-72. https://doi.org/10.4103/jfmpc.jfmpc_353_19
PMid:31463259
26. Balgimbekov SA, Tashenova GK. Features of hematological parameters in students with different levels of motor activity in modern education. *Bul Perm State Univ S Biol.* 2014;2:57-60.