



Promising Indicators in Probiotic-recommendations in COVID-19 and its Accompanying Diseases

Vardan Tsaturyan^{1,2}, Almagul Kushugulova³, Susanna Mirzabekyan⁴, Ketevan Sidamonidze⁵, David Tsereteli⁶, Tamas Torok⁷, Astghik Pepoyan^{4*}

¹Department of Military Therapy Yerevan State Medical University, Yerevan, Armenia; ²International Association for Human and Animals Health Improvement, Yerevan, Armenia; ³Centre for Life Science, National laboratory Astana, Nazarbayev University, Nur-Sultan, Kazakhstan; ⁴Department of Food Safety and Biotechnology, Armenian National Agrarian University, Yerevan, Armenia; ⁵Lugar Center for Public Health Research, Tbilisi, Georgia; ⁶Department of Communicable Diseases, National Centre for Disease Control and Public Health, Tbilisi, Georgia; ⁷Lawrence Berkeley National Laboratory, Berkeley, California, United States

Abstract

Scientific data suggests the possible beneficial role of probiotics in treatments for COVID-19, but the species/strains-specificity and disease-specificity of probiotics need high attention in choosing the appropriate probiotic in diseases, in particularly in the COVID-19. We hope this review will raise awareness of the COVID-19 probiotic recommendations, highlighting the latest scientific information about virus/hydrogen peroxide/probiotics and the importance of finding out of a specific "criterion" for the probiotics' recommendation in this disease.

Edited by: Ana Vucurevic

Citation: Tsaturyan V, Kushugulova A, Mirzabekyan S, Sidamonidze K, Tsereteli D, Torok T, Pepoyan A. Promising Indicators in Probiotic-recommendations in COVID-19 and its Accompanying Diseases. Open Access Maced J Med Sci. 2022 Jan 17; 10(B):625-631. <https://doi.org/10.3889/oamjms.2022.7989>

Keywords: COVID-19; Probiotic-specificity; Oxidative stress; Immunobiologic; Gut microbiota

***Correspondence:** Dr. Astghik Pepoyan, Department of Food Safety and Biotechnology, Armenian National Agrarian University, Yerevan, Armenia. E-mail: apepoyan@gmail.com

Received: 21-Nov-2021

Revised: 30-Dec-2021

Accepted: 07-Jan-2022

Copyright: © 2022 Vardan Tsaturyan, Almagul Kushugulova, Susanna Mirzabekyan, Ketevan Sidamonidze, David Tsereteli, Tamas Torok, Astghik Pepoyan

Funding: This research was funded by State Committee of Science of the Republic of Armenia (grant numbers: 10-15/J-5 and 21AG-4D065) and Science Committee of the Ministry of Education and Science of the Republic of Kazakhstan (grant number AP09563198).

Competing Interests: The authors have declared that no competing interests exist

Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Introduction

Literature data indicate an association between COVID-19 severity and diabetes [1], [2]. Most COVID-19 patients are prone to impaired glucose metabolism; glycemic testing and control are important even if the patients have no pre-existing diabetes [3]. Furthermore, hypertension [1], [4], acute coronary syndrome [5], rheumatic [6], gastrointestinal [7], and neurologic features [8], [9] in SARS-CoV-2 infection have been reported. Potential associations between host blood characteristics and gut bacteria [10], [11], [12], as well as between gut microbiota and COVID-19 – accompanying diseases have been actively discussed [13]. Sever childhood respiratory illness in association with vitamin D deficiency has also been shown [14]. The clinical trials and experimental studies on COVID-19 treatments are

ongoing worldwide, increasing the obtained information on infected people, blood and organ system, genomics, and metabolomics. Despite of a safe and efficacious vaccines, respiratory tract infections will remain of concern for high morbidity and mortality rates among the elderly due to low level vaccine-induced immune response [15], [16]. Recently collected data appear to confirm the possible beneficial role of probiotics in treating COVID-19 patients [17], [18]; however, there is insufficient scientific evidence specific to COVID-19. Therefore, species/strain- and disease-specificity of probiotics need more attention [19]. Probiotics may have beneficial, harmful, or neutral impact on the host. For example, in an *in vivo* study of their radio-preventive/protective characteristics, 17 putative probiotic lactobacilli, including the strain *Lactobacillus acidophilus* DDS[®]-1 (from Lacto-G, a marketed symbiotic formulation), the commercial probiotic product Narine (*L. acidophilus* INMIA 9602 Er-2

strain 317/402), and several strains of *Lactobacillus rhamnosus*, *Lactobacillus plantarum*, *Lactobacillus casei*, *Lactobacillus fermentum*, *Lactobacillus delbrueckii*, and *Lactobacillus helveticus*, have shown varying impact on 4.5 Gy, whole-body X-ray irradiated rats before and after the irradiation [20]. In addition, since COVID-19 patients may rapidly “transform” their physiological state following an infection by the virus, potential probiotic-effects and the individual dietary, nutritional, medical, lifestyle, and environmental risks should be carefully investigated before any recommendation. Since it was recognized that the SARS-CoV-2 pandemic has been particularly deadly in older adults [21], [22], we hope to raise awareness about probiotics for COVID-19 patients and highlight the possible importance of hydrogen peroxide production for any probiotics recommendation in this disease.

Probiotics/Immunobiotics

Nowadays, the interaction of probiotics with the SARS-CoV-2 spike (S) proteins [22] and with the gut microbiome [23], [24], [25] is at the attention of the researchers. In general, the benefit of probiotics is determined by the complex interactions between probiotic bacteria, the host intestinal microbiota, and the gut epithelium [23], [24], [25]. Frequently, probiotic lactobacilli are able to (i) promote the expression and regulation of tight junctions and adherence junctions, resulting in the restoration of a defective epithelial barrier and (ii) interact with immune cells through pattern recognition receptors, such as Toll-like receptors, which on activation stimulate or suppress various immune responses [26], emphasizing host gut-blood [27], and gut-brain [28] linkages. Probiotic immunobiotics beneficially regulate the mucosal immune system [29] and have valuable antagonistic potential against nosocomial pathogens. Among the many proposed mechanisms by which immunobiotics mediate their effects is the modulation of the innate immune response both by anti-inflammatory [30], [31] and pro-inflammatory effects [32]. In addition, immunobiotics have been shown to enhance the adaptive immune response, for example, antibody formation [33]. Inhibition of adherence, attaching, and effacing microorganisms [34], modulation of mucosal barrier function [35], or inhibition of trophic migration [36] may also be important mechanisms, whereby immunobiotics may influence intestinal diseases [37]. There is also strong evidence that signaling molecules/determinants are preserved in immunobiotic strains [38], and certain immunogenic strains enhance immune function, especially in subjects with less than adequate immune response [39].

Human Viral Diseases, Oxidative Stress, Hydrogen Peroxide, and Probiotics

The prevalence of chronic diseases increases with age. Increased production of reactive oxygen species is involved in the pathogenesis of cardiovascular diseases, such as coronary atherosclerosis, hypertension, diabetic vascular complications, and heart failure [40]. These are also known risks for potentially worse COVID-19 outcome. The patients with severe COVID-19 history have also often been found to have elevated D-dimer, troponin, ferritin, C-reactive protein, and alanine aminotransferase [41], and circulating endothelial cell levels [42], as well as vitamin D deficiency [14]. It seems that there is a link between oxidative stress and a variety of pathological conditions, including COVID-19 [43] and its accompanying diseases, such as diabetes [44], hypertension [45], acute coronary syndrome [46], rheumatic [47], gastrointestinal [48], or neurologic features [49].

Both enzymatic and non-enzymatic pathways are involved in endogenous antioxidant defense mechanisms. The enzymes superoxide dismutase, catalase, the prosthetic group Se-containing glutathione peroxidase, and glutathione reductase are common antioxidants; important in endogenous antioxidant defense systems [44]. Iron is an essential element for virtually all cell types due to its role in energy metabolism, but free iron may induce cellular and organ damage through the free radicals [50]. Peroxidase uses various organic compounds: Polyphenols, aromatic amines, ascorbic acid, etc., to destroy hydrogen peroxide and organic peroxides (oxygen donors), and to form a high valent iron intermediate named Compound I. In erythrocytes and some other tissues, glutathione peroxidase protects membranes and hemoglobin from oxidation by peroxides. Peroxisomes, the cell organelles known as important centers in innate immune-, lipid-, inflammatory-, and redox-signaling networks [51], by regulating their number, shape, and protein content in response to changing environmental conditions [52], have the intrinsic ability to mediate and modulate H₂O₂-driven biological processes [53].

Free radicals/other reactive oxygen and nitrogen species show a double role, causing oxidative damage/tissue dysfunction and serving as molecular signals activating beneficial stress responses [54], [55]. H₂O₂ emerged as a major redox metabolite operative in redox sensing, signaling, and regulation [56]. Its action mainly depends on the cellular context, its local concentration, and the kinetics of its production and elimination [57]. Interestingly, long-lasting blood pressure lowering effects of nitrite are NO-independent and are mediated by hydrogen peroxide, persulfides, and oxidation of protein kinase G1 α redox signaling [58]. Although H₂O₂ is a strong oxidizing agent, it can accumulate in cells and tissues to relatively high concentrations due to it

slow reaction kinetics with most biomolecules. The removal of excess H_2O_2 by antioxidant enzymes is therefore central in minimizing cellular damage [59].

Beside the endogenous H_2O_2 produced by immune cells to kill pathogenic microbes, to inhibit other competing bacteria, lactic acid bacteria also produce H_2O_2 , using different enzymes that include pyruvate oxidase, lactate oxidase, NADH oxidase, and NADH flavin-dependent reductases [60]. There are indications that H_2O_2 -producing lactobacilli in the intestine play an important role in the repair of intestinal damage [61], whereas H_2O_2 -producing lactobacilli in the vagina control the growth of pathogenic bacteria and prevent tumorigenesis [62]. Unfortunately, pathogenic *Streptococcus pneumoniae* (through SpxB) and *Streptococcus pyogenes* (through LacD) are also known to produce H_2O_2 , possibly through the lactate oxidation pathway [63]. While the host interactions in general are unclear, it was shown that H_2O_2 released by *Streptococcus pneumoniae* inhibits host inflammasomes and is responsible for pathogen colonization [64]. Blockage of inflammasome activation by the oral commensal H_2O_2 -producing bacterium *Streptococcus oralis* was also reported [64].

There are numerous reports indicating the role of hydrogen peroxide in viral inactivation and in virus-host interaction. For example,

- Herpes simplex virus 1-associated catalase may protect the virus from inactivation in an oxidizing environment outside a host cell [65];
- Uncontrolled concentrations of H_2O_2 promote translation by the internal ribosome entry site element of hepatitis C virus in tissue-cultured cells through adaptation of oxidative stress in the host cell by mediating La cytoplasmic shuttling [66];

The addition of polyethylene glycol-conjugated catalase increased the specific enzyme activity, which, in turn, along with a respiratory syncytial virus (RSV) infection decreased H_2O_2 in the airways, and had an important defensive impact on RSV-induced disease/pathology. Therefore, it was concluded that the addition of catalase might represent a new pharmacological approach that should be studied in humans for the prevention and treatment of respiratory infections caused by RSV [67];

- H_2O_2 -producing lactobacilli in the vagina may control genital HIV-1 shedding [68], the growth of several pathogenic bacteria, and may prevent tumorigenesis [62]. Along these ideas, Krüger and Bauer showed that the lactobacilli-origin H_2O_2 intrinsically is not likely to assist the vaginal epithelium, as it origins apoptosis both in non-transformed and in transformed cells [69]. The authors suggested that a combination of lactobacilli and peroxidase, that is, the situation actually found for tumor

tissue *in vivo*, leads to the conversion of H_2O_2 to HOCl which does not affecting on non-malignant cells. When malignant cells, due to the abundance of extracellular peroxide anions, allow the formation of apoptosis. Subsequently, the combination of peroxide producing lactobacilli and peroxidase causes the selective elimination of malignant cells [69].

Discussion

Clinical trials and experimental studies have shown that probiotics, their components, or their sterilized variants (paraprobiotics) may be successfully used as biotherapeutic agents for the prevention and treatment of gastrointestinal diseases [70], [71], [72], [73], [74], and for resistance enhancement in case of intestinal viral infections [75], [76]. The possibility of mitigating antimicrobial resistance, which might be a result of a nosocomial, COVID-19-related infection [77], through probiotics has also been discussed in publication [78]. A study on the impact of the probiotic “Narine” (Vitamax-E, Armenia) on the number of antibiotic resistant gut commensal *Escherichia coli* in familial Mediterranean fever patients, an autosomal recessive inflammatory disease [79], showed that the probiotic therapy resulted in a reduction of the relative abundance of operational taxonomic units in the genus *Escherichia* and the number of multi-resistant *E. coli* isolates [78].

Appropriate inflammasome activation and appropriate concentration of hydrogen peroxide realized by bacteria [64] are vital for the host to handle foreign pathogens and tissue damage, while aberrant inflammasome activation can cause uncontrolled tissue responses, leading to various diseases, including autoinflammatory disorders, cardiometabolic diseases, cancer, and neurodegenerative diseases [80]. In short, research data indicate the important role of hydrogen peroxide in host-microbe interaction. Therefore, the assessment of blood H_2O_2 , blood catalase activities, and detection of inflammasome activity present promising indicators in recommending probiotics in COVID-19 and accompanying diseases.

Conclusions

Thus, socioeconomic and biological strategies are needed to combat COVID-19 [81], [82], [83]. The production of hydrogen peroxide might be considered as one of promising indicators in probiotic recommendation in COVID-19 and its accompanying

diseases. We hope that continuing investigations will raise awareness of potential COVID-19 probiotic recommendations, highlighting the latest scientific information about COVID-19/probiotics, and the importance of determining the specific “criteria” for the recommendation of probiotics’ use in this disease.

References

- Hussain A, Bhowmik B, do Vale Moreira NC. COVID-19 and diabetes: Knowledge in progress. *Diabetes Res Clin Pract.* 2020;162:108142. <https://doi.org/10.1016/j.diabres.2020.108142>
PMid:32278764
- Guan WJ, Liang WH, Zhao Y, Liang HR, Chen ZS, Li YM, et al. Comorbidity and its impact on 1590 patients with COVID-19 in China: A nationwide analysis. *Eur Respir J.* 2020;55(5):2000547. <https://doi.org/10.1183/13993003.00547-2020>
PMid:32217650
- Wang S, Ma P, Zhang S, Song S, Wang Z, Ma Y, et al. Fasting blood glucose at admission is an independent predictor for 28-day mortality in patients with COVID-19 without previous diagnosis of diabetes: A multi-centre retrospective study. *Diabetologia.* 2020;63(10):2102-11. <https://doi.org/10.1007/s00125-020-05209-1>
PMid:32647915
- Schiffri EL, Flack JM, Ito S, Muntner P, Webb RC. Hypertension and COVID-19. *Am J Hypertens.* 2020;33(5):373-74. <https://doi.org/10.1093/ajh/hpaa057>
PMid:32251498
- Metzler B, Siostrzonek P, Binder RK, Bauer A, Johannes Reinstadler S. Decline of acute coronary syndrome admissions in Austria since the outbreak of COVID-19: The pandemic response causes cardiac collateral damage. *Eur Heart J.* 2020;41(19):1852-53. <https://doi.org/10.1093/eurheartj/ehaa314>
PMid:32297932
- Misra DP, Agarwal V, Gasparyan AY, Zimba O. Rheumatologists’ perspective on coronavirus disease 19 (COVID-19) and potential therapeutic targets. *Clin Rheumatol.* 2020;39(7):2055-62. <https://doi.org/10.1007/s10067-020-05073-9>
PMid:32277367
- Tsaturyan VV, Kushugulova A, Sidamonidze K, Tsereteli D, Pepoyan AZ. Gut Microbiota Composition and Disease Severity in Patients with COVID-19: Armenian Population. *Topical Issues of Biological Safety in Modern Conditions. Online Meeting, Almaty 22-23 September; 2021.* p. 6-7. <https://doi.org/10.13140/RG.2.2.28336.48642>
- Mao L, Jin H, Wang M. Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in Wuhan, China. *JAMA Neurol.* 2020;77(6):683-90. <https://doi.org/10.1001/jamaneurol.2020.1127>
PMid:32275288
- Manji H, Carr AS, Brownlee WJ, Lunn MP. Neurology in the time of Covid-19. *J Neurol Neurosurg Psychiatry.* 2020;91(6):568-70. <https://doi.org/10.1136/jnnp-2020-323414>
PMid:32312872
- Pluznick J. Gut microbes and host physiology: What happens when you host billions of guests? *Front Endocrinol (Lausanne).* 2014;5:91. <https://doi.org/10.3389/fendo.2014.00091>
PMid:24982653
- Pepoyan AZ, Tsaturyan VV, Badalyan M, Weeks R, Kamiya S, Chikindas ML. Blood protein polymorphisms and the gut bacteria: Impact of probiotic *Lactobacillus acidophilus* Narine on *Salmonella* carriage in Sheep. *Benef Microbes.* 2020;11:183-9. <https://doi.org/10.3920/BM2019.0138>
PMid:32028777
- Pepoyan A, Harutyunyan N, Grigoryan A, Balayan M, Tsaturyan V, Manvelyan A, et al. Some clinical blood characteristics of patients with familial Mediterranean fever disease from an Armenian cohort. *Klin Lab Diagn.* 2015;60(6):46-8.
- Kazemian N, Mahmoudi M, Halperin F, Wu JC, Pakpour S. Gut microbiota and cardiovascular disease: Opportunities and challenges. *Microbiome.* 2020;8(1):36. <https://doi.org/10.1186/s40168-020-00821>
PMid:32169105
- Rhodes JM, Subramanian S, Laird E, Griffin G, Kenny RA. Perspective: Vitamin D deficiency and COVID-19 severity plausibly linked by latitude, ethnicity, impacts on cytokines, ACE2 and thrombosis. *J Intern Med.* 2021 289(1):97-115. <https://doi.org/10.1111/joim.13149>
PMid:32613681
- Oh SJ, Lee JK, Shin OS. Aging and the immune system: The impact of immunosenescence on viral infection, immunity and vaccine immunogenicity. *Immune Netw.* 2019;19(6):e37. <https://doi.org/10.4110/in.2019.19.e37>
PMid:31921467
- Knoll MD, Wonodi C. Oxford-AstraZeneca COVID-19 vaccine efficacy. *Lancet.* 2020;397(10269):72-74. [https://doi.org/10.1016/S0140-6736\(20\)32623-4](https://doi.org/10.1016/S0140-6736(20)32623-4)
PMid:33306990
- Baud D, Agri VD, Gibson GR, Reid G, Giannoni E. Using probiotics to flatten the Curve of coronavirus disease COVID-2019 pandemic. *Front Public Health.* 2020;8:186. <https://doi.org/10.3389/fpubh.2020.00186>
PMid:32574290
- Villena J, Kitazawa H. The modulation of mucosal antiviral immunity by immunobiotics: Could they offer any benefit in the SARS-CoV-2 pandemic? *Front Physiol.* 2020;11:699. <https://doi.org/10.3389/fphys.2020.00699>
PMid:32670091
- McFarland LV, Evans CT, Goldstein E. Strain-specificity and disease-specificity of probiotic efficacy: A systematic review and meta-analysis. *Front Med.* 2018;5:124. <https://doi.org/10.3389/fmed.2018.00124>
PMid:29868585
- Pepoyan AZ, Manvelyan AM, Balayan MH, McCabe G, Tsaturyan VV, Melnikov VG, et al. The effectiveness of potential probiotics *Lactobacillus rhamnosus* Vahe and *Lactobacillus delbrueckii* IAHAI in irradiated rats depends on the nutritional stage of the host. *Probiotics Antimicrob Proteins.* 2020;12(4):1439-50. <https://doi.org/10.1007/s12602-020-09662-7>
PMid:32462507
- Armitage R, Nellums LB. COVID-19 and the consequences of isolating the elderly. *Lancet.* 2020;5(5):e256. [https://doi.org/10.1016/s2468-2667\(20\)30061-x](https://doi.org/10.1016/s2468-2667(20)30061-x)
PMid:32199471
- Tsaturyan V, Pepoyan A. Probiotics for COVID-19: Bacterocin, Host Microbe Interaction and Molecular Docking. *International BioThreat Reduction Symposium (IBTRS) 28 June-2 July 2021. Kyiv, Ukraine; 2021.* p. 61. <https://doi.org/10.13140/RG.2.2.19134.82242>
- Pepoyan A, Balayan M, Malkasyan L, Manvelyan A, Bezhanyan T, Paronikyan R, et al. Effects of probiotic *Lactobacillus acidophilus* strain INMIA 9602 Er 317/402 and putative probiotic lactobacilli on DNA damages in small

- intestine of Wistar rats *in vivo*. Probiotics Antimicrob Proteins. 2020;11:905-9. <https://doi.org/10.1007/s12602-018-9491-y>
24. Kossumov A, Mussabay K, Pepoyan A, Tsaturyan V, Sidamonidze K, Tsereteli D, et al. Digestive system and SARS-COV-2: New era of microbiome study and gastrointestinal tract manifestations during the Covid-19 pandemic. Open Access Maced J Med Sci. 2021;9(F):676-82. <https://doi.org/10.3889/oamjms.2021.7470>.
 25. Lewis Z, Sidamonidze K, Tsaturyan V, Tsereteli D, Khachidze N, Pepoyan A, et al. The fecal microbial community of breast-fed infants from Armenia and Georgia. Sci Rep. 2017;7:40932. <https://doi.org/10.1038/srep40932>
PMid:28150690
 26. Ussar S, Fujisaka S, Kahn C. Interactions between host genetics and gut microbiome in diabetes and metabolic syndrome. Molecul Metab 2016;5:795-803. <https://doi.org/10.1016/j.molmet.2016.07.004>
PMid:27617202
 27. Pepoyan A, Balayan M, Manvelyan A, Galstyan L, Pepoyan S, Petrosyan S, et al. Probiotic *Lactobacillus acidophilus* strain INMIA 9602 Er 317/402 administration reduces the numbers of *Candida albicans* and abundance of *Enterobacteria* in the gut microbiota of familial mediterranean fever patients. Front Immunol. 2018;9:1426. <https://doi.org/10.3389/fimmu.2018.01426>
PMid:29997616
 28. Clemmensen C, Müller TD, Woods SC, Berthoud HR, Seeley RJ, Tschöp MH. Gut-brain cross-talk in metabolic control. Cell. 2017;168:758-74. <https://doi.org/10.1016/j.cell.2017.01.025>
PMid:28235194
 29. Villena J, Kitazawa H. Editorial: Immunobiotics interactions of beneficial microbes with the immune system. Front Immunol. 2017;8:1580. <https://doi.org/10.3389/fimmu.2017.01580>
PMid:29250061
 30. Imaoka A, Shima T, Kato K, Mizuno S, Uehara T, Matsumoto S, et al. Anti-inflammatory activity of probiotic *Bifidobacterium*: Enhancement of IL-10 production in peripheral blood mononuclear cells from ulcerative colitis patients and inhibition of IL-8 secretion in HT-29 cells. World J Gastroenterol 2008;14(16):2511-6. <https://doi.org/10.3748/wjg.14.2511>
PMid:18442197
 31. Cosseu C, Devine DA, Dullaghan E, Gardy JL, Chikatamarla A, Gellatly S, et al. The commensal *Streptococcus salivarius* K12 downregulates the innate immune responses of human epithelial cells and promotes host-microbe homeostasis. Infect Immun. 2008;76(9):4163-75. <https://doi.org/10.1128/IAI.00188-08>
PMid:18625732
 32. Galstyan L, Tsaturyan V, Pepoyan A. Efficiency of pre and probiotic therapy for the management of periodic disease and hypoxic ischemic encephalopathy of newborns: NLRP3 inflammasome. PARMA (in Armenian) 2008;57:49-51.
 33. Vaarala O. Immunological effects of probiotics with special reference to lactobacilli. Clin Exp Allergy. 2003;33:1634-40. <https://doi.org/10.1111/j.1365-2222.2003.01835.x>
PMid:14656348
 34. Mack D, Michail S, Wei S, McDougall L, Hollingsworth M. Probiotics inhibit enteropathogenic *E. coli* adherence *in vitro* by inducing intestinal Mucin gene expression. Am J Physiol. 1999;276:941-50. <https://doi.org/10.1152/ajpgi.1999.276.4.G941>
PMid:10198338
 35. Mack D, Ahne S, Hyde L, Wei S, Hollingsworth M. Extracellular MUC3 mucin secretion follows adherence of *Lactobacillus* strains to intestinal epithelial cells *in vitro*. Gut. 2003;52(6):827-33. <https://doi.org/10.1136/gut.52.6.827>
PMid:12740338
 36. Michail S, Abernathy F. *Lactobacillus plantarum* inhibits the intestinal epithelial migration of neutrophils induced by enteropathogenic *Escherichia coli*. J Pediatr Gastroenterol Nutr. 2003;36():385-91. <https://doi.org/10.1097/00005176-200303000-00017>
PMid:12604980
 37. Pepoyan A, Trchounian A. Biophysics, molecular and cellular biology of probiotic activity by bacteria. In: Trchounian A, editors. Bacterial Membranes. Ultrastructure, Bioelectrochemistry, Bioenergetics and Biophysics. Kerala, India: Trivandrum, Signpost; 2020. p. 275-87.
 38. Bu H, Wang X, Zhu Y, Williams R, Hsueh W, Zheng X, et al. Lysozyme-modified probiotic components protect rats against polymicrobial sepsis: Role of macrophages and cathelicidin-related innate immunity. J. Immunol. 2006;177:8767-76. <https://doi.org/10.4049/jimmunol.177.12.8767>
 39. Gill H, Rutherford K, Cross M, Gopal P. Enhancement of immunity in the elderly by dietary supplementation with the probiotic *Bifidobacterium lactis* HN019. Am J Clin Nutr. 2001;74:833-9. <https://doi.org/10.1093/ajcn/74.6.833>
PMid:11722966
 40. Petrie JR, Guzik TJ, Touyz RM. Diabetes, hypertension, and cardiovascular disease: Clinical insights and vascular mechanisms. Can J Cardiol. 2018;34(5):575-84. <https://doi.org/10.1016/j.cjca.2017.12.005>
PMid:29459239
 41. Sultan S, Sultan M. COVID-19 cytokine storm and novel. Med Hypotheses. 2020;144:109875. <https://doi.org/10.1016/j.mehy.2020.109875>
PMid:18442197
 42. Guervilly C, Burtsey S, Sabatier F, Cauchois RL, Lano G, Abdili E, et al. Circulating endothelial cells as a marker of endothelial injury in severe COVID-19. J Infect Dis. 2020;222(11):1789-93. <https://doi.org/10.1093/infdis/jiaa528>
PMid:32812049
 43. Liguori I, Russo G, Curcio F, Bulli G, Aran L, Della-Morte D, et al. Oxidative stress, aging, and diseases. Clin Interv Aging. 2018;13:757-72. <https://doi.org/10.2147/CIA.S158513>
PMid:29731617
 44. Matough FA, Budin SB, Hamid ZA, Alwahaibi N, Mohamed J. The role of oxidative stress and antioxidants in diabetic complications. Sultan Qaboos Univ Med J. 2012;12(1):5-18. <https://doi.org/10.12816/0003082>
PMid:22375253
 45. Rodrigo R, González J, Paoletto F. The role of oxidative stress in the pathophysiology of hypertension. Hypertens Res. 2011;34(4):431-40. <https://doi.org/10.1038/hr.2010.264>
PMid:21228777
 46. Vichova T, Motovska Z. Oxidative stress: Predictive marker for coronary artery disease. Exp Clin Cardiol. 2013;18(2):e88-91. PMid:23940453
 47. Quiñonez-Flores CM, González-Chávez SA, Del Río Nájera D, Pacheco-Tena C. Oxidative stress: Predictive marker for coronary artery disease. Exp Clin Cardiol. Biomed Res Int. 2016;2016:6097417. <https://doi.org/10.1155/2016/6097417>
 48. Saha SK, Lee SB, Won J, Choi HY, Kim K, Yang GM, et al. Correlation between oxidative stress, nutrition, and cancer initiation. Int J Mol Sci. 2018;18(7):1544. <https://doi.org/10.3390/ijms18071544>
 49. Patel M. Targeting oxidative stress in central nervous system disorders. Trends Pharmacol Sci. 2016;37(9):768-78. <https://doi.org/10.1016/j.tips.2016.06.007>
PMid:27491897

50. Abbaspour N, Hurrell R, Kelishadi R. Review on iron and its importance for human health. *J Res Med Sci.* 2014;19(2):164-74. PMID:24778671
51. Wang B, Apanasets O, Nordgren M, Fransen M. Dissecting peroxisome-mediated signaling pathways: A new and exciting research field. In: Brocard C, Hartig A, editors. *Molecular Machines Involved in Peroxisome Biogenesis and Maintenance*. 1st ed. Wien, Austria: Springer; 2014. p. 255-73.
52. Fransen M. Peroxisome dynamics: Molecular players, mechanisms, and (Dys) functions. *Int Sch Res Notices.* 2012;2012:714192. <https://doi.org/10.5402/2012/714192>
53. Lismont C, Revenco I, Fransen M. Peroxisomal hydrogen peroxide metabolism and signaling in health and disease. *Int J Mol Sci.* 2015;20(15):3673. <https://doi.org/10.3390/ijms20153673> PMID:31357514
54. Di Meo S, Reed TT, Venditti P, Victor VM. Role of ROS and RNS Sources in physiological and pathological conditions. *Oxid Med Cell Longev.* 2016;2016:1245049. <https://doi.org/10.1155/2016/1245049> PMID:27478531
55. Di Meo S, Venditti P. Evolution of the knowledge of free radicals and other oxidants. *Oxid Med Cell Longev.* 2020;2020:9829176. <https://doi.org/10.1155/2020/9829176> PMID:32411336
56. Sies H. Hydrogen peroxide as a central redox signaling molecule in physiological oxidative stress: Oxidative eustress. *Redox Biol.* 2017;11:613-9. <https://doi.org/10.1016/j.redox.2016.12.035> PMID:28110218
57. Auten RL, Davis JM. Oxygen toxicity and reactive oxygen species: The devil is in the details. *Pediatr Res.* 2009;66(2):121-7. <https://doi.org/10.1203/PDR.0b013e3181a9eafb> PMID:19390491
58. Feelisch M, Akaike T, Griffiths K, Ida T, Pryszyzna O, Goodwin JJ, et al. Long-lasting blood pressure lowering effects of nitrite are NO-independent and mediated by hydrogen peroxide, persulfides, and oxidation of protein kinase G1 α redox signalling. *Cardiovasc Res.* 2020;116(1):51-62. <https://doi.org/10.1093/cvr/cvz202> PMID:31372656
59. Doskey CM, Buranasudja V, Wagner BA, Wilkes JG, Du J, Cullen JJ, et al. Tumor cells have decreased ability to metabolize H₂O₂: Implications for pharmacological ascorbate in cancer therapy. *Redox Boil.* 2016;10:274-84. <https://doi.org/10.1016/j.redox.2016.10.010> PMID:27833040
60. Hertzberger R, Arents J, Dekker HL, Pridmore RD, Gysler C, Kleerebezem M, et al. H₂O₂ production in species of the *Lactobacillus acidophilus* group: A central role for a novel NADH-dependent flavin reductase. *Appl Environ Microbiol.* 2014;80:2229-39. <https://doi.org/10.1128/AEM.04272-13>
61. Singh AK, Hertzberger RY, Knaus UG. Hydrogen peroxide production by lactobacilli promotes epithelial restitution during colitis. *Redox Boil.* 2018;16:11-20. <https://doi.org/10.1016/j.redox.2018.02.003> PMID:29471162
62. Klebanoff SJ, Hillier SL, Eschenbach DA, Walterdorff AM. Control of the microbial flora of the vagina by hydrogen peroxide-generating lactobacilli. *J Infect Dis.* 1991;164:94-100. <https://doi.org/10.1093/infdis/164.1.94> PMID:1647428
63. Seki M, Iida K, Saito M, Nakayama H, Yoshida S. Hydrogen peroxide production in *Streptococcus pyogenes*: Involvement of lactate oxidase and coupling with aerobic utilization of lactate. *J Bacteriol.* 2018;186(7):2046-51. <https://doi.org/10.1128/JB.186.7.2046-2051> PMID:15028688
64. Erttman S, Gekara N. Hydrogen peroxide release by bacteria suppresses inflammasome-dependent innate immunity. *Nat Commun.* 2019;10(1):3493. <https://doi.org/10.1038/s41467-019-11169-x> PMID:31375698
65. Newcomb WW, Brown JC. Internal catalase protects herpes simplex virus from inactivation by hydrogen peroxide. *J Virol.* 2012;86:11931-4. <https://doi.org/10.1128/JVI.01349-12> PMID:22915822
66. Chan SW. Hydrogen peroxide induces la cytoplasmic shuttling and increases hepatitis C virus internal ribosome entry site-dependent translation. *J Gen Virol.* 2016;97:2301-15. <https://doi.org/10.1099/jgv.0.000556> PMID:27436793
67. Ansar M, Ivanciuc T, Garofalo RP, Casola A. Increased lung catalase activity confers protection against experimental RSV infection. *Sci Rep.* 2020;10:3653. <https://doi.org/10.1038/s41598-020-60443-2>
68. Balkus JE, Mitchell C, Agnew K, Liu C, Fiedler T, Cohn SE, et al. Detection of hydrogen peroxide-producing *Lactobacillus* species in the vagina: A comparison of culture and quantitative PCR among HIV-1 seropositive women. *BMC Infect Dis.* 2012;12:188. <https://doi.org/10.1186/1471-2334-12-188> PMID:22889380
69. Krüger H, Bauer G. Lactobacilli enhance reactive oxygen species-dependent apoptosis-inducing signaling. *Redox Biol.* 2017;11:715-24. <https://doi.org/10.1016/j.redox.2017.01.015> PMID:28193594
70. Balayan M, Manvelyan A, Marutyan S, Isajanyan M, Tsaturyan V, Pepoyan A, et al. Impact of *Lactobacillus acidophilus* INMIA 9602 Er-2 and *Escherichia coli* M-17 on some clinical blood characteristics of familial Mediterranean fever disease patients from the Armenian cohort. *Int J Probiotics Prebiotics.* 2015;10:91-5.
71. Bagherpour G, Ghasemi H, Zand B, Zarei N, Roohvand F, Ardakani EM, et al. Oral administration of recombinant *Saccharomyces boulardii* expressing ovalbumin-CPE fusion protein induces antibody response in mice. *Front Microbiol.* 2018;9:723. <https://doi.org/10.3389/fmicb.2018.00723> PMID:29706942
72. Pepoyan AZ, Balayan MH, Manvelyan AM, Mamikonyan V, Isajanyan M, Tsaturyan VV, et al. *Lactobacillus acidophilus* INMIA 9602 Er-2 strain 317/402 probiotic regulates growth of commensal *Escherichia coli* in gut microbiota of familial Mediterranean fever disease subjects. *Lett Appl Microbiol.* 2017;64:254-60. <https://doi.org/10.1111/lam.12722> PMID:28140472
73. Pepoyan A, Balayan M, Manvelyan A, Pepoyan S, Malkhasyan L, Bezhanian T, et al. Radioprotective effects of lactobacilli with antagonistic activities against human pathogens. *Biophys J.* 2018;114:665a. <https://doi.org/10.1016/j.bpj.2017.11.3586>
74. Pepoyan AZ, Pepoyan ES, Galstyan L, Harutyunyan NA, Tsaturyan VV, Torok T, et al. The effect of immunobiotic/psychobiotic *Lactobacillus acidophilus* strain INMIA 9602 Er 317/402 narine on gut prevotella in familial Mediterranean fever: Gender-associated effects. *Probiotics Antimicrob Prot* 2021;13:1306-15. <https://doi.org/10.1007/s12602-021-09779-3> PMID:34132998
75. Kanauchi O, Andoh A, AbuBakar S, Yamamoto N. Probiotics and paraprobiotics in viral infection: Clinical application and effects on the innate and acquired immune systems. *Curr Pharm Des.* 2018;24:710-7. <https://doi.org/10.2174/1381612824666180116163411>

- PMid:29345577
76. Eguchi K, Fujitani N, Nakagawa H, Miyazaki T. Prevention of respiratory syncytial virus infection with probiotic lactic acid bacterium *Lactobacillus gasser* SBT2055. *Sci Rep*. 2019;18:4812. <https://doi.org/10.1038/s41598-019-39602-7>
77. Carter B, Collins JT, Barlow-Pay F, Rickard F, Bruce E, Verduri A, et al. Nosocomial COVID-19 infection: Examining the risk of mortality. The COPE-nosocomial study (COVID in older people). *J Hosp Infect*. 2020;106(2):376-84. <https://doi.org/10.1016/j.jhin.2020.07.013>
PMid:32702463
78. Pepoyan AZ, Balayan MA, Arutyunyan NA, Grigoryan AG, Tsaturyan VV, Manvelyan AM, et al. Antibiotic resistance of *Escherichia coli* of the intestinal microbiota in patients with familial Mediterranean Fever *Klin Med (Mosk)*. 2015;93:37-9.
PMid:26596057
79. Touitou I, Pepoyan A. Concurrence of Crohn's and familial Mediterranean fever diseases for Armenian cohort. *Inflamm Bowel Dis*. 2008;14:S39.
<https://doi.org/10.1097/00054725-200812003-00128>
80. Zheng D, Liwinski T, Elinav E. Inflammasome activation and regulation: Toward a better understanding of complex mechanisms. *Cell Discov*. 2020;6:36. <https://doi.org/10.1038/s41421-020-0167-x>
PMid:32550001
81. Telles CR, Roy A, Ajmal MR, Mustafa SK, Ahmad MA, de la Serna JM, et al. The impact of COVID-19 management policies tailored to airborne SARS-CoV-2 transmission: Policy analysis. *JMIR Public Health Surveill*. 2021;7(4):e20699. <https://doi.org/10.2196/20699>
PMid:33729168
82. Khalid MS, Aljohani MM, Alomrani NA, Oyouni AA, Alzahrani O, Ahmad OA, et al. COVID-19 and immune function "a significant" zinc. *Orient J Chem* 2020;36:1026-36.
83. Khalid MS, Ahmad MA, Baranova V, Deineko Z, Lyashenko V, Oyouni AA. Using wavelet analysis to assess the impact of COVID-19 on changes in the price of basic energy resources. *Int J Emerg Trends Eng Res*. 2020;8(7):2907-12.