



Hormonal Perturbations in Occupationally Exposed Nickel Workers

Safia Beshir*, Khadiga Salah Ibrahim, Weam Shaheen, Eman M. Shahy

Environmental and Occupational Medicine Department, National Research Centre, Dokki, Giza, Egypt

Abstract

Citation: Beshir S, Ibrahim KS, Shaheen W, Shahy EM. Hormonal Perturbations in Occupationally Exposed Nickel Workers. OA Maced J Med Sci. <http://dx.doi.org/10.3889/oamjms.2016.046>

Key words: Nickel; FSH; LH; Testosterone; sexual problems.

***Correspondence:** Safia Beshir Ahmed. Professor of Environmental Health, Environmental and Occupational Medicine Department, National Research Centre, 33 EL Bohouth st. (former EL Tahrir st.) - Dokki - Giza - Egypt - P.O. 12622. Tel.: 002 01006285633. E-mail: safiabeshir123@yahoo.com

Received: 07-Feb-2016; **Revised:** 13-Mar-2016; **Accepted:** 14-Mar-2016; **Online first:** 31-Mar-2016

Copyright: © 2016 Safia Beshir, Khadiga Salah Ibrahim, Weam Shaheen, Eman M. Shahy. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Competing Interests: The authors have declared that no competing interests exist.

BACKGROUND: Nickel exposure is recognized as an endocrine disruptor because of its adverse effects on reproduction.

AIM: This study was designed to investigate the possible testiculo-hormonal perturbations on workers occupationally exposed to nickel and to assess its effects on human male sexual function.

METHODS: Cross-sectional comparative study, comprising 105 electroplating male non-smoker, non-alcoholic workers exposed to soluble nickel and 60 controls was done. Serum luteinizing hormone, follicle stimulating hormone, testosterone levels and urinary nickel concentrations were determined for the studied groups.

RESULTS: Serum luteinizing hormone, follicle stimulating hormone, urinary nickel and the simultaneous incidence of more than one sexual disorder were significantly higher in the exposed workers compared to controls. The occurrence of various types of sexual disorders (decreased libido, impotence and premature ejaculation) in the exposed workers was 9.5, 5.1 and 4.4 folds respectively than the controls.

CONCLUSIONS: Exposure to nickel produces possible testiculo-hormonal perturbations in those exposed workers.

Introduction

There is increasing prevalence of various abnormalities in human male reproductive system [1]. It may be due to stress, lifestyle factors and presence of a variety of endocrine-altering environmental chemicals. Occupational activities may involve constant exposure to toxic agents and may have a detrimental effect on human reproduction [1]. As compared to other mammals, human males are of relatively low fertility and hence may be at a greater risk for toxicants influencing reproduction [2].

Nickel is a silver-white metallic chemical element that is naturally present in the Earth's crust [3]. Pure nickel metal is used in electroplating, as a chemical catalyst, and in the manufacture of alkaline batteries, coins, welding products, magnets, electrical

contacts and electrodes. Nickel salts are used in electroplating, ceramics, pigments, and as intermediates (e.g. catalysts, formation of other nickel compounds) [4, 5].

The mammalian male reproductive system can adversely be affected by nickel as shown in many experimental studies [6, 7]. Nickel-induced male reproductive toxic effects coupled with low dietary protein, induce severe changes, including altered physiologic functions, biochemical defects, and structural disorders [6, 7]. Experimental reports suggested that nickel exposure causes decrease in weight of testicular and accessory sex organs: epididymides, seminal vesicles and prostate gland, and decrease in testicular steroidogenic enzymes activities [7-9]. High doses of nickel have shown testicular toxicity involving oxidative stress in mice. This is evidenced by increased lipid peroxidation, DNA damage, and apoptosis in the testes,

morphological sperm head abnormalities, and decreased fertility [8, 9].

Relatively few data are available regarding the possible reproductive effects of nickel in human males. Some reports suggested significant positive correlation between high blood nickel level in welders and morphologically abnormal sperms [10, 11]. However, the involvement of oxidative stress mechanisms is the basic mode of action of nickel-induced toxicity in male reproductive dysfunction [12, 13]. Nickel exposure elevates testicular lipid peroxidation and suppresses antioxidant enzyme activities in rats [14]. Kakela and his colleagues [6] found that NiCl_2 induced shrinkage of seminiferous tubules and decreased the number of spermatogonia in the tubules in male rats. Free radical generation from the reaction of Ni-thiol complexes and molecular oxygen, and lipid hydroperoxides could play an important role in the mechanism(s) of Ni toxicity [15].

Nickel was recognized as an endocrine disruptor because of its adverse effect on reproduction [12] and disruption of steroidogenesis and spermatogenesis [16]. It may interfere with the reproductive hypothalamic hormones: luteinizing hormone (LH) and follicle stimulating hormone (FSH) as well as testosterone [17, 18].

The FSH and LH hormones are secreted by the anterior pituitary gland in response to gonadotropin-releasing hormone, produced by the hypothalamus. The secretion of FSH and LH is regulated through negative feedback by steroid hormones and by inhibin (a non-steroidal gonadal substance). In males, LH acts on the Leydig cells in the testes to stimulate the synthesis of testosterone. The action of FSH in maintaining spermatogenesis in the seminiferous tubules of the testes is augmented by LH and testosterone [19]. Alterations caused by endocrine disruptors can be temporary or permanent [20]. Endocrine disruptors can cause reproductive anomalies (morphological and functional gonadal dysfunction, e.g. infertility and decreased libido) and congenital mal-formations (altered embryonic and fetal intrauterine development) [21].

Occupational nickel exposure results in elevated levels of nickel in blood, urine and body tissues⁴. It is known as a potentially harmful element for humans. Its concentration in the environment can rise due to industrial activities [22, 23].

This study was designed to investigate the possible testiculo-hormonal perturbations on workers occupationally exposed to nickel and to assess its effects on human male sexual function.

Materials and Methods

The study was a cross-sectional comparative study, comprising 105 electroplating male non-smoker, non-alcoholic workers exposed to soluble nickel compound (by inhalation and dermal absorption) in a factory in Helwan district (Cairo, Egypt). Their age ranged from 31 to 60 yr (mean 41.13 ± 5.37) with duration of exposure ranging from 7 to 30 yr (15.41 ± 7.11). The control population included 60 apparently healthy non-smoker, non-alcoholic subjects from the National Research Centre not occupationally exposed to Nickel or other heavy metals, matched for age ranged from 29-59 yr (40.3 ± 6.14).

After taking consent from each individual, they were interviewed and completed a questionnaire that included personal data, detailed occupational history, and marital, sexual, surgical, medical and family histories. Five ml of venous blood sample were collected from all participants either exposed (105) or controls (60) during morning work shift by the authors of the current study. The collected venous blood samples were centrifuged at 3000 rpm for 10 min at 4°C, and the serum was collected and frozen at -20°C for later analysis of serum LH, FSH and testosterone. Urine specimen was collected in plastic containers previously acid washed by 1 M HNO_3 and rinsed extensively with water. Also, the urine was frozen and kept at -20°C until analyzed for urinary nickel and urinary creatinine (Cr).

Reproductive endocrine hormones were determined by immunosorbent assay (ELISA techniques) using commercial kits for serum testosterone (Biosource Europe SA Laboratories) and luteinising hormone (LH) (Elitech Diagnostics Laboratories) and follicle stimulating hormone (FSH) (Eurogenetics).

Urinary nickel was determined by the atomic absorption analysis in graphite furnace [24]. The standard addition technique was used for calibration. The nickel content of urine was expressed as micrograms per gram creatinine ($\mu\text{g/g Cr}$) excretion to compensate for an effect of muscle mass on Cr excretion²⁵. Determination of urinary Cr to adjust the values of the urinary nickel was carried out by using the Jaffe method without deproteinisation [26].

Data Analysis

The data were statistically analyzed using SPSS18 program. Independent t-test and Chi-square test were used to detect the statistical differences in the quantitative and qualitative data respectively between the two groups. Pearson's bivariate correlation coefficient was also calculated. The differences were considered significant at a level of $p < 0.05$.

Results

The urinary nickel and serum testosterone, LH and FSH levels of exposed and control subjects are given in Table 1. There were no significant differences between the control and the exposed groups regarding the age. There was a significant difference in urinary nickel level between the two studied groups. This was accompanied with significantly higher serum concentrations of LH and FSH. At the same time, no difference was found in the levels of serum testosterone.

Table 1: The characteristics of the studied groups

Parameters	Control (60) (Mean ± SD)	Exposed (105) (Mean ± SD)	P-value
Age (yr)	40.3 ± 6.14	41.13 ± 5.37	>0.05
Duration of exposure (yr)	-	15.41 ± 7.11	-
Urinary Ni (µg/g Cr)	1.67 ± 0.87	4.18 ± 1.50	<0.001
Serum Testosterone(ng/ml)	5.69 ± 1.91	5.44 ± 1.75	>0.05
Serum LH (mIU/ml)	4.47 ± 0.96	6.03 ± 1.23	<0.05
Serum FSH (mIU/ml)	7.19 ± 1.67	8.68 ± 2.47	<0.05

Table 2 shows that 35% of the controls and 42.8% of the exposed group complain of sexual disorders. In the exposed group, there was increase of simultaneous incidence of more than one sexual disorder (33.3%). The distribution of various types of sexual disorders (decreased libido, impotence, premature ejaculation) was higher in the exposed workers. It was found that an odd ratio of occurrence of those sexual disorders among the exposed group was 9.5, 5.1 and 4.4 times the controls respectively (Table 3).

Table 2: Incidence of sexual disorders in the studied groups

Parameters	Control (60)		Exposed (105)	
	No	%	No	%
Subjects with sexual disorders	21	35	45	42.8
Subjects complaining of one sexual disorder	15	25	10	9.5
Subjects complaining of more than one sexual disorder	6	10	35	33.3

Positive correlations were found between the levels of serum FSH and the duration of exposure ($r = 0.276$, $P < 0.05$) on one side, and the levels of urinary nickel ($r = 0.346$, $P < 0.01$) on the other side.

Table 3: Distribution of sexual disorders in the studied groups

Sexual symptoms	Control (60)		Exposed (105)		χ^2	95% CI		Odds ratio
	No	%	No	%		Lower limit	Upper limit	
Decreased libido	3	5	35	33.3	<0.0001	2.777	32.497	9.5
Impotence	6	10	38	36.2	<0.0002	2.008	12.971	5.1
Premature ejaculation	9	15	46	43.8	<0.0001	1.972	9.898	4.4
Infertility	3	5	8	7.6	>0.05			

Discussion

Nickel can be found practically in all environmental compartments. It originates from natural and artificial sources. The general population is exposed to nickel through nickel alloys and nickel-plated materials, such as coins, steel, and jewelry,

and residual nickel may be found in soaps, fats and oils [27]. The urinary nickel level relative to creatinine concentration in normal healthy adults is $<2 \mu\text{g/g}$ [28]. Occupational exposure is common for workers involved in mining, smelting, welding, casting, spray-painting and grinding, electroplating, production and use of nickel catalysts, polishing of nickel-containing alloys, and other jobs where nickel and nickel compounds are produced or used [5]. Nickel was higher in the urine of workers who were exposed to soluble nickel compounds than the workers exposed to less soluble compounds. The levels of nickel in biological fluids increase remarkably in persons with increased occupational or environmental exposure and decline rapidly when exposure is reduced or stopped. Thus, measurements of nickel, particularly in the urine, serum or hair, may serve as indices of exposure [29]. As there is a good correlation between the concentrations of nickel in the air and those present in the urine of exposed subjects, the measurement of urinary nickel represents the most appropriate test for the evaluation of an occupational exposure to this metal [30].

In our study a significant urine nickel levels of the exposed workers compared to the control group may be due to high levels of nickel fumes and fine particles in their work atmosphere. This result was in agreement with results of De Sio et al. [31] and Sancini et al. [32] studies.

A significant high concentration of LH and FSH with no difference in the levels of testosterone was found in the nickel exposed workers in our study, when compared to the control group. Elevated LH and FSH concentrations are sensitive indicators of Leyding and Sertoli cell failure. In many clinically apparent hypogonadism syndromes, LH and FSH are almost invariably raised even when testosterone concentrations are normal [33]. This finding could be explained by direct toxic action of nickel on the testes leading to reduced production of testosterone or by decreasing testicular sensitivity to gonadotrophic actions. In men, intact hypothalamo-pituitary axis is capable of compensating testosterone reduction by increasing FSH and LH secretions, bring back the testosterone production to the normal level.

The nickel interference with the reproductive hypothalamic hormones LH and FSH and the testosterone has been observed in vitro studies and studies on laboratory animals [34, 35]. These results were partly confirmed also in studies on human subjects exposed to nickel [17, 19].

In our study, positive correlations were found between the levels of serum FSH and the duration of exposure and the levels of urinary nickel. These findings were in agreement with that of De Sio et al. [31] who found a positive constant correlation between the values of urinary nickel and plasma FSH. The relationship between urinary nickel and the increase in plasma FSH depends on the action of this

metal. The endocrine disruption could be linked to the suppression of the neuroendocrine control in the testicles (with effects on the synthesis and release of testosterone), central nervous system (with effects on FSH, and LH), or both locations simultaneously [36, 37].

The nickel - exposed group in the present study showed significant increase in the incidence of sexual problems in the form of premature ejaculation, impotence and decreased libido, which are the most common male sexual problems as stated by Cleveland Clinic [38].

Testosterone is necessary to maintain male secondary sex characteristics, libido, and probably potency. Thus patients with endocrine abnormalities may present with variety of symptoms, elevated levels of the gonadotropins, FSH and LH in the presence of decreased testosterone levels indicating primary testicular dysfunction.

Future studies are warranted to overcome the current study's limitation of relatively small sample size.

As a conclusion to the present study, occupational exposure to nickel produces possible testiculo-hormonal perturbations in those workers. Most of the nickel occupationally exposed workers showed compensated primary hypogonadism (elevated LH and FSH values concurrent with normal testosterone concentration) that may be reversible if the exposure ceases.

Medical examination and measurement of urinary level of nickel are recommended periodically for workers occupationally exposed to nickel to detect any side effects or complications of exposure early. Those at high risk should be avoided from further exposure.

References

1. Toppari J, Larsen JC, Christiansen P, et al. Male reproductive health and environmental xenoestrogens. *Environ Health Perspect.* 1996;104 (4):741-803. <http://dx.doi.org/10.1289/ehp.96104s4741> PMID:8880001 PMCID:PMC1469672
2. Working PK. Male reproductive toxicology: comparison of the human to animal models. *Environ Health Perspect.* 1988;77:37-44. <http://dx.doi.org/10.1289/ehp.887737> PMID:3289906 PMCID:PMC1474524
3. Arita A, Niu J, Qu Q et al. Global levels of histone modifications in peripheral blood mononuclear cells of subjects with exposure to nickel. *Environ Health Perspect.* 2012; 120:198-203. <http://dx.doi.org/10.1289/ehp.1104140> PMID:22024396 PMCID:PMC3279455
4. IARC. Chromium, nickel and welding. IARC Monogr Eval Carcinog Risks Hum. 1990; 49:1-648. PMID:2232124
5. HSDB. Fact Sheet" Hazardous Substances Data Bank." National Library of Medicine. <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB> and search on CAS number. Last accessed: 10/22/2009.
6. Kakela R, Kakela A, Hyvarinen H. Effects of nickel chloride on reproduction of the rat and possible antagonistic role of selenium. *Comp Biochem Physiol C Pharmacol Toxicol Endocrinol.* 1999; 123:27-37. [http://dx.doi.org/10.1016/S0742-8413\(99\)00006-7](http://dx.doi.org/10.1016/S0742-8413(99)00006-7)
7. Obone E, Chakrabarti SK, Bai C et al. Toxicity and bioaccumulation of nickel sulfate in Sprague-Dawley rats following 13 weeks of subchronic exposure. *J Toxicol Environ Health.* 1999;57:379-401. <http://dx.doi.org/10.1080/009841099157593>
8. Das KK, Dasgupta S. Effect of nickel on testicular nucleic acid concentrations of rats on protein restriction. *Biol Trace Elem Res.* 2000;73:175-180. <http://dx.doi.org/10.1385/BTER:73:2:175>
9. Kong L, Tang M, Zhang T, et al. Nickel nanoparticles exposure and reproductive toxicity in healthy adult rats. *Int J Mol Sci.* 2014;15(11):21253-69. <http://dx.doi.org/10.3390/ijms151121253> PMID:25407529 PMCID:PMC4264223
10. Danadevi K, Rozati R, Reddy PP, et al. Semen quality of Indian welders occupationally exposed to nickel and chromium. *Reprod Toxicol.* 2003;17:451-456. [http://dx.doi.org/10.1016/S0890-6238\(03\)00040-6](http://dx.doi.org/10.1016/S0890-6238(03)00040-6)
11. Hjollund NH, Bonde JP, Jensen TK et al. Semen quality and sex hormones with reference to metal welding. *Reprod Toxicol.* 1998; 12(2):91-5. [http://dx.doi.org/10.1016/S0890-6238\(97\)00156-1](http://dx.doi.org/10.1016/S0890-6238(97)00156-1)
12. Stinson TJ, Jaw S, Jeffery EH, et al. The relationship between nickel chloride-induced peroxidation and DNA strand breakage in rat liver. *Toxicol Appl Pharmacol.* 1992;117:98-103. [http://dx.doi.org/10.1016/0041-008X\(92\)90222-E](http://dx.doi.org/10.1016/0041-008X(92)90222-E)
13. Chen CY, Huang YL, Lin TH. Association between oxidative stress and cytokine production in nickel-treated rats. *Arch Biochem Biophys.* 1998;356:127-132. <http://dx.doi.org/10.1006/abbi.1998.0761> PMID:9705202
14. Gupta AD, Dhundasi SA, Ambekar JG, et al. Effect of Lascorbic acid on antioxidant defense system in testes of albino rats exposed to nickel sulphate. *J Basic Clin Physiol Pharmacol* 2007; 18:87-95. <http://dx.doi.org/10.1515/JBCPP.2007.18.4.255>
15. Das KK, Buchner V. Effect of nickel exposure on peripheral tissues: Role of oxidative stress in toxicity and possible protection by ascorbic acid. *Rev Environ Health.* 2007; 22:133-49. <http://dx.doi.org/10.1515/REVEH.2007.22.2.157>
16. Sun Y, Ou Y, Cheng M et al. Binding of nickel to testicular glutamate-ammonia ligase inhibits its enzymatic activity. *Mol Reprod Dev.* 2011; 78(2):104-15. <http://dx.doi.org/10.1002/mrd.21275> PMID:21254280
17. Wirth JJ, Mijal RS. Adverse effects of low level heavy metal exposure on male reproductive function. *Syst Biol Reprod Med.* 2010;56(2):147-67. <http://dx.doi.org/10.3109/19396360903582216> PMID:20377313
18. Apostoli P, Catalani S. Metal ions affecting reproduction and development. *Met Ions Life Sci.* 2011;8:263-303. PMID:21473384
19. Pizent A, Tariba B, Živković T. Reproductive toxicity of metals in men. *Arh Hig Rada Toksikol.* 2012; 63(1):35-46. <http://dx.doi.org/10.2478/10004-1254-63-2012-2151>
20. Gronowski AM, Landau-Levine M. Reproductive endocrine function. In Burtis CA, Ashwood ER, eds. *Tietz Textbook of Clinical Chemistry* 5th ed. Philadelphia, Pa: WB Saunders, 1999:877-78.

21. Waissmann W. Health surveillance and endocrine disruptors. *Cad Saúde Pública*. 2002; 18:511-17.
<http://dx.doi.org/10.1590/S0102-311X2002000200016>
PMid:11923893
22. Nelson P. Epidemiology, biology, and endocrine disruptors. *Occup Environ Med*. 2003; 60:541-42.
<http://dx.doi.org/10.1136/oem.60.8.541>
PMid:12883013 PMCid:PMC1740595
23. Eliades T, Pratsinis H, Kletsas D et al. Characterization and cytotoxicity of ions released from stainless steel and nickel-titanium orthodontic alloys. *Am J Orthod Dentofacial Orthop*. 2004; 125:24–29.
<http://dx.doi.org/10.1016/j.ajodo.2003.09.009>
24. Long-zhu J, Zhe-ming N. Determination of nickel in urine and other biological samples by graphite furnace atomic absorption spectrometry. *Fresenius' Zeitschrift für analytische Chemie*. 1985; 321:72-76.
<http://dx.doi.org/10.1007/BF00464491>
25. Baxmann AC, Ahmed MS, Marques NC, et al. Influence of Muscle Mass and Physical Activity on Serum and Urinary Creatinine and Serum Cystatin C. *Clin J Am Soc Nephrol*. 2008; 3(2):348–54.
<http://dx.doi.org/10.2215/CJN.02870707>
PMid:18235143 PMCid:PMC2390952
26. Bartels H. Determination of serum and urinary creatinine by Jaffe's method without deproteinisation in a 2-point reaction rate measurement in 2 minutes. *Clin Chim Acta*. 1971; 32: 81-84.
27. ATSDR. Toxicological Profile for Nickel. Agency for Toxic Substances and Disease Registry. 1997.
<http://www.atsdr.cdc.gov/toxprofiles/tp15.pdf>. 293 pp
28. Agency for Toxic Substances and Disease Registry (ATSDR). Toxicological Profile for Nickel. Atlanta, GA: U.S. Department of Health and Human Services, Public Human Service, 2005.
29. Savolainen H. Biochemical and clinical aspects of nickel toxicity. *Rev Environ Health*. 1996; 11:167-73.
<http://dx.doi.org/10.1515/REVEH.1996.11.4.167>
PMid:9085432
30. Campurra G. Agenti chimici. In: *Manuale medicina del lavoro*. Italy: Wolters Kluwer, 2010; 868-69.
31. De Sio S, Casale T, Rosati MV et al. Correlation between urinary nickel and FSH plasma values in workers occupationally exposed to urban stressors. *Prevention and research*. 2013; 2 (4):131–40.
<http://dx.doi.org/10.11138/pr/2013.2.4.131>
32. Sancini A, De Sio S, Giofrè PA et al. Correlation between urinary nickel and testosterone plasma values in workers occupationally exposed to urban stressors. *Ann Ig*. 2014; 26(3):237-54.
PMid:24998215
33. Steinberger E. Current status of studies concerned with evaluation of toxic effects of chemicals on the testes. *Environ Health Perspect*. 1981; 38:29-33.
<http://dx.doi.org/10.1289/ehp.813829>
34. Lukac N, Bardos L, Stawarz R et al. In vitro effect of nickel on bovine spermatozoa motility and annexin V-labeled membrane changes. *J Appl Toxicol*. 2011; 31(2): 144-9.
PMid:20737413
35. Murawska-Ciałowicz E, Bal W, Januszewska L, et al. Oxidative stress level in the testes of mice and rats during nickel intoxication. *Scientific World Journal*. 2012; 2012: 395741
<http://dx.doi.org/10.1100/2012/395741>
PMid:22448131 PMCid:PMC3290116
36. Rodamilans M, Martinez-Osaba MJ, To-Figueras J, et al. Inhibition of intratesticular testosterone synthesis by inorganic lead. *Toxicol Lett*. 1988; 42 (3): 285-390.
[http://dx.doi.org/10.1016/0378-4274\(88\)90113-0](http://dx.doi.org/10.1016/0378-4274(88)90113-0)
37. Yang Y, Lu XS, Li DL, et al. Effects of environmental lead pollution on blood lead and sex hormone levels among occupationally exposed group in an E-waste dismantling area. *Biomed Environ Sci*. 2013; 26 (6): 474-84.
PMid:23816581
38. Sexual Dysfunction in Males - Cleveland Clinic
http://my.clevelandclinic.org/health/diseases_conditions/hic_An_Overview_of_Sexual_Dysfunction/hic_Sexual_Dysfunction_in_Males