



O Blood Group as Risk Factor for Preeclampsia among Sudanese Women

Abdelmageed Elmugabil¹, Duria A. Rayis², Mohamed A. Ahmed², Ishag Adam^{2*}, Gasim I. Gasim³

¹Faculty of Medicine, El Imam El Mahdi University, Kosti, Sudan; ²Faculty of Medicine, University of Khartoum, Khartoum, Sudan; ³Faculty of Medicine and Health Sciences, Alneelain University, Khartoum, Sudan

Abstract

Citation: Elmugabil A, Rayis DA, Ahmed MA, Adam I, Gasim GI. O Blood Group as Risk Factor for Preeclampsia among Sudanese Women. Open Access Maced J Med Sci. <http://dx.doi.org/10.3889/oamjms.2016.108>

Keywords: Preeclampsia; blood groups; Sudan; pregnancy.

***Correspondence:** Ishag Adam. Faculty of Medicine, University of Khartoum, P.O. Box 102, 11111, Khartoum, Sudan. Tel +249912168988. Fax +249183771211. E-mail: ishagadam@hotmail.com

Received: 06-Aug-2016; **Revised:** 21-Sep-2016; **Accepted:** 22-Sep-2016; **Online first:** 24-Sep-2016

Copyright: © 2016 Abdelmageed Elmugabil, Duria A. Rayis, Mohamed A. Ahmed, Ishag Adam, Gasim I. Gasim. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This research did not receive any financial support.

Competing Interests: The authors have declared that no competing interests exist.

AIM: To investigate blood groups and the other possible risk factors for preeclampsia among Sudanese women.

MATERIAL AND METHODS: A case – control study was conducted at Saad Abualila Hospital, Khartoum, Sudan during the period of July 2013 through December 2014. The cases were women with preeclampsia and healthy pregnant women were the controls.

RESULTS: Two hundred eighty pregnant women were enrolled (140 in each arm of the study). Around one-quarter of all women (280) were primiparae (74.0, 26.4%), the majority were housewives (201, 71.7%). Seventy-nine (28.2%) were illiterate or had no informal education. Around half of the women (130, 46.4%) had O blood group. Binary logistic regression showed association between preeclampsia and lack of antenatal care (OR = 2.75, 95% CI = 1.172–6.494, P = 0.020) as well as O blood group (OR = 1.78, 95% CI = 1.088–2.934, P=0.022).

CONCLUSION: The current study showed that women with blood group O were at higher risk of preeclampsia.

Introduction

Preeclampsia is a common complication of pregnancy, where its incidence is approximately one tenth of births [1]. It is a major cause of increased maternal and perinatal fatality worldwide [2]. The definite aetiology of preeclampsia remains ambiguous, nevertheless, previous researchers have delineated a number of risk factors including low education, primiparity, family history of hypertension, obesity, younger or advanced maternal age, and the ethnic background [3–7].

Some biochemical and haematological factors such as serum creatinine level, serum uric acid level, anaemia were also considered as predictors of preeclampsia [8, 9]. A Proper understanding of epidemiological and clinical risk factors of preeclampsia is essential for earlier diagnoses and

follow-up of women at higher risk of this disease. Most published researches on preeclampsia were carried at good resource settings with only a few exceptions [3].

Previous researchers exploring blood group in women with preeclampsia revealed different findings [10]. While some studies reported the associations of different blood groups with preeclampsia, other studies failed to reach similar conclusions [11–15]. There are an extremely high maternal mortality and morbidity in Sudan, where 4.2% out of all obstetric complications and about one fifth of maternal deaths are attributed to preeclampsia/eclampsia [16–18].

The current study was carried out to assess the possible risk factors for preeclampsia including blood groups. The data obtained is of paramount importance for the health planners, health providers, and supports the previous research on preeclampsia in Sudan [19–22].

Material and Methods

A case – control study was conducted at Saad Abualila Hospital, Khartoum, Sudan. Saad Abualila Hospital is a tertiary care hospital for women who attended the prenatal care at the hospital as well as for referrals from the other clinics and hospitals. All women with high-risk pregnancy or obstetric/medical complications are referred to the hospital. Medical files were reviewed for the women during the period of July 2013 through December 2014. A case was defined as a woman who had preeclampsia, which was defined as hypertension associated with proteinuria in a pregnant woman. Hypertension was defined as blood pressure ≥ 140 mmHg systolic or diastolic blood pressure ≥ 90 mmHg diastolic emerging after 20 weeks of gestation in a woman who was normotensive before this period. Proteinuria was defined as excretion of ≥ 300 mg of protein in 24 h urine sample or $\geq 2+$ on the dipstick. A consecutive file for each case was taken as control. The controls were pregnant women presented for delivery, with blood pressure values $< 139/89$ mmHg and no proteinuria recorded in the pregnant health card during the antenatal visits or at the time of delivery. Women with a twin pregnancy or diabetics were excluded because these are known predictors for preeclampsia [7].

The obstetric and medical history (age, parity, education, antenatal care, weight and height) were extracted from the files. Weight and height were used to calculate the body mass index, which was expressed as weight in kg/height in m^2 . Blood groups subtypes were recorded from the files of both the cases and the controls and expressed as O, A, B and AB.

A total sample size of 145 participants in each arm of the study was calculated to investigate the difference rate of the O blood group in the women with preeclampsia and the controls. We assumed the rate of O blood as 56% and 40% for preeclampsia women and the controls depending on our previous work in the same setting [22]. This rate would provide 80% power to detect a 5% difference at $\alpha = 0.05$, with an assumption that complete data might not be available for 12% of participants.

Statistics

SPSS for Windows (version 16.0) was used for data analyses. Studied variables were described with means (M) and standard deviations (SD). Proportions of the studied groups were expressed in percentages (%). The difference of mean (SD) and proportion was compared between the two studied groups using T-test and χ^2 , respectively. Binary logistic regression analyses were performed, where preeclampsia was the dependent variable and

medical/ obstetrics characteristics (age, parity, and residence), blood groups (O vs. non-O blood group), BMI were the independent variables. Odds ratio and 95% CI were calculated. $P < 0.05$ was considered statistically significant.

Ethics

The study received ethical clearance from the Research Board at the Department of Obstetrics and Gynaecology, Faculty of Medicine, University of Khartoum, Sudan

Results

Two hundred eighty pregnant women were enrolled (140 in each arm of the study). Around one-quarter of all women (280) were primiparae (74.0, 26.4%), the majority were housewives (201, 71.7%). Seventy-nine (28.2%) were illiterate or had no informal education. Around half of these women (130, 46.4%) had O blood group.

Table 1: Comparing the obstetrics and clinical characteristics of preeclamptic and controls women at Khartoum, Sudan

Variable	Preeclampsia (n = 140)	Controls (n = 140)	P
<i>The mean (SD)</i>			
Age, year	28.8 (6.2)	28.1 (6.0)	0.315
Parity	1.7 (2.1)	1.7 (1.4)	0.974
Body mass index, Kg/m ²	24.5 (3.0)	24.4 (2.2)	0.648
Number (%) of			
Education < secondary level	38 (27.1)	41 (29.3)	0.410
Lack of antenatal care	35 (25)	20 (14.3)	<0.001
O	86 (61.4)	64 (45.7)	
A	24 (17.1)	37 (26.4)	
B	27 (19.3)	26 (18.6)	0.007
AB	3 (2.1)	13 (9.3)	

There was no significant difference in the mean (SD) of the age, parity and BMI between women with preeclampsia and controls. Thirty-six (12.9%) and 104 (37.1) were severe and mild preeclampsia, respectively. Compared with the controls, significantly higher numbers of women with preeclampsia were illiterate and had O blood group, (Table 1, Fig. 1).




Figure 1: Percentage of blood group among preeclamptic and controls at Khartoum, Sudan

In binary logistic regression, women who had no antenatal care (OR = 2.75, 95%CI = 1.172–6.494, P = 0.020) and women with O blood group (OR = 1.78, 95%CI = 1.088–2.934, P = 0.022) were at higher risk of preeclampsia (Table 2).

Table 2: Binary logistic regression of the predictors of preeclampsia, Sudan

Variables	OR	95%CI	P
Age, year	1.05	1.001–1.113	0.048
Parity	0.88	0.739–1.050	0.158
Education < secondary level	1.19	0.647–2.203	0.571
Lack of antenatal care	2.75	1.172–6.494	0.020
Body mass index, Kg/m ²	0.98	0.897–1.090	0.821
O vs. none O blood group	1.78	1.088–2.934	0.022

Discussion

Perhaps this is the first study investigating the association between blood groups and preeclampsia in Africa. In this study, women who had no antenatal care were more likely to develop preeclampsia than those who had an antenatal follow-up. Previously, Bilano *et al.* described such a finding in their secondary analysis for the world health organisation facility based, a multi-country survey [23]. An explanation for such an association is the fact that antenatal visits help to detect and treat pregnancy-induced hypertension at an early stage before the development of complications.

An important finding of the current study is the association between blood group O and preeclampsia. This goes with the previous findings, where blood group O has been reported as a risk factor for preeclampsia among 386 cases and 342 controls [24]. Likewise, blood group A has been reported as a risk factor for preeclampsia [15].

A large Swedish cohort study showed that the blood group O is rather protective against preeclampsia [14]. Interestingly, a meta-analysis conducted by Franchini *et al.* demonstrated that the ABO system was consistently involved in preeclampsia. According to Franchini *et al.*, the non-O blood groups are risk factors while blood group O is protective against preeclampsia [10].

There is a great controversy about whether blood groups could have an association with preeclampsia. Scott *et al.* found no association between the ABO system and preeclampsia in their case-control study on a population of 46 patients [25]. However, Scott *et al.* study had relatively small sample size compared with the other studies that established an association between the ABO system and preeclampsia. Scott *et al.*, results are further supported by Witsenburg *et al.*, a report in 36 cases and 272 controls and Clark *et al.*, findings in 4250 pregnancies, 66 of which are preeclamptic women

[11, 12]. A separate population-based nested case-control study exploring 248 study cases demonstrated AB blood group as a risk factor for preeclampsia [13].

In conclusion, the current study showed that women with blood group O were at higher risk of preeclampsia.

References

- Robillard P-Y, Hulsey TC, Dekker GA, Chaouat G. Preeclampsia and human reproduction. An essay of a long term reflection. *J Reprod Immunol.* 2003; 59(2):93–100. [http://dx.doi.org/10.1016/S0165-0378\(03\)00040-8](http://dx.doi.org/10.1016/S0165-0378(03)00040-8)
- Duley L. The global impact of pre-eclampsia and eclampsia. *Semin Perinatol.* 2009;33(3):130–7. <http://dx.doi.org/10.1053/j.semperi.2009.02.010> PMID:19464502
- Conde-Agudelo A, Belizan JM. Risk factors for pre-eclampsia in a large cohort of Latin American and Caribbean women. *BJOG.* 2000;107(1):75–83. <http://dx.doi.org/10.1111/j.1471-0528.2000.tb11582.x> PMID:10645865
- Lee C-J, Hsieh T-T, Chiu T-H, Chen K-C, Lo L-M, Hung T-H. Risk factors for pre-eclampsia in an Asian population. *Int J Gynecol Obstet.* 2000;70(3):327–33. [http://dx.doi.org/10.1016/S0020-7292\(00\)00240-X](http://dx.doi.org/10.1016/S0020-7292(00)00240-X)
- Roberts JM, Pearson G, Cutler J, Lindheimer M. Summary of the NHLBI Working Group on research on hypertension during pregnancy. *Hypertension.* 2003;41(3 Pt 1):437–45.
- Sibai B, Dekker G, Kupferminc M. Pre-eclampsia. *Lancet.* 2005;365(9461):785–99. [http://dx.doi.org/10.1016/S0140-6736\(05\)71003-5](http://dx.doi.org/10.1016/S0140-6736(05)71003-5)
- Adeyinka DA, Oladimeji O, Adekanbi TI, Adeyinka FE, Falope Y, Aimakhu C. Outcome of adolescent pregnancies in southwestern Nigeria: a case-control study. *J Matern Fetal Neonatal Med.* 2010; 23(8):785–9. <http://dx.doi.org/10.3109/14767050903572166> PMID:20082596
- Ali AA, Rayis DA, Abdallah TM, Elbashir MI, Adam I. Severe anaemia is associated with a higher risk for preeclampsia and poor perinatal outcomes in Kassala hospital, eastern Sudan. *BMC Res Notes.* 2011;4(1):311. <http://dx.doi.org/10.1186/1756-0500-4-311> PMID:21867566 PMCID:PMC3224576
- Ndayambagye EB, Nakalembe M, Kaye DK. Factors associated with persistent hypertension after puerperium among women with pre-eclampsia/eclampsia in Mulago hospital, Uganda. *BMC Pregnancy Childbirth.* 2010;10:12. <http://dx.doi.org/10.1186/1471-2393-10-12> PMID:20222993 PMCID:PMC2848130
- Franchini M, Mengoli C, Lippi G. Relationship between ABO blood group and pregnancy complications: a systematic literature analysis. *Blood Transfus.* 2016;1–8. PMID:27483484
- Witsenburg CPJ, Rosendaal FR, Middeldorp JM, Van Der Meer FJM, Scherjon SA. Factor VIII levels and the risk of pre-eclampsia, HELLP syndrome, pregnancy related hypertension and severe intrauterine growth retardation. *Thromb Res.* 2005;115(5):387–92. <http://dx.doi.org/10.1016/j.thromres.2004.09.009> PMID:15733972
- Clark P, Walker ID, Govan L, Wu O, Greer IA. The GOAL study: A prospective examination of the impact of factor V Leiden and ABO(H) blood groups on haemorrhagic and thrombotic pregnancy outcomes. *Br J Haematol.* 2008;140(2):236–40. PMID:18028481
- Hiltunen LM, Laivuori H, Rautanen A, Kaaja R, Kere J, Krusius T, et al. Blood group AB and factor V Leiden as risk factors for pre-eclampsia: A population-based nested case-control study. *Thromb Res.* 2009;124(2):167–73. <http://dx.doi.org/10.1016/j.thromres.2008.11.012> PMID:19110300
- Lee BK, Zhang Z, Wikman A, Lindqvist PG, Reilly M. ABO and

RhD blood groups and gestational hypertensive disorders: a population-based cohort study. BJOG. 2012 Sep;119(10):1232–7.
<http://dx.doi.org/10.1111/j.1471-0528.2012.03421.x>
PMid:22734590

15. Phaloprakarn C, Tangjitgamol S. Maternal ABO blood group and adverse pregnancy outcomes. J Perinatol. 2013;33(2):107–11.
<http://dx.doi.org/10.1038/jp.2012.73> PMid:22678143

16. Leiberman JR, Fraser D, Kasis A, Mazor M. Reduced frequency of hypertensive disorders in placenta previa. Obstet Gynecol. 1991;77:83–86. PMid:1984232

17. Ali AA, Adam I. Lack of antenatal care, education, and high maternal mortality in Kassala hospital, eastern Sudan during 2005–2009. J Matern Fetal Neonatal Med. 2011;1077–8.
<http://dx.doi.org/10.3109/14767058.2010.545908> PMid:21231847

18. Ali AA, Okud A, Khojali A, Adam I. High incidence of obstetric complications in Kassala Hospital, Eastern Sudan. J Obstet Gynaecol (Lahore). 2012;32(2):148–9.
<http://dx.doi.org/10.3109/01443615.2011.637140> PMid:22296425

19. Bakheit KH, Bayoumi NK, Eltom AM, Elbashir MI, Adam I. Cytokines profiles in Sudanese women with preeclampsia. Hypertens Pregnancy. 2009;28(2):224–9.
<http://dx.doi.org/10.1080/10641950802601245> PMid:19437232

20. Bakheit KH, Ghebremeskel K, Pol K, Elbashir MI, Adam I. Erythrocyte omega-3 and omega-6 fatty acids profile in Sudanese women with pre-eclampsia. J Obstet Gynaecol. 2010;30(2):151–4.
<http://dx.doi.org/10.3109/01443610903391005> PMid:20143974

21. Bakheit KH, Ghebremeskel K, Zaiger G, Elbashir MI, Adam I. Erythrocyte antioxidant enzymes and plasma antioxidant vitamins in Sudanese women with pre-eclampsia. J Obstet Gynaecol. 2010;30(2):147–50. <http://dx.doi.org/10.3109/01443610903249448>
PMid:20143973

22. Adam I, Elhassan EM, Mohammed A a, Salih MM, Elbashir MI. Malaria and pre-eclampsia in an area with unstable malaria transmission in Central Sudan. Malar J. 2011;10(1):258.
<http://dx.doi.org/10.1186/1475-2875-10-258> PMid:21899731
PMCID:PMC3224261

23. Bilano VL, Ota E, Ganchimeg T, Mori R, Souza JP. Risk factors of pre-eclampsia/eclampsia and its adverse outcomes in low- and middle-income countries: a WHO secondary analysis. PLoS One. 2014 Jan;9(3):e91198.
<http://dx.doi.org/10.1371/journal.pone.0091198> PMid:24657964
PMCID:PMC3962376

24. Amin NS, Tahir SA, Abadi NA, Kubba K. Association of blood groups with preeclamptic toxemia. Med Sci Res. 1989;17:861–2.

25. Scott JR, Beer AE, Stastny P. Immunogenetic factors in preeclampsia and eclampsia. Erythrocyte, histocompatibility, and Y-dependent antigens. JAMA. 1976;235(4):402–4.
<http://dx.doi.org/10.1001/jama.1976.03260300028025> PMid:54446