

Multiple Primary Recurrent Basaliomas (mPR-BCCs) of the Scalp with Cranial Bone Invasion

Georgi Tchernev^{1*}, James W. Patterson², Torello Lotti³, Serena Gianfaldoni⁴, Jacopo Lotti⁵, Katlein França⁶, Atanas Batashki⁷, Uwe Wollina⁸

¹Medical Institute of Ministry of Interior (MVR), Department of Dermatology and Dermatologic Surgery, General Skobelev 79, 1606 Sofia, Bulgaria; ²Onkoderma - Policlinic for Dermatology, Venereology and Dermatologic Surgery, 26 General Skobelev blvd., Sofia, Bulgaria; ³Department of Pathology, University of Virginia Health System, 1215 Lee Street, Box 800214, Charlottesville, VA 22908, USA; ⁴Chair of Dermatology, University of Rome G. Marconi, Rome 00186, Italy; ⁵Dermatology and Venereology, University G. Marconi of Rome Rome, Rome 00192, Italy; ⁶Department of Nuclear, Subnuclear and Radiation Physics, University of Rome "G. Marconi", Rome, Italy; ⁷Institute for Bioethics & Health Policy, Department of Dermatology & Cutaneous Surgery; Department of Psychiatry & Behavioral Sciences, University of Miami Miller School of Medicine - Miami, FL, USA; ⁸Abdominal and Thoracic Surgery, Department of Special Surgery, Medical University of Plovdiv, bul. "Peshtersko shose" Nr 66, 4000 Plovdiv, Bulgaria; ⁸Department of Dermatology and Allergology, Academic Friedrichstrasse 41, 01067, Dresden, Germany

Abstract

Citation: Tchernev G, Patterson JW, Lotti T, Gianfaldoni S, Lotti J, França K, Batashki A, Wollina U. Multiple Primary Recurrent Basaliomas (mPR-BCCs) of the Scalp with Cranial Bone Invasion. Open Access Maced J Med Sci. 2017 Jul 25; 5(4):564-565. <https://doi.org/10.3889/oamjms.2017.145>

Keywords: basal cell carcinoma; cranial bone invasion; adequate therapy; surgery; selctrosurgery.

***Correspondence:** Professor Dr Georgi Tchernev PhD, Chief of 1) Department of Dermatology, Venereology and Dermatologic Surgery, Medical Institute of Ministry of Interior (MVR), General Skobelev 79, 1606 Sofia; 2) Onkoderma- Policlinic for Dermatology and Dermatologic Surgery, General Skobelev 26, Sofia, Bulgaria. GSM: 00359885588424. E-mail: georgi.tchernev@yahoo.de

Received: 06-Apr-2017; Revised: 22-Apr-2017; Accepted: 23-Apr-2017; Online first: 25-Jul-2017

Copyright: © 2017 Georgi Tchernev, James W. Patterson, Torello Lotti, Serena Gianfaldoni, Jacopo Lotti, Katlein França, Atanas Batashki, Uwe Wollina. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

Funding: This research did not receive any financial support.

Competing Interests: The authors have declared that no competing interests exist.

We present a 68-year-old patient with multiple primary infiltrative BCCs in the scalp area, initially treated 14 years ago with superficial contact X-ray therapy, end dose 60 greys, followed by electrocautery (x2) several years later. He presented in the dermatologic policlinic for diagnosis and therapy of two additional, newly-formed pigmented lesions, and because of an uncomfortable, itchy, burning sensation in the area where lesions had been treated years before. Screening cranial computer-tomography (CT) examination revealed two deformities in the form of tumor-mediated osteolysis, affecting the diploe of the tabula externa on the left parietal and parasagittal areas. Complete excision with removal of periosteum and partial removal of the tabula externa was planned with neurosurgeons at a later stage. BCC is one of the most common malignant skin tumours of the head and neck region (about 90% of cases) and is characterised by a significant potential for local infiltration and destructive growth. Recurrent, invasive BCC of the scalp and calvarium is a difficult problem for which universally accepted treatment protocols had not been established. The primary treatment of aggressive BCCs is surgical, with a thorough examination of excision margins to ensure complete resection. Procedural-based options include standard excision, curettage, curettage with electrodesiccation, and Mohs micrographic surgery (MMS), with MMS being the gold standard for the definitive treatment of BCC. Improper removal or electrocautery (as in our case) of the several aggressive forms of BCC seems to be a particular problem, and not only for dermatologic surgeons. The risk of subsequent invasion and destruction of the cranium, underlying dura, and cranial nerves by basal cell carcinoma (BCC) is extremely low, with an estimated incidence of 0.03%, but is a potential complication over time. Computed tomography is the modality of choice for detecting tumour invasion into bone, which commonly appears as irregular demineralization or osteolysis.

We present a 68-year-old patient with multiple primary infiltrative BCCs in the scalp area, initially treated 14 years ago with superficial contact X-ray therapy, end dose 60 greys, followed by electrocautery (x2) several years later (Fig. 1a). He presented in the dermatologic policlinic for diagnosis and therapy of two additional, newly-formed pigmented lesions, and because of an uncomfortable, itchy, burning sensation in the area where lesions had been treated years before (Fig. 1a-d).

Screening cranial computer-tomography (CT) examination revealed two deformities in the form of

tumor-mediated osteolysis, affecting the diploe of the tabula externa on the left parietal and parasagittal areas. Complete excision with removal of periosteum and partial removal of the tabula externa was planned with neurosurgeons at a later stage. BCC is one of the most common malignant skin tumours of the head and neck region (about 90% of cases) and is characterised by a significant potential for local infiltration and destructive growth [1]. Recurrent, invasive BCC of the scalp and calvarium is a difficult problem for which universally accepted treatment protocols had not been established [2]. The primary

treatment of aggressive BCCs is surgical, with a thorough examination of excision margins to ensure complete resection [3]. Procedural-based options include standard excision, curettage, curettage with electrodesiccation, and Mohs micrographic surgery (MMS), with MMS being the gold standard for the definitive treatment of BCC [4].




Figure 1: Lesions of the scalp in our patient. 1a) Ulcerated lesions - see horizontally oriented arrows. Horizontally oriented arrows also show histopathologically verified infiltrative BCCs that had been treated in the past. Vertically oriented arrows show newly pigmented BCCs; 1b-d) Horizontally oriented arrows show the older BCCs, treated in the past via radiation and electrodesiccation

Improper removal or electrocautery (as in our case) of the several aggressive forms of BCC seems to be a particular problem, and not only for dermatologic surgeons. The risk of subsequent invasion and destruction of the cranium, underlying dura, and cranial nerves by basal cell carcinoma

(BCC) is extremely low, with an estimated incidence of 0.03%, but is a potential complication over time [5]. Computed tomography is the modality of choice for detecting tumour invasion into bone, which commonly appears as irregular demineralization or osteolysis [5].

References

1. Rasulić LG, Jovanović MD. Surgical treatment and dilemmas in the treatment of basal cell carcinomas with intracranial propagation. *Vojnosanit Pregl.* 2014;71(11):1045-8. <https://doi.org/10.2298/VSP1411045R> PMID:25536808
2. Cho GJ, Wang F, Garcia SM, Viner J, Hoffman WY, McDermott MW, Pomerantz JH. Recalcitrant Invasive Skin Cancer of the Scalp: Combined Extirpation and Microsurgical Reconstruction Without Cranioplasty. *J Craniofac Surg.* 2017;28(2):325-330. <https://doi.org/10.1097/SCS.0000000000003384> PMID:28045814
3. Naumann IC, Cordes SR. Giant basal cell carcinoma of the forehead with extensive intracranial involvement. *Ann Otol Rhinol Laryngol.* 2007;116(9):663-6. <https://doi.org/10.1177/000348940711600907> PMID:17926588
4. Blackmon J, Machan M, Rajpara A, Beatty R. Recurrent basal cell carcinoma with intracranial invasion: a case report and literature review. *Dermatol Online J.* 2014;20(7). pii: 13030/qt11t0g5h8.
5. Kleydman Y, Manolidis S, Ratner D. Basal cell carcinoma with intracranial invasion. *J Am Acad Dermatol.* 2009;60(6):1045-9. <https://doi.org/10.1016/j.jaad.2008.10.005> PMID:19467376