

# Medium Sized Congenital Melanocytic Nevus with Suspected Progression to Melanoma during Pregnancy: What's the Best for the Patient?

Georgi Tchernev<sup>1,2\*</sup>, Gabriela Atanasova Dzhelyatova<sup>3</sup>, Uwe Wollina<sup>4</sup>, Ilia Lozev<sup>5</sup>, Torello Lotti<sup>6</sup>

<sup>1</sup>Medical Institute of Ministry of Interior (MVR), Department of Dermatology, Venereology and Dermatologic Surgery, Sofia, Bulgaria; <sup>2</sup>"Onkoderma" - Polyclinic for Dermatology and Dermatologic Surgery, Sofia, Bulgaria; <sup>3</sup>Medical Institute of the Ministry of Interior, Dermatology and Dermatologic Surgery, Sofia, Sofia, Bulgaria; <sup>4</sup>Städtisches Klinikum Dresden - Department of Dermatology and Allergology, Dresden, Sachsen, Germany; <sup>5</sup>Medical Institute of Ministry of Interior Department of General, Vascular and Abdominal Surgery, Sofia, Bulgaria; <sup>6</sup>University G. Marconi of Rome - Dermatology and Venereology, Rome, Italy;

## Abstract

**Citation:** Tchernev G, Dzhelyatova GA, Wollina U, Lozev I, Lotti T. Medium Sized Congenital Melanocytic Nevus with Suspected Progression to Melanoma during Pregnancy: What's the Best for the Patient? Open Access Maced J Med Sci. <https://doi.org/10.3889/oamjms.2018.016>

**Keywords:** Congenital melanocytic nevus (CMN); Pregnancy; Melanoma; Malignant transformation

**\*Correspondence:** Georgi Tchernev. Department of Dermatology, Venereology and Dermatologic Surgery, Medical Institute of Ministry of Interior (MVR-Sofia), General Skobelev 79, 1606 Sofia, Bulgaria; Onkoderma - Polyclinic for Dermatology, Venereology and Dermatologic Surgery, General Skobelev 26, 1606, Sofia, Bulgaria. E-mail: [georgi\\_tchernev@yahoo.de](mailto:georgi_tchernev@yahoo.de)

**Received:** 20-Aug-2017; **Revised:** 17-Sep-2017; **Accepted:** 24-Sep-2017; **Online first:** 13-Jan-2018

**Copyright:** © 2018 Georgi Tchernev, Gabriela Atanasova Dzhelyatova, Uwe Wollina, Ilia Lozev, Torello Lotti. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

**Funding:** This research did not receive any financial support

**Competing Interests:** The authors have declared that no competing interests exist

**BACKGROUND:** Congenital melanocytic nevi (CMN) are pigmented skin lesions usually present at birth. Rare varieties can develop and become clinically very large. Although they are benign nevocytic neoplasms, all CMN may be precursors of the melanoma, regardless of their size. Individual risk of malignant transformation of melanocyte is determined by simultaneous action of exogenous and endogenous factors. The major exogenous risk factor is ultraviolet radiation. Leading roles among the endogenous factors are attributed to skin phenotype, gene mutation, sex hormones and their significance.

**CASE REPORT:** We present a case of a 27 – year – old pregnant female patient with a congenital melanocytic nevus, which increased significantly in size, during her pregnancy. Estrogen levels increase during pregnancy and clinical evidence has suggested that melanocytes are estrogen – responsive. Nevi in a pregnant patient would exhibit increased expression of estrogen receptor  $\beta$  (ER $\beta$ ) and thus enhanced the potential to respond to altered estrogen levels.

**CONCLUSION:** All pigmented skin lesions should be carefully observed during pregnancy by a dermatologist due to the increased risk of malignant transformation, associated with the endocrine dependence. All lesions with visible changes should be removed surgically with appropriate anaesthesia.

## Introduction

Despite aesthetic, congenital melanocytic nevi can cause health problems. Usually they are classified by size: small (< 1.5 cm in diameter), medium (1.5 – 19.9 cm) and large or giant ( $\geq 20$ ) [1]. Independently of their size, all congenital melanocytic nevi are associated with increased risk of development of melanoma [1]. The risk of malignant transformation is higher in giant congenital nevi, and they should be carefully monitored biopsied if indicated [2].

The most prominent and predictable progression could be seen in the middle sized melanocytic nevi by dermoscopic and clinical evaluation, because:

1. The giant congenital melanocytic nevi often show areas which are clinically and dermoscopically difficult to differentiate from melanoma [3]. In these cases (patients with giant congenital melanocytic nevi) surgical excision is rarely due to the enlarged size of the lesions [4][5]. Confocal laser dermoscopy and PET CT can be useful to diagnose melanoma [6].

2. Small congenital melanocytic nevi or so-

called congenital pseudomelanomas are often clinically and dermoscopically indistinguishable from real melanomas. Histopathological verification of the above-mentioned lesions are also subjected to lively discussions; therefore differentiation of melanoma is extremely difficult.

These two facts are giving a new perspective on diagnosis and choosing of the most appropriate treatment option for the medium-sized melanocytic nevi, namely by surgery [3][4][5]. Progression of normal and dysplastic nevi to melanoma during pregnancy is an interesting topic which at the moment does not find a definitive solution [7].

## Case report

We present a case of a 27 – year - old female patient, with a pigmented lesion measuring 3 x 5 cm, located above the right gluteal area since early childhood. The lesion was asymptomatic and had not shown any changes in size or colour for the last 20 years. There was no evidence of significant comorbidities or medical treatment. During pregnancy, the patient noticed peripheral enlargement of the lesion as well as the intensification of the dark hue. The latest changes prompted the patient to seek medical consultation at the dermatological clinic. A large melanocytic nevus was established within the clinical examination, located above the right gluteal area with asymmetric shape, uneven boundaries at the periphery, no uniform colour in the different areas of the lesion as well as the difference in diameter – east, west, north, south, but no elevation of the lesion.



Figure 1: a, b) Clinical view of the lesion located above the right gluteal area; c, d) Consecutive stages within the excision of the lesion

The diagnosis of medium-sized congenital melanocytic nevus was confirmed by the medical history, dermoscopic and clinical signs of dysplasia

and progression during pregnancy. The lesion was surgically removed under local anaesthesia (Fig.1 a, b, c, d). The histopathological evaluation concluded the diagnosis of medium-sized congenital pigmented congenital melanocytic nevus with minimal cytological atypia and clear surgical margins.

Since pregnancy is a sure risk factor for the progression of normal nevi to dysplastic or dysplastic nevi to melanoma, we recommend surgical treatment as a preferable option.

## Discussion

It is well known that the frequency and prognosis of melanoma in women are influenced by hormonal and reproductive factors [7]. It is also well established that the prognosis and survival rate in premenopausal women is better than postmenopausal [8]. In the last years there has been increased interest and discussion about the impact of pregnancy on nevi and their malignant transformation [7][9][10]. New theories and approaches have been advanced to explain the interplay between hormones and pathological changes in nevi [11]. One of the hypotheses is the influence of estrogen expression. Beneficial and protective effects on the skin have estrogen receptors: estrogen – receptor  $\alpha$  (ER $\alpha$ ) and estrogen - receptor  $\beta$  (ER $\beta$ ) [12]. Significant differences in the concentrations of these receptors have been established in sections of melanocytic lesions and those with healthy skin as well as in pregnant and non- pregnant women's skin [13]. Subtype  $\beta$  is a predominant receptor in melanocytes and its protective function is well known [12]. ER $\beta$  is antagonist against uncontrolled cell- proliferation and tumor growth [12][14]. An increased in the immunoreactivity for ER $\beta$  was observed in normal nevi during pregnancy [15]. The immunoreactivity for ER $\beta$  was found to decreases with such deeply extending cells [16][17]. Loss of ER $\beta$  expression and its presumed inhibitory effects may promote transformation into melanoma, which is a key event in neoplastic progression [18]. Several studies show reduced expression of ER $\beta$  in metastatic stages of malignant melanoma [19], in the presence of a greater thickness of the dysplastic nevi [20]. The clinical implications of such altered ER $\beta$  expressions remains underestimated. Different hypothesis explains the higher risk of malignancy during pregnancy with the increased levels of male sex hormones- androgens [21]. There is a theory that endocrine effect reduces after first pregnancy. During second and third pregnancy the risk of development of malignant melanoma (MM) is lower due to the presence of antibodies against tumor-associated fetal antigens. Thus, during first pregnancy the risk of malignant transformation is increased, while every subsequent

pregnancy has a protective effect [22]. Recently, mutations in two tumor suppressive genes - BAP1 (BRCA - associated protein 1) and BRAF (V - raf murine sarcoma virus oncogene homolog B1) have been associated with increased susceptibility for development of MM and other atypical epithelial lesions [23][24]. Screening for mutation/loss/inactivation of BAP1 and BRAFV600E can be performed by immunohistochemistry. Most melanocytic lesions show positive BAP1 nuclear staining. BRAFV600E is positive in 5% of congenital melanocytic nevus [23][25]

The potential relationship between dysplastic nevi and malignant transformation during pregnancy is underestimated [26][27]. However, all pigment skin lesions should be carefully observed during this period [26][27]. In our case of a 27 – years - old pregnant woman, with CMN, which significantly increased its size and changed its colour and therefore, we decide to remove the lesions surgically, because of the increased risk of malignant transformation.

## References

- Eds. Bologna J, Jorizzo J, Schaffer J, *Dermatology* 3rd ed. Elsevier Sanders. 2012; P 1871-6.
- Viana ACL, Gontijo B, Bittencourt FV. Giant congenital melanocytic nevus. *An Bras Dermatol*. 2013; 88(6): 863-78. <https://doi.org/10.1590/abd1806-4841.20132233> PMID:24474093 PMCID:PMC3900335
- Chokoeva AA, Fioranelli M, Rocchia MG, Lotti T, Wollina U, Tchernev G. Giant congenital melanocytic nevus in a bulgarian newborn. *J Biol Regul Homeost Agents*. 2016; 30(2 Suppl 2): 57-60. PMID:27373137
- Goldman A, Wollina U, Tchernev G, Chokoeva AA, Lotti T. Medium-sized congenital melanocytic nevus of the forehead, glabella and temple – surgical treatment and long-term follow-up. *J Biol Regul Homeost Agents*. 2016; 30(2 Suppl 2): 53-8. PMID:27373136
- Chokoeva AA, Tchernev G, Trayanova E, Patterson JW, Lotti T, Wollina U. Giant congenital melanocytic nevus localized in the axillary area: serial excisions as optimal treatment option. *J Biol Regul Homeost Agents*. 2015; 29(1 Suppl):123-8. PMID:26016980
- Chokoeva A, Tchernev G. [Significance of the clinical and dermatoscopic findings, histopathology and reflex confocal microscopy (RCM) in determining the origin of melanocytic lesions: an analysis based on a clinical case]. *Akush Ginekol (Sofia)*. 2014; 53(6):56-62.
- Trayanova E, Chokoeva AA, Tchernev G, Patterson JW, Wollina U, Lotti T. Dysplastic nevi, melanoma and pregnancy- where is the relationship? *J Biol Regul Homeost Agents*. 2015; 29(1 Suppl): 87-90. PMID:26016974
- Daryanani D, Plukker JT, De Hullu JA, Kuiper H, Nap RE, Hoekstra HJ. Pregnancy and early- stage melanoma. *Cancer*. 2003; 97: 2248-53. <https://doi.org/10.1002/cncr.11321> PMID:12712479
- Brenner S, Tamir E. Early detection of melanoma: the best strategy for a favorable prognosis. *Clin Dermatol*. 2002; 20(3): 203-11. [https://doi.org/10.1016/S0738-081X\(02\)00233-X](https://doi.org/10.1016/S0738-081X(02)00233-X)
- Neil SM, Eves P, Richardson B, Molife R, Lorigan P, Wagner M, Layton C, Morandini R, Ghanem G. Oestrogenic steroids and melanoma cells influence invasion of melanoma cells in vitro. *Pigment Cell Res*. 2000; (13S8): 68-72.
- Pennoyer JW, Grin CM, Driscoll MS, et al. Changes in size of melanocytic nevi during pregnancy. *J Am Acad Dermatol*. 1997; 36: 378-382. [https://doi.org/10.1016/S0190-9622\(97\)80212-5](https://doi.org/10.1016/S0190-9622(97)80212-5)
- de Giorgi V, Gori A, Grazzini M, Rossari S, Scarfi F, Corciova S, Verdelli A, Lotti T, Massi D. Estrogens, estrogen receptors and melanoma. *Expert Rev Anticancer Ther*. 2011; 11(5):739-47. <https://doi.org/10.1586/era.11.42> PMID:21554049
- Grill HJ, Benes P, Manz B, Schramm P, Morsches B, Korting GW, Pollow K. Steroid hormone receptors in human melanoma. *Arch Dermatol Res*. 1982; 272:97-101. <https://doi.org/10.1007/BF00510399> PMID:7165327
- Folkerd EJ, Downsett M. Influence of sex hormones on cancer progression. *J Clin Oncol*. 2010; 28:4038-44. <https://doi.org/10.1200/JCO.2009.27.4290> PMID:20644089
- Schmidt AN, Nanney LB, Boyd AS, King LE, Jr, Ellis DL. Oestrogen receptor- beta expression in melanocytic lesion. *Exp Dermatol*. 2006; 15: 971-780. <https://doi.org/10.1111/j.1600-0625.2006.00502.x> PMID:17083364
- Kincannon J, Boutzale C. The physiology of pigmented nevi. *Pediatrics*. 1999; 104:1042-1045. PMID:10506262
- Krengel S. Nevogenesis- new thoughts regarding a classical problem. *Am J Dermatopathol*. 2005; 27:456-465. <https://doi.org/10.1097/01.dad.0000175532.27368.3f> PMID:16148419
- Slominski A, Wortsman J, Tuckey RC, Paus R. Differential expression of HPA axis homolog in the skin. *Mol Cell Endocrinol*. 2007; 266:143-149. <https://doi.org/10.1016/j.mce.2006.12.012> PMID:17197073 PMCID:PMC1839836
- De Giorgi V, Mavilia C, Massi D, et al. Estrogen receptor expression in cutaneous melanoma: a real-time reverse transcriptase-polymerase chain reaction and immunohistochemical study. *Arch Dermatol*. 2009; 145:30-6. <https://doi.org/10.1001/archdermatol.2008.537> PMID:19153340
- De Giorgi V, Gori A, Gandini S, et al. Estrogen receptor beta and melanoma. A comparative study. *Br J Dermatol*. 2013; 168:513-9. <https://doi.org/10.1111/bjd.12056> PMID:23013061
- Rampen F. Malignant melanoma: sex differences in survival after evidence of distant metastasis. *Br J Cancer*. 1980; 42(1):52-7. <https://doi.org/10.1038/bjc.1980.202> PMID:7426329 PMCID:PMC2010477
- Smith MA, Fine JA, Barnhill RL, Berwick M. Hormonal and reproductive influences and risk of melanoma in women. *Int J Epidemiol*. 1998; 27(5):751-7. <https://doi.org/10.1093/ije/27.5.751> PMID:9839729
- Piris A, Mihm MC Jr, Hoang MP, BAP1 and BRAFV600E expression in benign and malignant melanocytic proliferation. *Hum. Pathol*. 2015; 46(2):239-45. <https://doi.org/10.1016/j.humpath.2014.10.015> PMID:25479927
- Wang A, Papneja A, Hycza M, Al-Habeeb A, Ghazarian D, Gene of the month: BAP1. *J Clin Patol*. 2016; 69(9):750-3. <https://doi.org/10.1136/jclinpath-2016-203866> PMID:27235536
- Busam KJ, Sung J, Wiesner T, von Deimling A, Jungbluth A. Combined BRAF(V600E)-positive melanocytic lesions with large epithelioid cells lacking BAP1 expression and conventional nevomelanocytes. *Am J Surg Pathol*. 2013; 37(2):193-9. <https://doi.org/10.1097/PAS.0b013e318263648c> PMID:23026932
- Trayanov I, Trayanova E, Chokoeva A, Tchernev G. [Congenital melanocytic nevus of the shoulder with rapid growth progression during pregnancy. Successful surgical approach]. *Akush Ginekol (Sofia)*. 2015; 54(8):51-6.
- Trayanova E, Trayanov I, Chokoeva A, Tchernev G. [Malignant melanoma--the influence of hormonal factors on the progression and prognosis. Comparative analysis based on two clinical cases]. *Akush Ginekol (Sofia)*. 2015; 54(7):45-51.