

# Correlation of Procalcitonin and C-Reactive Protein with Intra-Abdominal Hypertension in Intra-Abdominal Infections: Their Predictive Role in the Progress of the Disease

Bedri Braha<sup>1</sup>, Dafina Mahmutaj<sup>1</sup>, Mehmet Maxhuni<sup>1</sup>, Burim Neziri<sup>2</sup>, Shaip Krasniqi<sup>2\*</sup>

<sup>1</sup>Clinic of Surgery, University Clinical Center of Kosovo, Prishtina, Kosovo; <sup>2</sup>Faculty of Medicine, University of Prishtina "Hasan Prishtina", Prishtina, Kosovo

## Abstract

**Citation:** Braha B, Mahmutaj D, Maxhuni M, Neziri B, Krasniqi S. Correlation of Procalcitonin and C-Reactive Protein with Intra-Abdominal Hypertension in Intra-Abdominal Infections: Their Predictive Role in the Progress of the Disease. Open Access Maced J Med Sci. <https://doi.org/10.3889/oamjms.2018.112>

**Keywords:** Intra-abdominal infection; Intra-abdominal hypertension; Procalcitonin; C - reactive protein

**\*Correspondence:** Shaip Krasniqi. Faculty of Medicine, University of Prishtina "Hasan Prishtina", Prishtina, Kosovo. E-mail: [shaip.krasniqi@uni-pr.edu](mailto:shaip.krasniqi@uni-pr.edu)

**Received:** 11-Oct-2017; **Revised:** 23-Nov-2017; **Accepted:** 25-Nov-2017; **Online first:** 05-Mar-2018

**Copyright:** © 2018 Bedri Braha, Dafina Mahmutaj, Mehmet Maxhuni, Burim Neziri, Shaip Krasniqi. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

**Funding:** This research did not receive any financial support

**Competing Interests:** The authors have declared that no competing interests exist

**AIM:** To analyse the correlation of procalcitonin (PCT) and C-reactive protein (CRP) values with increased intra-abdominal pressure and to evaluate their predictive role in the progression of Intra-abdominal infections.

**MATERIALS AND METHODS:** A non-randomized prospective study conducted in the group of 80 patients. We have measured the PCT, CRP and intra-abdominal pressure (IAP).

**RESULTS:** According to IAH grades (G), there was a significant difference of PCT values: G I  $3.6 \pm 5.1$  ng/ml, G II  $10.9 \pm 22.6$  ng/ml, G III  $15.2 \pm 30.2$  ng/ml ( $p = 0.045$ ) until CRP values were increased in all IAH groups but without distinction between the groups: GI  $183 \pm 64.5$ , GII  $196 \pm 90.2$ , GIII  $224 \pm 96.3$  ( $p = 0.17$ ). According to the severity of the infection, we yielded increased values of PCT, IAP and CRP in septic shock, severe sepsis and SIRS/sepsis resulting in significant differences of PCT and IAP.

**CONCLUSION:** Based on the results of our research, we conclude that the correlation of PCT values with IAH grades is quite significant while the CRP results remain high in IAH but without significant difference between the different grades of IAH.

## Introduction

Intraabdominal infections are defined as peritoneal inflammation in response to microorganisms [1]. Despite the achievements in the diagnosis, management, and proper treatment of patients with intra-abdominal infections the presence of secondary peritonitis reflects the critical prognosis, which has a high mortality rate of up to 30% [2] [3] [4] [5] [6]. Early prognosis estimation of complicated intra-abdominal infections is important in evaluating the severity of the disease since intra-abdominal infections might advance to severe sepsis and septic shock [4]. It is often associated with hypoperfusion followed by tissue damage and organ dysfunction following increased severity of systemic response to infection [7] [8] [9]. There are some promising biomarkers for sepsis that are currently used for relatively few diagnostic purposes [10]. Studies are aimed to find a single marker at low cost, that can be

measured quickly, to repeat measurement easy, and that measurement does not cause disturbances in patients [11]. From this group of biomarkers, CRP and PCT are most commonly used in clinical practice [10]. CRP is the acute phase protein synthesised in the liver and is released after inflammation and tissue damage [12] [13].

In many studies, the role of PCT in the diagnosis and treatment of systemic and local infections was investigated, including intra-abdominal infections [14]. PCT in secondary peritonitis has improved diagnostic value for identification of patients with severe sepsis endangered by septic shock [15]. Bacterial infection is a strong PCT stimulus, while viral systemic inflammation results in PCT values to be low [14][16]. In recent years, there was expressed interest in intra-abdominal hypertension (IAH) that is defined as a pathological increase of intra-abdominal pressure above 12 mmHg [17]. IAH is determined by the elasticity of the abdominal wall/diaphragm, the volume

of internal organs and the presence of ascites, blood, and intestinal contents. Some abdominal clinical conditions (peritonitis, ileus, volvulus, pancreatitis, complications after abdominal surgery) and systemic conditions (sepsis, mechanical ventilation, pneumonia, and burns) found to be related with IAH [17].

The purpose of this prospective study is to compare the values of PCT and CRP with IAP grades to Intra-abdominal infections and determination of the correlation between them, and correlation of these parameters about the severity of the infection, which points to their predictive role in evaluating the prognosis of the disease.

## Materials and Methods

This is a prospective, controlled clinical trial involving hospitalised patients at the University Clinical Center of Kosovo in Pristina, registered in the period from July 2016 to February 2017. The research was conducted at the University Clinical Center of Kosovo, Pristina, where 80 patients were hospitalised as abdominal cases in which Intra-abdominal hypertension was present. The research protocol has been approved by the Ethics Committee within the University of Prishtina – Faculty of Medicine and the consent form is completed by all participants or their family members.

**Inclusion Criteria:** Age over 18 years, the presence of SIRS due to Intra-abdominal infections, the presence of intra-abdominal hypertension, consent from the patient's relatives.

**Exclusion criteria:** Urinary trauma, urinary bladder injury, immunosuppression, chronic kidney failure, liver cirrhosis, SIRS is not present, long-term use of corticoids, acute hepatic insufficiency, diabetes.

### Protocol and Treatment

The baseline criterion during the research was the measurement of the IAP before the operation every four hours, and it was evaluated as IAH if the average of these measurements was  $> 12$  mmHg. The measurement was performed even after the operation, where the measurement was performed three times in 24h, and the mean was calculated for each day. The duration of the measurement has continued until the normalisation of the IAP. The Intra-abdominal pressure measurement was done by the method evaluated by Kron. PCT and CRP measurement were performed before surgery, after the 1<sup>st</sup>, 4<sup>th</sup> and 7<sup>th</sup> days and the following days after the surgery if necessary. The PCT marker level measurement is done with the Elecsys BRAHMS

method. Laboratory measurements and analyses that are relevant to the designation of IAH / ACS (Intra-abdominal Hypertension / Abdominal Compartment Syndrome), MODS (Multiple Organ Dysfunction Score), APACHE II (Acute Physiology and Chronic Health Evaluation), SOFA (Sequential Organ Failure Assessment), IMP (Index Mannheim Peritonitis), MAP (Mean Arterial Pressure) were realized. The fluid balance in the body is determined based on the weight, ordinated juices, and diuresis APP (APP = MAP - IAP) before and after surgery, FG (FG = MAP - 2IAP) before and after surgery. Based on the clinical, laboratory and general findings, the severity of Intra-abdominal infection has been classified into three groups: SIRS/sepsis, severe sepsis and septic shock. At the same time, based on the values of IAP, patients are grouped into three groups: Grade I: IAP 12 - 15 mmHg, Grade II: IAP 16 - 20 mmHg, Grade III: IAP 21 - 25 mmHg. There are also cases involving ACS in GIII. In the realised measurements, we did not have cases with IAP  $> 25$  mm Hg. The PCT and CRP relationships with IAP grades, APP, FG, diuresis and fluid balance also were analysed in correlation with these parameters according to the severity of the infection.

### Statistical Analysis

Statistical data processing was performed with the statistical package SPSS 22.0. The arithmetic average, the standard deviation, the minimum and the maximum values were calculated. Spearman correlation coefficient was used for non-parametric data. Qualitative data testing was done with the Pearson's chi-squared test ( $\chi^2$ ) and the exact Fisher test of quantitative data that had a normal distribution with T-test and One Way ANOVA, while those with non-normal distribution with the Mann-Whitney test or Kruskal Wallis test. The difference is significant if  $P < 0.05$ .

## Results

In the study, procalcitonin and CRP were analysed about IAH of diagnoses of different structures with Intra-abdominal infections.

In Table 1, we presented the patient's related characteristics and the diagnosis structure of these patients about the three grades of IAH. In the total cohort of patients, male patients were dominant in comparison to females (54 vs 26). The average age of the participants was 49.4 years (SD  $\pm$  18.3 years), ranging from 18 to 56 years. The results showed a significant difference ( $P < 0.003$ ) between the age of the patients, the gender and the degree of IAH. First-degree hypertension was registered in 46 or 57.5%,

second degree in 24 or 30.0% and third degree in 10 or 12.5% patients. Regarding the diagnosis structure, acute perforated appendicitis with diffuse peritoneal diffuse has dominated all degrees of IAH. In cases with SIRS/sepsis, the presence of GI hypertension was more common in severe sepsis, and septic shock. GII and GIII of IAH were more pronounced. Abdominal Compartment Syndrome was present at 4 (5%) cases (Table 1).

**Table 1: Characteristics of the patients involved in the study**

	Intra-abdominal Hypertension			
	G I	G II	G III	P
Age	43.3 ± 17.5	57.3 ± 17.5	58.2 ± 14.2	0.003 <sup>a</sup>
Female	15 (57.7%)	6 (23.1%)	5 (19.2%)	0.366 <sup>b</sup>
Male	31 (57.4%)	18 (33.3%)	5 (9.3%)	0.366 <sup>b</sup>
Appendicitis acuta gangrenosa perforativa.	21	6	4	
Peritonitis diffusa fibro purulenta				
Cholecystitis gangrenosa	2	2	-	
Dehiscencia suttura duodeni ,peritonitis diffusa fibropurulenta	1	-	-	
Duodenopancreatocomia cefalica,dehiscencia anastomosis pancreato-jejunalis	-	-	1	
Hernia ventralis per magna.Incarceratio intestine ilei et perforation ilei	-	-	1	
Dehiscencia entero - entero anastomosis	4	5	2	
Perforatio diverticulum Meckeli,Peritonitis seropurulenta	-	1	-	
Perforatio in loci GEA,Peritonitis diffusa	1	-	-	
Perforation liberal	4	2	-	
Perforatio sigmae,Colitis ulcerosa,peritonitis diffusa stercoralis	1	-	-	
St post hysterectomiam,Peritonitis fibropurulenta	-	1	-	
Tu perforans colonias colonias	-	1	1	
Ulcus bulbi duodeni perforans.	10	3	1	
Peritonitis fibropurulenta	-	1	-	
Ulcus ventriculi perforans abscessus multiples intestini	-	1	-	
Vulnus sclopetarium abdominis,Laesio intestine ,peritonitis difusa	2	2	-	
SIRS/Sepsis	41 (70.69%)	14 (24.14%)	3 (5.17%)-ACS(n=1)	
Severe Sepsis	4 (25%)	6 (37.5%)	6 (37.5%)-ACS(n=2)	
Septic Shock	1 (16.7%)	4 (66.6%)	1 (16.7%)-ACS(n=1)	

a - Kruskal Wallis test, b- X<sup>2</sup> - test.


The PCT and mortality rates determined by the scales: APACHE, SOFA, IMP, MODS before surgery, which have been observed in higher results for IAH patients. All these parameters were statistically significant by IAH grades. CRP results to be increased at all levels of IAH but without statistical significance among groups. MODS was 0 % in 19 patients out of which 3 were with GII and 16 were with GI of IAH, at the same time, PCT in these cases was < 2 ng / ml except for only one case with PCT values of 10 ng/ml. SOFA mortality was < 10 % in 18 patients, 14 with GI and 4 with GII of IAH, and PCT > 2 ng/ml in one patient. The GII, GIII / ACS, values of FG, APP, diuresis and fluid balance were significantly reduced, and we have distinguished significant statistical significance of these parameters by IAH grades (Table 2).

**Table 2: Predictive parameters in admission in accordance with intra - abdominal pressure degrees**

	Abdominal Urgencies n = 80			P
	GI (n = 46)	GII (n = 24)	GIII (n = 6)/CSA (n = 4)	
PCT	3.6 ± 5.1	10.9 ± 22.6	15.2 ± 30.2	0.045 <sup>a</sup>
CRP	183.7 ± 64.5	196.3 ± 90.2	224.4 ± 96.3	0.17 <sup>a</sup>
FG	56.0 ± 10.8	48.7 ± 25	37.1 ± 16.2	0.005 <sup>b</sup>
APP	69.1 ± 10.8	66.1 ± 24.2	56.3 ± 17.3	0.096 <sup>b</sup>
Diuretics and fluid balance	1.0 ± 0.5	0.6 ± 0.4	0.4 ± 0.5	<0.0001 <sup>a</sup>
SOFA-mortality	11.1 ± 14.2	28.8 ± 27.0	39.6 ± 24.9	0.0001 <sup>a</sup>
APACHE II- mortality	12.3 ± 14.7	31.8 ± 26.9	40.4 ± 22.8	<0.0001 <sup>a</sup>
IMP- mortality	15.0 ± 12.7	29.3 ± 24.2	38.6 ± 22.4	0.003 <sup>b</sup>
MODS- mortality	7.3 ± 12.3	19.3 ± 22.5	26.2 ± 29.8	0.0004 <sup>a</sup>


A - Kruskal Wallis test; b - One Way ANOVA. PCT - procalcitonin, CRP - C reactive protein, FG - Filtration gradient, APP - abdominal perfusion pressure, SOFA - Sequential Organ Failure Assessment, APACHE - Acute Physiology and Chronic Health Evaluation IMP - Index Mannheim peritonitis, MODS - Multiple Organ Dysfunction Score.

The analysis of the values of IAP and PCT in all cases involved in the research results with a significant positive correlation of a medium scale (r = 0.42, P < 0.0001) (Fig. 1).



**Figure 1: Correlation between intra-abdominal pressure and PCT values**

A significant positive correlation of a middle degree was observed with the correlation of Sperman (r = 0.44, P < 0.0001) between IAP and CRP values (Fig. 2).



**Figure 2: Correlation between intra-abdominal pressure and CRP values**

Intra-abdominal pressures, analysed according to the severity of the infection, resulting in higher values for all groups. However, the statistical difference lies within the category of the infection before surgery and the following 4 - day post-surgery period. Even in PCT, we gained statistical significance by infection category before surgery, in day 1 and 4 post - surgery. CRP values were increased in all groups, but there was no significant difference according to the severity of the infection (Table 3).

**Table 3: Correlation of PCT and CRP with IAH according to the severity of the infection**

	Abdominal Urgencies			P
	Septic Shock	Severe Sepsis	SIRS/sepsis	
IAP before Surgery	18.4 ± 2.8	18.3 ± 3.2	14.7 ± 2.3	0.0002 <sup>a</sup>
IAP after surgery - day 1	16.4 ± 4.0	16.4 ± 3.2	11.8 ± 2.1	<0.0001 <sup>b</sup>
IAP after surgery - day 2	16.5 ± 4.1	14.76 ± 3.6	10.1 ± 2.3	<0.0001 <sup>a</sup>
IAP after surgery - day 3	15.8 ± 0.97	12.2 ± 3.71	9.8 ± 2.06	0.0031 <sup>a</sup>
IAP after surgery - day 4	17.62 ± 8.37	11.31 ± 6.25	9.45 ± 0.72	0.0500 <sup>a</sup>
PCT before surgery	49.2 ± 46.6	9.3 ± 12.1	3.1 ± 4.6	0.0002 <sup>a</sup>
PCT after surgery - day 1	42.4 ± 48.2	6.5 ± 7.6	2.1 ± 2.6	0.0024 <sup>a</sup>
PCT after surgery - day 4	43.3 ± 37.3	3.3 ± 3.4	0.8 ± 0.8	<0.0001 <sup>a</sup>
CRP before surgery	189.9 ± 114.4	187.0 ± 90.2	140.6 ± 73.5	0.1447 <sup>a</sup>
CRP after surgery - day 1	210.3 ± 117.1	124.7 ± 80.2	115.1 ± 64.7	0.894 <sup>a</sup>
CRP after surgery - day 4	124.2 ± 98.2	101.2 ± 48.1	97.1 ± 51.8	0.264 <sup>a</sup>

a - Kruskal Wallis test; One Way ANOVA test.

## Discussion

In this clinical study, PCT and CRP correlation with IAH and their predictive role in evaluating intra-abdominal infections with peritonitis were evaluated. Our findings indicate a significant positive correlation of a middle degree ( $r = 0.44$ ,  $P < 0.0001$ ) between IAH and PCT values of intra-abdominal infections. Medium-scale correlation was also found between CRP and IAH in intra-abdominal infections ( $r = 0.42$ ,  $P < 0.0001$ ).

Moreover, we have investigated the correlation of PCT, CRP, APACHE II, MODS, SOFA, IMP, APP, FG, diuresis and fluid balance about IAH grades. The results showed a statistical significance of these parameters, and exposure of IAH yielded a predictive role of PCT and IAH in the course of the disease. There is relevant evidence that proves the PCT correlation with IAH, involving only acute pancreatitis. Berenau et al., analysed the correlation between maximum IAP values and mean PCT values in acute pancreatic patients, which was highly significant [18]. Bettina M. Rau et al., [19] has established conclusion that pancreatic patients with acute pancreatitis, with infected pancreatic necrosis and MODS or those who have died have significant association with PCT's high values compared to complications, while Bezmarevic concludes that in the first 24 hours acceptance of acute pancreatitis to determine the severity of the disease PCT has higher sensitivity and similarity to APACHE II while CRP has low sensitivity in this regard, at the same time the growth of IAP is followed by increased PCT, so there is a correlation between these parameters and have a predictive role of disease progression [20].

In our study, the PCT values were high in severe sepsis and septic shock patients. High PCT values in septic shock were also found in other studies [21] [22] [23]. Despite that CRP values are high in all IAH groups of our study; there is no significant difference between groups. Many authors conclude that patients with a severe degree of IAH manifest, evident clinical changes in cardiovascular function [24] [25] [26], splanchnic hypoperfusion [27] and acute renal failure [28] [29]. The high values of the IAP result in organ dysfunction, which results in high values of the MODS escalation [30] [31]. The development of purulent peritonitis increases the IMP degrees that correspond to the growth of the IAP [32]. In other studies, results showed that the IAH presence was associated with significant increases in the SOFA scaling indicating that the IAH has negative effects not only on the abdomen but also on the function of other organs [33] at the same time the high values of intra-abdominal pressure are in correlation with high APACHE II values [33] and complied to the data from our study. Studies conducted to confirm that patients with IAH and high scores of SOFA and APACHE II have a worse clinical prognosis for hospital treatment

and intensive care [33] [34] [35] [37]. Taking into account the correlation of APACHE II with PCT it has been concluded that the increase in values of this marker is related to the severity of the disease and is the early marker of sepsis [29] APACHE II shows significant PCT correlation in dysfunctional and non-dysfunctional patients, whereas CRP correlation with APACHE II was not significant [28]. Similar data are also provided by Lopez et al. [34].

In our study of GII, GIII/ACS we had a decrease in the GF, APP, diuretic and fluid balance values and we have gained a distinction of statistical significance of these parameters by the IAH grades. Even in other studies, patients who did not survive had high levels of IAP and low values of APP, diuresis and fluid balance and FG [33] [37] [38]. In the research work of Kovac, it turns out that APP has a significant correlation with MODS, APACHE II and SIRS until the IAP shows no correlation with these parameters [32].

PCT and IAP transmitted before and after the operation resulted in higher values of probability of septic shock and severe sepsis have a significant difference in these parameters according to the severity of the infection, indicating that the deterioration of the overall state of the average of these parameters grows. CRP values are increased, but there is no significant difference according to the severity of the infection. Nargis findings show that PCT is more predictive than CRP in identifying and assessing the severity of sepsis even though both markers cannot differentiate infectious and non-infectious clinical syndromes [39]. In our study, we also found that PCT is more reliable in assessing the severity of the disease. PCT remains elevated in patients with severe sepsis and septic shock; this increases the risk of mortality regardless of the level [15] [39] [40].

The study performed by Sameh in 25 patients diagnosed as SIRS, sepsis, severe sepsis and septic shock with aetiology from all organs, analysed the PCT and CRP values and its data resulted that CRP does not show any degree of differentiation of severity of sepsis [41]. Similar data was also given in the study conducted by Castelli [42] that corresponds to the data from our research.

In conclusion, the increase in IAP is followed by increased PCT and CRP values. In this study, we have identified the correlation in the values of PCT and CRP to intra-abdominal infections where IAH is present. PCT results are high in all cases with IAH, and there is a significant statistical significance by IAH grades in the first 24 hours. However, CRP results are high but without a significant difference to IAH grades. High PCT and IAP values are evident in severe sepsis and septic shocks that are correlated with low FG, APP, diuresis and fluid balance and high mortality rate scores. The role of PCT in intra-abdominal infections is of greater diagnostic and predictive value compared

to CRP values; however, the role of CRP remains important for the identification of other clinical circumstances.

## References

- Wittmann DH, Schein M, Condon RE. Management of secondary peritonitis. *Ann Surg.* 1996; 224(1):10–8. <https://doi.org/10.1097/0000658-199607000-00003> PMID:8678610 PMCID:PMC1235241
- Malangoni MA, Inui T. Peritonitis – the Western experience. *World J Emerg Surg [Internet].* 2006;1(1):25. <https://doi.org/10.1186/1749-7922-1-25> PMID:16953882 PMCID:PMC1592073
- Menichetti F, Sganga G. Definition and Classification of Intra-abdominal Infections. *J Chemother.* 2009; 21(Suppl 1):3–4. <https://doi.org/10.1179/joc.2009.21.Supplement-1.3> PMID:19622444
- Pieracci FM, Barie PS. Management of Severe Sepsis of Abdominal Origin. *Scand J Surg.* 2007; 96(3):184–96. <https://doi.org/10.1177/145749690709600302> PMID:17966743
- Mulari K, Leppäniemi A. Severe Secondary Peritonitis following Gastrointestinal Tract Perforation. *Scand J Surg.* 2004; 93(3):204–8. <https://doi.org/10.1177/145749690409300306> PMID:15544075
- Mulier S, Penninckx F, Verwaest C, Filez L, Aerts R, Fieuws S, et al. Factors Affecting Mortality in Generalized Postoperative Peritonitis: Multivariate Analysis in 96 Patients. *World J Surg.* 2003; 27(4):379–84. <https://doi.org/10.1007/s00268-002-6705-x> PMID:12658477
- Hotchkiss RS, Karl IE. The Pathophysiology and Treatment of Sepsis. *N Engl J Med.* 2003; 348(2):138–50. <https://doi.org/10.1056/NEJMr021333> PMID:12519925
- Annane D, Bellissant E, Cavaillon J-M. Septic shock. *Lancet (London, England).* 2005; 365(9453):63–78. [https://doi.org/10.1016/S0140-6736\(04\)17667-8](https://doi.org/10.1016/S0140-6736(04)17667-8)
- Weigard M, Horner C, Bardenheuer H. The SIRS. *Best Pr Res Clin Anaesthesiol.* 2004; 18(3):455–75.
- Pierrakos C, Vincent J-L. Sepsis biomarkers: a review. *Crit Care.* 2010; 14(1):R15. <https://doi.org/10.1186/cc8872> PMID:20144219 PMCID:PMC2875530
- Marshall J, Innes M. Intensive care unit management of intra-abdominal infection. *Crit Care Med.* 2003; 31(8):2228–2237. <https://doi.org/10.1097/01.CCM.0000087326.59341.51> PMID:12973184
- Póvoa P, Almeida E, Moreira P, Fernandes A, Mealha R, Aragão A, et al. C-reactive protein as an indicator of sepsis. *Intensive Care Medicine.* 1998; 24:1052–1056. <https://doi.org/10.1007/s001340050715> PMID:9840239
- Póvoa P. C-reactive protein: a valuable marker of sepsis. *Intensive Care Med.* 2002; 28(3):235–43. <https://doi.org/10.1007/s00134-002-1209-6> PMID:11904651
- Mathew B, Roy D D, Kumar TV. The use of procalcitonin as a marker of sepsis in children. *J Clin Diagn Res.* 2013; 7(2):305–7. <https://doi.org/10.7860/JCDR/2013/4739.2753>
- Pupelis G, Drozdova N, Mukans M, Malbrain ML. Serum procalcitonin is a sensitive marker for septic shock and mortality in secondary peritonitis. *Anestezijol Intens Ter.* 2014; 46(4):262–73. <https://doi.org/10.5603/AIT.2014.0043> PMID:25293477
- Pfister R, Kochanek M, Leygeber T, Brun-Buisson C, Cuquemelle E, Machado MB, et al. Procalcitonin for diagnosis of bacterial pneumonia in critically ill patients during 2009 H1N1 influenza pandemic: a prospective cohort study, systematic review and individual patient data meta-analysis. *Crit Care.* 2014; 18(2):R44. <https://doi.org/10.1186/cc13760> PMID:24612487 PMCID:PMC4056761
- Papavramidis TS, Marinis AD, Pliakos I, Kesisoglou I, Papavramidou N. Abdominal compartment syndrome - Intra-abdominal hypertension: Defining, diagnosing, and managing. *J Emerg Trauma Shock.* 2011; 4(2):279–91. <https://doi.org/10.4103/0974-2700.82224> PMID:21769216 PMCID:PMC3132369
- Bereanu AS, Sava M. The evolution of the values of procalcitonin and intra-abdominal pressure in severe acute pancreatitis. *Acta Medica Transilvanica.* 2016; 21(1).
- Rau BM, Kemppainen EA, Gumbs AA, Büchler MW, Wegscheider K, Bassi C, et al. Early assessment of pancreatic infections and overall prognosis in severe acute pancreatitis by procalcitonin (PCT): a prospective international multicenter study. *Ann Surg.* 2007; 245(5):745–54. <https://doi.org/10.1097/01.sla.0000252443.22360.46> PMID:17457167 PMCID:PMC1877072
- Bezmarevic M, Mirkovic D, Soldatovic I, Stamenkovic D, Mitrovic N, Perisic N, Marjanovic I, Mickovic S KM. Correlation between procalcitonin and intra-abdominal pressure and their role in prediction of the severity of acute pancreatitis. *Pancreatology.* 2012; 12(4):337–43. <https://doi.org/10.1016/j.pan.2012.05.007> PMID:22898635
- Rau BM, Kemppainen EA, Gumbs AA, Büchler MW, Wegscheider K, Bassi C, et al. Early assessment of pancreatic infections and overall prognosis in severe acute pancreatitis by procalcitonin (PCT): a prospective international multicenter study. *Ann Surg.* 2007; 245(5):745–54. <https://doi.org/10.1097/01.sla.0000252443.22360.46> PMID:17457167 PMCID:PMC1877072
- Brunkhorst F, Wegscheider K, Forycki Z, Brunkhorst R. Procalcitonin for early diagnosis and differentiation of SIRS, sepsis, severe sepsis and septic shock. *Crit Care.* 1999; 3(Suppl 1):P095.
- Nakamura A, Wada H, Ikejiri M, Hatada T, Sakurai H, Matsushima Y, et al. Efficacy of procalcitonin in the early diagnosis of bacterial infections in a critical care unit. *Shock.* 2009; 31(6):587–92. <https://doi.org/10.1097/SHK.0b013e31819716fa> PMID:19060784
- Cheatham ML, Malbrain MLNG. Cardiovascular implications of abdominal compartment syndrome. *Acta Clin Belg.* 2007; 62(Suppl1):98–112. <https://doi.org/10.1179/acb.2007.62.s1.013>
- Quintel M, Pelosi P, Caironi P, Meinhardt JP, Luecke T, Herrmann P, et al. An Increase of Abdominal Pressure Increases Pulmonary Edema in Oleic Acid-induced Lung Injury. *Am J Respir Crit Care Med.* 2004; 169(4):534–41. <https://doi.org/10.1164/rccm.200209-1060OC> PMID:14670801
- Ameloot K, Gillebert C, Desie N MM. Hypoperfusion, Shock States, and Abdominal Compartment Syndrome (ACS). *Surg Clin North Am.* 2012; 92(2):207–20. <https://doi.org/10.1016/j.suc.2012.01.009> PMID:22414408
- Cresswell AB, Wendon JA. function and non-invasive hepatosplanchnic monitoring in patients with abdominal hypertension. *Acta Clin Belg.* 2007; 62(Suppl 1):113–8. <https://doi.org/10.1179/acb.2007.62.s1.014>
- De Laet IE. Current insights in intra-abdominal hypertension and abdominal compartment syndrome. *Med Intensiva.* 2007; 31(2):88–99. [https://doi.org/10.1016/S0210-5691\(07\)74781-2](https://doi.org/10.1016/S0210-5691(07)74781-2)
- Dalfino L, Tullo L, Donadio I, Malcangi V, Brienza N. Intra-abdominal hypertension and acute renal failure in critically ill patients. *Intensive Care Med.* 2008; 34(4):707–13. <https://doi.org/10.1007/s00134-007-0969-4> PMID:18157662
- Kovač N, Širanović M, Mazul-Sunko B. Clinical significance of intra-abdominal pressure and abdominal perfusion pressure in patients with acute abdominal syndrome. *Signa vitae.* 2007; 2(2):14–7. <https://doi.org/10.22514/SV22.102007.3>
- De Waele JJ, De laet I. Intra-Abdominal Hypertension and MODS. In: *Sepsis Management.* Berlin, Heidelberg: Springer Berlin Heidelberg, 2012:59–71. [https://doi.org/10.1007/978-3-642-03519-7\\_6](https://doi.org/10.1007/978-3-642-03519-7_6)

32. Plaudis H, Rudzats A, Melberga L, Kazaka I, Suba O, Pupelis G. Abdominal negative-pressure therapy: a new method in countering abdominal compartment and peritonitis - prospective study and critical review of literature. *Ann Intensive Care*. 2012; 2(Suppl 1):S23. <https://doi.org/10.1186/2110-5820-2-S1-S23> PMID:23281649 PMCID:PMC3527158
33. Vidal MG, Weisser JR, Gonzalez F, Toro MA, Loudet C, Balasini C, et al. Incidence and clinical effects of intra-abdominal hypertension in critically ill patients. *Crit Care Med*. 2008; 36(6):1823–31. <https://doi.org/10.1097/CCM.0b013e31817c7a4d> PMID:18520642
34. López F. Procalcitonin (PCT), C reactive protein (CRP) and its correlation with severity in early sepsis. *Clin Rev Opin*. 2011; 3(3):26–31.
35. Pelosi P, Quintel M, Malbrain MLNG. Effect of Intra-abdominal pressure on respiratory mechanics. *Acta Clin Belg*. 2007; 62(Suppl 1):78–88. <https://doi.org/10.1179/acb.2007.62.s1.011>
36. Dornikova G. Diagnostic and prognostic value of procalcitonin and C-reactive protein: systematic review. 22nd European Congress of Clinical Microbiology and infectious diseases (ECCMID), 2012. PMID:22973655
37. Cheatham ML, Safcsak K. Is the evolving management of intra-abdominal hypertension and abdominal compartment syndrome improving survival?. *Crit Care Med*. 2010; 38(2):402–7. <https://doi.org/10.1097/CCM.0b013e3181b9e9b1>
38. Regueira T, Bruhn A, Hasbun P, Aguirre M, Romero C, Llanos O, et al. Intra-abdominal hypertension: incidence and association with organ dysfunction during early septic shock. *J Crit Care*. 2008; 23(4):461–7. <https://doi.org/10.1016/j.jcrc.2007.12.013> PMID:19056007
39. Nargis W, Ibrahim M, Ahamed BU. Procalcitonin versus C-reactive protein: Usefulness as biomarker of sepsis in ICU patient. *Int J Crit Illn Inj Sci*. 2014; 4(3):195–9. <https://doi.org/10.4103/2229-5151.141356> PMID:25337480 PMCID:PMC4200544
40. Harbarth S. Diagnostic Value of Procalcitonin, Interleukin-6, and Interleukin-8 in Critically Ill Patients Admitted with Suspected Sepsis. *Am J Respir Crit Care Med*. 2001; 164(3):396–402. <https://doi.org/10.1164/ajrccm.164.3.2009052> PMID:11500339
41. Maraghi S EL, Yehia A, Mahmoud O, Abd Hamid A EL, El Maraghi S. Procalcitonin Versus C-Reactive Protein at Different SOFA Scores in I.C.U. Sepsis: Diagnostic Value and Therapeutic Implications. *Med J Cairo Univ*. 2014; 82(1):29–36.
42. Castelli GP, Pognani C, Meisner M, Stuani A, Bellomi D, Sgarbi L. Procalcitonin and C-reactive protein during systemic inflammatory response syndrome, sepsis and organ dysfunction. *Crit Care*. 2004; 8(4):R234-42. <https://doi.org/10.1186/cc2877> PMID:15312223 PMCID:PMC522844