

Mathematical Model for Forecasting the Influence of Atmospheric Pollution on Population Morbidity in Stara Zagora Municipality (Bulgaria)

Magdalena Platikanova^{1*}, Petya Hristova², Hristina Milcheva³

¹Trakia University, Medical Faculty, Hygiene, Epidemiology and Infectious Diseases, 11 Armeyska, Stara Zagora 6000, Bulgaria; ²Trakia University, Medical Faculty, Stara Zagora, Bulgaria; ³Trakia University, Medical Faculty, Stara Zagora, Bulgaria

Abstract

Citation: Platikanova M, Hristova P, Milcheva H. Mathematical Model for Forecasting the Influence of Atmospheric Pollution on Population Morbidity in Stara Zagora Municipality (Bulgaria). Open Access Maced J Med Sci. 2018 May 20; 6(5):934-939. <https://doi.org/10.3889/oamjms.2018.205>

Keywords: Atmospheric pollution; Morbidity; Mathematical model; Atmospheric pollutants; PM₁₀

***Correspondence:** Magdalena Platikanova. Trakia University, Medical Faculty, Hygiene, Epidemiology and Infectious Diseases, 11 Armeyska, Stara Zagora 6000, Bulgaria. E-mail: mplatikanova@abv.bg

Received: 19-Feb-2018; **Revised:** 11-Apr-2018; **Accepted:** 12-Apr-2018; **Online first:** 18-May-2018

Copyright: © 2018 Magdalena Platikanova, Petya Hristova, Hristina Milcheva. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Funding: This research did not receive any financial support

Competing Interests: The authors have declared that no competing interests exist

AIM: This paper aims to create a mathematical model for forecasting the morbidity of the population in the Republic of Bulgaria and the Stara Zagora Municipality in particular as a consequence of the atmospheric pollution.

SUBJECTS AND METHODS: This model is based on a formula which determines the correlation between the average annual concentrations of atmospheric pollutants SO₂, PM₁₀, Pb aerosols, NO₂ and H₂S) and the morbidity of the population based on the number of people who visited their GPs in a relation with a chronic health problem or emergency condition and the number of hospitalisations in two age groups (newborn to 17 years olds and 18 and older) as well as for the entire population in the period 2009-2013, making it possible to predict morbidity levels.

RESULTS: The expected morbidity level predictions based on the number of people who visited their GPs in Municipality are lower, while hospitalisation level predictions are higher. This model has been created and tested and is applicable in all residential areas.

CONCLUSIONS: A new, very sensitive, mathematical model has been created and tested (average margin of error from 0.61% to 2.59%) and is applicable in all residential areas.

Introduction

Stara Zagora Municipality is situated in the central part of South Bulgaria, in the Upper Thracian Plain, on the south hills of Sredna Sarnena Gora mountain (N 42° 25', E 25° 37'). It covers an area of 1019.1 square km and is located at 240 m above sea level. It comprises 52 towns and villages with a total of 156,662 residents, 137,834 of whom live in the city of Stara Zagora. Stara Zagora Municipality is amongst the most industrially developed municipalities in Bulgaria, which is the main cause of air, water and soil pollution. One of the key threats to public health associated with the environment in Stara Zagora is

atmospheric air pollution. Its protection is governed by the Clean Air Act [1] as well as by its associated regulations – Regulation No. 14/1997 [2]; Regulation No. 11/2007 [3]; Regulation No. 12/2010 [4], where all Bulgarian rules and regulations are harmonised with those of the EU (Directive 2008/50/EC). Atmospheric air quality monitoring and control are carried out throughout the year by the Regional Inspectorate of Environment and Water Directorate.

In the past 20 years, the atmospheric air in Stara Zagora Municipality has been amongst the most polluted in the country [5] [6] [7]. The atmospheric air quality in the municipality is contingent upon the operation of numerous local (industrial enterprises, domestic heating, heavy traffic) and regional (the

three thermal power stations of the Maritza East power complex situated 40 km south of Stara Zagora and Zmeevo, a military testing ground 7 km to the southwest) emission sources. These sources mainly generate sulphur dioxide (SO₂), breathable PM₁₀, nitrogen dioxide (NO₂), lead aerosols and hydrogen sulphide (H₂S). Two trans-European corridors go through the territory of the municipality. These characteristics coupled with weather conditions and the topography of the area facilitate the migration of pollutants and affect the morbidity of the population. An ecological task of top priority is to prevent atmospheric air pollution from the major pollutants (PM₁₀, sulphur and nitrogen oxides, ozone), given their role as a risk factor in the aetiology, pathogenesis and spreading of some diseases (cardiovascular diseases, respiratory diseases etc.)

The purpose of this study is to analyse air pollution and its impact on the morbidity amongst the population of Stara Zagora Municipality in the period 2009 - 2013, as well as to create a mathematical model of predicting morbidity levels.

Subjects and Methods

On the territory of Stara Zagora Municipality, sample-taking and analyses of the atmospheric pollutants (sulphur dioxide, PM₁₀, nitrogen oxides, lead aerosols and hydrogen sulphide) are conducted in an automatic measuring station, two differential optical absorption spectroscopic systems and one unit for the manual taking of samples. Standards in compliance with European law are used in the analysis of the pollutants. Information about the morbidity based on the number of people who visited their GPs in a relationship with a chronic health problem or emergency condition and the number of hospitalisations has been obtained from the Regional Health Inspectorate – form 365, 1A, appendices 5 and 6. The study covers the two age groups (from 0 to 7-year-olds, 18 and older) and the entire population of Stara Zagora Municipality.

The influence of atmospheric pollution on the morbidity of the population in Stara Zagora Municipality in the period 2009-2013 is based on disease class according to the International Classification of Diseases, revision X.

The correlation between the examined atmospheric pollutants and morbidity can be calculated using the following formula:

n

$$Y_i = e^{ei} K_0 \prod X_{ij}^{K_j}$$

j=1

Where:

Y_i - the dependant variable (result = the number of predicted diseases);

i - the year;

n – number of factors;

X_{ij} - the value of factor j for the period i;

K_j – powers (coefficients) – the constants of each dependence;

K_0 - the module (the equaliser of the dimensions of each dependence);

ε_i - statistical error.

Module K_0 and the powers K_j are obtained using an integral exponential function by applying the least squares method used by Frenkely and described in Multifactor correlation model for labour productivity [8].

In this particular case, the factors (pollutants) are $X_{i,1}$ -SO₂, $X_{i,2}$ -PM₁₀, $X_{i,3}$ -NO₂, $X_{i,4}$ -Pb aerosols). The magnitude of each factor (X_i) is calculated in $\mu\text{g}/\text{m}^3$.

Results

The analysis of systemically controlled pollutants (SO₂, PM₁₀, NO₂, H₂S and lead aerosols) by months and years for the period 2009-2013. Showed that the mean monthly concentrations of SO₂, NO₂, H₂S and lead aerosols did not exceed the permissible levels. The only observed pollutant in which there are cases of exceeding mean monthly concentrations relative to the threshold value is PM₁₀. At only one of the measurement points, all PM₁₀ average annual PM₁₀ values for 2009, 2010 and 2011 are above the limit value of 40 $\mu\text{g}/\text{m}^3$. Contributing to this is probably the lively traffic on the boulevard where the points are located. About the annual average concentrations, the results for all atmospheric pollutants indicate that they are within the boundaries of the norm and safety for human health. The complex assessment of atmospheric pollution in the municipality of Stara Zagora shows a slight fluctuation of the aggregate indices by years, without any sudden changes, and confirms the tendency to reduce the systemically observed pollutants: sulfur dioxide, PM₁₀ nitrogen oxides, lead aerosols and hydrogen sulphide for the period 2009-2013.

As regards the morbidity based on the number of people who visited their GPs in a relationship with a chronic health problem or emergency condition, the greatest share for the entire population take the cardiovascular diseases (22.99%). They are more frequent (27.77%) in the age group of 18 and older, or more as compared to the age group 0-17 (0.97%). Second are respiratory diseases

(17.34%) for the entire population. In age group 0-17, the relative share is rather high, 47.20%. Third, come to the diseases of the genitourinary system (9.76%).

As regards incidence of diseases that require hospitalisation, the leading diseases with almost the same share are the diseases of the digestive system (11.90%) and cardiovascular diseases (11.85), followed by respiratory diseases (10.86%) and genitourinary systems (8.88%). In age group 0-17, the greatest share is the share of the respiratory diseases (36.93%). Most common in this disease class is pneumonia (82.44%). Children most frequently suffer from infectious diseases where disturbed immune balance is observed. Cardiovascular diseases are most common amongst adults (14.26%).

During the examined period, there is a tendency in Stara Zagora Municipality towards decreasing the incidence of diseases based on the number of people who visited their GPs in a relationship with a chronic health problem or emergency condition and an increase based on hospitalisation as compared to the base year (2009).

The correlation between atmospheric pollution and the morbidity of the population as well as tracking the influence of each pollutant on morbidity is determined using the values of the mathematical expression X_{ij}^{Kj} based on the mathematical model used. The data analysis after the computations shows that PM_{10} has the greatest influence on the morbidity of the population in Stara Zagora Municipality based on the number of people who visited their GPs in a relationship with a chronic health problem or emergency condition and the number of hospitalisations. The correlation is particularly strong in the age group 0-17 by the number of people who visited their GPs in a relationship with a chronic health problem or emergency condition ($r = 0.762, p < 0.05$).

Sulphur dioxide exhibits a positive correlation with morbidity based on the number of hospitalisations in the age group 18 and older at the beginning of the studied period (2009-2013) ($r = 0.430, p < 0.05$), and at the end of the period it is negative. Lead aerosols too have a pronounced influence on morbidity based on the number of hospitalisations amongst the population aged 18 and older in Stara Zagora Municipality ($r = 0.501, p < 0.05$).

We found the additive action of some pollutants as SO_2+PM_{10} on malignant diseases for the whole population by hospitalisation [9] [10] [11] [12] [13] and cardiovascular diseases for the whole population is affected by the additive effect of $PM_{10}+Pb$ aerosols [14] [15] [16].

After applying the Least squares method on the morbidity based on the number of hospitalisations and on the number of people who visited their GPs in a relationship with a chronic health problem or emergency condition amongst the entire population in

Stara Zagora Municipality, the following coefficients are obtained as an independent variable:

$$K_0 e^{19.7723} \quad K_1 - 0.1039 \quad K_2 - 1.8949 \quad K_3 - 1.5106 \quad K_4 - 0.2254$$

The following formula will be used to calculate the predicted future morbidity count:

$$e^{19.7723} X_{i,1}^{Y_{prognosis}} X_{i,2}^{0.1039} X_{i,3}^{1.8949} X_{i,4}^{1.5106} X_{i,4}^{0.2354} =$$

The result calculated based on the formula above makes it possible to forecast morbidity with a significant precision-the average margin of error for the studied period ranges from 0.61 to 2.59%. The expected morbidity level predictions based on the number of people who visited their GPs in a relationship with a chronic health problem or emergency condition in Stara Zagora Municipality are lower, while hospitalisation level predictions are higher as compared to the numbers obtained during the period of the study (Table 1).

Table 1: Registered and projected morbidity of the population in Stara Zagora Municipality by the number of people who visited their GPs in a relationship with a chronic health problem or emergency condition and the number of hospitalisations for the period 2009-2013

Year	Number of registered diseases		Number of projected diseases		Error for the year %
	Diseases by the number of people who visited their GPs in a relationship with a chronic health problem or emergency condition		Diseases by the number of hospitalisations		
2009	333425	105396	34038	105396	0.68
2010	339482	196226	37277	196226	0.42
2011	402741	108831	37194	108831	0.72
2012	321726	141328	32019	141328	0.56
2013	321968	104115	45777	104115	0.67
The average margin of error for the period = 0.61%					
Year	Number of registered diseases		Number of projected diseases		Error for the year %
	Diseases by the number of people who visited their GPs in a relationship with a chronic health problem or emergency condition		Diseases by the number of hospitalisations		
2009	333425	105396	34038	105396	2.09
2010	339482	196226	37277	196226	4.26
2011	402741	108831	37194	108831	1.92
2012	321726	141328	32019	141328	3.41
2013	321968	104115	45777	104115	1.27
The average margin of error for the period = 2.59%					

Discussion

The majority of cardiovascular diseases can be explained by the fact that people older than 65 are more vulnerable as regards hypertonic disease, cerebrovascular diseases and ischemic heart disease. These diseases, in particular, indirectly associated with air pollution, seasons and climate factors, increase the possibility of the occurrence of thrombosis [17] [18] [19]. The respiratory system is affected to no lesser extent, and data shows that it is the most sensitive to atmospheric pollutants and children are the most sensitive group [20] [21] [22]. This is correlated to the immature immunity, high vulnerability and the body's predisposition to different respiratory diseases, especially in the early childhood years. Frequent pneumonia in hospitalised children

are influenced by meteorological factors, which is confirmed by the studies of some authors [23] [24]. It is established that atmospheric pollutants have a suppressive effect on the production of immunoglobulin for this age group [25] [26] [27].

Similar to our observations of the effect of nitrogen dioxide on the respiratory system, described Nickmilder M. et al., 2007 which establish the inflammatory changes in the airways, but they bind them to the low-level contamination with nitrogen oxides and ozone [28].

Numerous studies of this age group show a positive relationship even between low concentrations of lead aerosols and damage to hemopoiesis, nervous tissue, hearing, changes in neurobehavioral functions, and intellectual [29] [30] [31] [32]. Reports of blocking the bioactivity of porphyrin synthesis enzymes, impaired haem- synthesis and the development of anemia by the action of lead is found in research by de Carvalho RP et al., 2006; Zimmermann M. B. et al., 2006; Muwakkit S. et al., 2008; Rondó P. K. et al., 2011, Park S. et al., 2014 [33] [34] [35] [36] [37]. A lot of studies show a relationship between atmospheric pollution and malignant diseases. The analyzing of the data in morbidity by the number of people who visited their GPs in a relation with a chronic health problem or emergency condition of the groups from 0 to 17 years, 18 and more and the whole population shows high correlation dependence between sulphur dioxide and the increasing of bladder and lung cancer leading to high mortality [38] [39] [40] [41]. For the influence of PM₁₀ on all diseases, report Raaschou-Nielsen O. et al., 2013; Ponomareva L. A et al. V.V. Inogamova, 2013; Jevtić M. et al., 2014 [42] [43] [44]. Some factors like quality of water, food and feeding can affect the occurrence of diseases of the genitourinary system.

Using the mathematical model formula, it is possible to calculate the predicted value of the number of diseases based on all average annual concentrations of the pollutants for the years in question. The formula is a multiplication product because all pollutants together affect the morbidity and since the concentrations of each factor (pollutant) change over time, an integral exponential function is included too. The calculation of the average development rate for the different pollutants makes it possible to predict the level of atmospheric pollution for each coming year and, hence, the morbidity for the same period. The decrease in atmospheric pollution for the studied period is associated with the commissioning and effective operation of desulphurisation in Maritza East power complex, the efforts directed toward maintaining the road infrastructure, the gasification of Stara Zagora City, reduced use of wood and coal for heating in the municipality, the use of more environmentally-friendly fuels. Determining the projected morbidity levels confirms the tendencies in the dynamics and structure

of morbidity based on the number of people who visited their GPs in a relationship with a chronic health problem or emergency condition and the number of hospitalisations amongst the population in Stara Zagora Municipality for the period 2009-2013.

In conclusion, the main methods of medical, mathematical statistics (regression, correlation analysis etc.) do not make it possible to cover fully and precisely the actual correlations between the different factors and the result. In fact, these connections are non-linear which is why it is necessary to use a multi-factor correlation model with non-linear dependences. The functions used in the current study constitute a connection between two factors (atmospheric pollutant and morbidity) and exclude the interaction of all factors. The proposed dependence provides a theoretical model of the interaction, which is quite close to the real one. Determining the dependences makes it possible to realise and control the fluctuations of the result through the impact of one factor or another. A disadvantage of the model can be assumed to be the implicit nature of the functional interdependencies. In reality, this does not have a considerable effect on the practical application of the model in every residential area.

Using the created mathematical model and the atmospheric pollution data, it is possible to forecast the morbidity in every residential area.

Acknowledgements

I hereby thank Valentin Popov - Teaching Fellow, St Andrews University, Fife, Scotland for his cooperation and the great help he offered me while preparing the mathematical model.

References

1. Clean Air Act. Last amendment: Bulgarian State Gazette № 38, May 18, 2012.
2. Ordinance № 14 on standards for permissible concentrations of harmful substances in the air of the settlements. Bulgarian State Gazette № 88, October 03, 1997.
3. Ordinance № 11 on the norms of arsenic, cadmium, nickel and polycyclic hydrocarbons in air. Bulgarian State Gazette № 42, May 29, 2007.
4. Ordinance № 12 on the norms of sulphur dioxide, nitrogen dioxide, particulate matter, lead, benzene, carbon monoxide and ozone in the atmosphere. Bulgarian State Gazette № 58, July 30, 2010.
5. Takuchev N. Analysis of Stara Zagora Air Pollution in the Period 2002-2004, Type of Pollutants, Regime, Climatic and Relief Prerequisites, Sources. Journal of Ecology and Future. 2008; 7: 3-

- 12.
6. Petkov G, Kostadinova G. Air pollution and its impacts on ecosystems. *Agricultural science and technology*. 2008; 54:177-222.
7. Kostadinova G, Petkov G, Georgieva N, Dospatliev L, Dermendjieva D. Assessment of Air Pollution in Stara Zagora Region, Bulgaria. *Journal of Balkan Ecology*. 2011; 14:341-355.
8. Frenkely AA. Multifactor correlation model for labor productivity. 1965; 143-145.
9. Nyberg F, Gustavsson P, Järup L, Bellander T, Berglund N, Jakobsson R, Pershagen G. Urban Air Pollution and Lung Cancer in Stockholm. *Epidemiology*. 2000; 11(5):487-495. <https://doi.org/10.1097/00001648-200009000-00002> PMID:10955399
10. Benedetti M, Lavarone I, Comba P. Cancer Risk Associated with Residential Proximity to Industrial Sites. *Archives of Environmental Health: An International Journal*. 2001; 56(4):342-349. <https://doi.org/10.1080/00039890109604466> PMID:11572278
11. Scott AV, Wang B, Xu Z, Schlatter Y, Wang L, Xu X. Particulate matter, sulfur dioxide, and daily mortality in Chongqing, China. *Environ Health Perspect*. 2003; 111(4):562-567.
12. Krewski D, Burnett R, Jerrett M, Pope CA, Rainham D, Calle E, Thurston G, Thun M. Mortality and Long-Term Exposure to Ambient Air Pollution: Ongoing Analyses Based on the American Cancer Society Cohort. *Journal of Toxicology and Environmental Health, Part A*. 2005; 68(13-14):1093-1109.
13. Beelen R, Hoek G, van den Brandt P, Goldbohm R, Fischer P, Schouten L, Armstrong B, Brunekreef B. Long-Term Exposure to Traffic-Related Air Pollution and Lung Cancer Risk. *Epidemiology*. 2008; 19(5):702-720. <https://doi.org/10.1097/EDE.0b013e318181b3ca> PMID:18633326
14. Banza CLN, Nawrot TS, Haufroid V, Decrée S, De Putter T, Smolders E et al. High human exposure to cobalt and other metals in Katanga, a mining area of the Democratic Republic of Congo (Article). *Environmental Research*. 2009; 109(6):745-752. <https://doi.org/10.1016/j.envres.2009.04.012> PMID:19486963
15. Chen L, Xu Z, Liu M, Huang Y, Fan R, Su Y, Hu G, Peng X. Lead exposure assessment from study near a lead-acid battery factory in China. *Science of The Total Environment*. 2012; 429:191-198. <https://doi.org/10.1016/j.scitotenv.2012.04.015> PMID:22578522
16. Mielke HW, Gonzales C, Powell E, Mielke PW. Evolving from reactive to proactive medicine: Community lead (Pb) and clinical disparities in Pre- and post-Katrina new Orleans. *International Journal of Environmental Research and Public Health*. 2014; 11(7):7482-7491. <https://doi.org/10.3390/ijerph110707482> PMID:25050655 PMCid:PMC4113889
17. Dales RE, Cakmak S, Vidal CB. Air pollution and hospitalization for venous thromboembolic disease in Chile. *J Thromb Haemost*. 2010; 8:669-674. <https://doi.org/10.1111/j.1538-7836.2010.03760.x> PMID:20088925
18. Martinelli N, Girelli D, Ciqolini D, Sandri M, Ricci G, Rocca G, et al. Access rate emergency department for venous thromboembolism in relationship with coarse and fine particulate matter air pollution. *PLoS One*. 2012; 7:e34831.
19. Damjanović Z, Jovanović M, Stojanović M. Correlation between the climatic factors and the pathogenesis of deep vein thrombosis. *Hippokratia*. 2013; 17:203-206.
20. Brent R, Weitzman M (supplement Editor). The vulnerability, sensitivity and resiliency of the developing embryo, infant, child and adolescent to the effects of environmental, chemical, drugs and physical agents as compared to the adult. *Pediatrics*. 2004; 113:932-1172.
21. Bateson TF, Schwartz J. Children's Response to Air Pollutants. *Journal of Toxicology and Environmental Health, Part A*. 2007; 71:238-243. <https://doi.org/10.1080/15287390701598234> PMID:18097949
22. Salvi S. Health effects of ambient air pollution in children. *Paediatric Respiratory Reviews*. 2007; 8:275-280. <https://doi.org/10.1016/j.prrv.2007.08.008> PMID:18005894
23. Almasri M, Diza E, Papa A, Eboriadou M, Souliou E. *Mycoplasma pneumoniae* respiratory tract infections among Greek children. *Hippokratia*. 2011; 15:147-152.
24. Chen ZR, Yan YD, Wang YQ, Zhu H, Shao XJ, Xu J, Ji W. Epidemiology of community-acquired *Mycoplasma Pneumoniae* respiratory tract infections among hospitalized Chinese children, including relationships with meteorological factors. *Hippokratia*. 2013; 17:20-26.
25. McDermott MR., Befus AD, Bienenstock J. The structural basis for immunity in the respiratory tract. *Int Rev Exp Pathol*. 1982; 23:47-112.
26. Milligan MS, Warren JS, Smith CW, Anderson DC, Yen CG, Rudolph AR, Ward PA. Lung injury after deposition of Ig A immune complexes: requirements for CD18 and L-arginine. *J Immunol*. 1992; 148:3086-3092.
27. Lukacs NW, Ward PA. Inflammatory mediators, cytokines and adhesion molecules in pulmonary inflammation and injury. *Adv Immunol*. 1996; 82:257-304.
28. Nickmilder M, de Burbure C, Sylviane C, Xavier D, Alfred B, Alain D. Increase of Exhaled Nitric Oxide in Children Exposed to Low Levels of Ambient Ozone. *Journal of Toxicology and Environmental Health*. 2007; 70(3-4):270-274.
29. Alvarenga KF, Morata TC, Lopes AC, Feniman MR, LJ Corteletti. Brainstem auditory evoked potentials in children with lead exposure. *Brazilian Journal of Otorhinolaryngology*. 2014; 2014.
30. Chen K, Huang L, Yan B, Li H, Sun H, Bi J. Effect of Lead Pollution Control on Environmental and Childhood Blood Lead Level in Nantong, China: An Interventional Study. *Environmental Science & Technology*. 2014; 48(21):12930-12936. <https://doi.org/10.1021/es502994j> PMID:25294690
31. Jeong KS, Park H, Ha E, Hong YC, Ha M, Park H et al. Evidence that cognitive deficit in children is associated not only with iron deficiency, but also with blood lead concentration: A preliminary study. *Journal of Trace Elements in Medicine and Biology*. 2014.
32. Kicinski M, Vrijens J, Vermier G, Hond ED, Schoeters G, Nelen V et al. Neurobehavioral function and low-level metal exposure in adolescents. *International Journal of Hygiene and Environmental Health*. 2014.
33. De Carvalho RP, Carvalho MFH, Souzaet MC, Moraes F. Lead, hemoglobin, zinc protoporphyrin and ferritin concentration in children. *Revista de Saude Publica*. 2006; 40(1):71-76. <https://doi.org/10.1590/S0034-89102006000100012>
34. Zimmermann MB, Muthayya S, Moretti D, Kurpadet A, Hurrell ARF. Iron fortification reduces blood lead levels in children in Bangalore, India. *Pediatrics*. 2006; 117(6):2014-2021. <https://doi.org/10.1542/peds.2005-2440> PMID:16740843
35. Muwakkit S, Nuwayhid I, Nabulsi M, Hajj RAI, Khoury R, Mikati M et al. Iron deficiency in young Lebanese children: Association with elevated blood lead levels. *Journal of Pediatric Hematology/Oncology*. 2008; 30(5):382-386. <https://doi.org/10.1097/MPH.0b013e318165b283> PMID:18458574
36. Rondó PK, Conde A, Souza MC, Sakuma A. Iron deficiency anaemia and blood lead concentrations in Brazilian children. *Trans R Soc Trop Med Hyg*. 2011; 105:525-530. <https://doi.org/10.1016/j.trstmh.2011.05.012> PMID:21742361
37. Park SK, Sim CS, Lee H, Et H, Kim Y. Effects of iron therapy on blood lead concentrations in infants. *Journal of Trace Elements in Medicine and Biology*. 2014; 28(1):56-59. <https://doi.org/10.1016/j.jtemb.2013.11.003> PMID:24315962
38. Englander V, Söberg A, Hagmar L, Attewell R, Schütz A, Möller T, Skerfving S. Mortality and cancer morbidity in workers exposed to sulphur dioxide in a sulphuric acid plant. *Occupational Environmental Health*. 1988; 61(3):157-162. <https://doi.org/10.1007/BF00381012>

39. Lee JW, Teschke K, Kauppinen T, Andersen A, Jäppinen P, Szadkowska-Stanczyk I et al. Mortality from lung cancer in workers exposed to sulfur dioxide in the pulp and paper industry. *Environ Health Perspect*. 2002; 110(10):991-995. <https://doi.org/10.1289/ehp.02110991> PMID:12361923
PMCID:PMC1241024
40. Nafstad P, Håheim LL, Oftedal B, Gram F, Holme I, Hjermann I et al. Lung cancer and air pollution: a 27 year follow up of 16,209 Norwegian men. *Thorax*. 2003; 58:1071-1076. <https://doi.org/10.1136/thorax.58.12.1071> PMID:14645978
PMCID:PMC1746548
41. Gurjar BR, Jain A, Sharma A, Gupta P, Nagpure AS, Lelieveld J. Human health risks in megacities due to air pollution. *Atmospheric Environment*. 2010; 44(36):4606–4613. <https://doi.org/10.1016/j.atmosenv.2010.08.011>
42. Raaschou-Nielsen O, Andersen ZJ, Beelen R, Samoli E, Stafoggia M, Weinmayr G et al. Air pollution and lung cancer incidence in 17 European cohorts: prospective analyses from the European Study of Cohorts for Air Pollution Effects (ESCAPE). *The Lancet Oncology*. 2013;14(9):813-822. [https://doi.org/10.1016/S1470-2045\(13\)70279-1](https://doi.org/10.1016/S1470-2045(13)70279-1)
43. LA Ponomareva VV, Inogamova. Atmospheric air pollution in Tashkent and its importance for the occurrence of the nephrology diseases. *Uzbekiston Tibbiy Zhurnali*. 2013; 2:71-73.
44. Jevtić M, Dragić N, Bijelović S, Popović M. Cardiovascular diseases and air pollution in Novi Sad, Serbia. *International journal of Occupational Medicine and Environmental Health*. 2014; 27(2):153-164. <https://doi.org/10.2478/s13382-014-0239-y> PMID:24549995