

# Transverse Skeletal Effects of Rapid Maxillary Expansion in Pre and Post Pubertal Subjects: A Systematic Review

Nouran Fouad Seif-Eldin<sup>\*</sup>, Sherif Aly Elkordy, Mona Salah Fayed, Amr Ragab Elbeialy, Faten Hussein Eid

*Department of Orthodontics, Faculty of Dentistry, Cairo University, Cairo, Egypt*

## Abstract

**Citation:** Seif-Eldin NF, Elkordy SA, Fayed MS, Elbeialy AR, Eid FH. Transverse Skeletal Effects of Rapid Maxillary Expansion in Pre and Post Pubertal Subjects: A Systematic Review. Open Access Maced J Med Sci. 2019 Feb 15; 7(3):467-477. <https://doi.org/10.3889/oamjms.2019.080>

**Keywords:** Rapid maxillary expansion; Rapid palatal expansion; Pre-pubertal; Post-pubertal; Skeletal effects

**\*Correspondence:** Nouran Fouad Seif-Eldin, Department of Orthodontics, Faculty of Dentistry, Cairo University, Cairo, Egypt. E-mail: [nouran.fouad@dentistry.cu.edu.eg](mailto:nouran.fouad@dentistry.cu.edu.eg)

**Received:** 17-Oct-2018; **Revised:** 26-Dec-2018; **Accepted:** 31-Dec-2018; **Online first:** 08-Feb-2019

**Copyright:** © 2019 Nouran Fouad Seif-Eldin, Sherif Aly Elkordy, Mona Salah Fayed, Amr Ragab Elbeialy, Faten Hussein Eid. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

**Funding:** This research did not receive any financial support

**Competing Interests:** The authors have declared that no competing interests exist

**OBJECTIVE:** The aim of this systematic review was to assess the transverse skeletal effects of rapid maxillary expansion (RME) in pre and post-pubertal subjects.

**MATERIAL AND METHODS:** Five databases were searched till May 2018; Pubmed, Cochrane, Scopus, Lilacs and Web of science in addition to the manual search of other sources. There were no language restrictions. Methodological Index for Non-Randomized Studies MINORS was used to assess the quality and risk of bias of the trials included.

**RESULTS:** Six studies were finally included in the qualitative analysis. A meta-analysis wasn't performed due to the heterogeneity of methodologies and outcomes. All of the included studies showed drawbacks in their structure yielding weak evidence. On the short term, RME caused an increase in the maxillary and lateral-nasal widths in pre-pubertal subjects by 3.4 mm and 3.3 mm, and by 2.8 and 2.2 mm respectively in post-pubertal subjects. Although statistically insignificant, the maxillary width increase was more than that of the post-pubertal subjects by 0.6 mm. Over the long term, expansion produced permanent increases in the transverse dimensions of both the dento-alveolar and skeletal components of the maxilla and circum-maxillary structures in pre-pubertal subjects. The post-pubertal subjects presented with a statistically significant increase only in the later-nasal width by 1.3 mm than the untreated controls with no permanent increase in the skeletal maxillary width.

**CONCLUSION:** The literature is very deficient regarding the use of skeletal age as a reference in the treatment of skeletal crossbites using RME. Only weak evidence exists supporting the increased maxillary and lateral-nasal widths after tooth-tissue borne RME in pre-pubertal subjects, with these effects being less in the post-pubertal ones.

## Introduction

Transverse maxillary deficiency is a common finding among populations, mostly presented with a unilateral or bilateral posterior crossbite. The posterior crossbite is reported to be the most prevalent type of malocclusion occurring between 8% and 22% [1]. Whether skeletal or dental, crossbites should be treated once diagnosed, as it is believed that skeletal crossbites affect TMJ functions, chewing patterns, breathing habits and tongue posture. The main goal of skeletal crossbite correction resulting from a maxillary

deficiency is achieving transverse skeletal expansion of the maxilla with the least dental effects allowing optimum coordination of the maxillary and mandibular dental arches.

In 1975, Melsen [2] divided the morphological development of the median palatine suture (MPS) into 3 stages, proposing that opening of the suture through maxillary expansion is best done before the age of 15 years as, after that age, the growth of the suture was observed to be ceased. This conclusion was also highlighted in 1977 when Person and Thilander [3] demonstrated in a histological study that the degree of suture obliteration increases from the juvenile period

to adulthood, yet complete suture closure was rarely found until the third decade of life. Accordingly, practitioners for decades have been using the chronological age to decide the treatment of choice for the transverse maxillary skeletal deficiency: slow or rapid maxillary expansion (SME/RME) for that who are under 15 years of age and surgically assisted rapid palatal expansion (SARPE) for those who are above that age [4].

Since categorising subjects by chronological age has many limitations compared to measuring the developmental status of individuals about specific stages of skeletal maturation, Fishman [5] in 1994 correlated between his famous eleven skeletal maturational stages and the percentage of the MPS closure using occlusal radiographs. Surprisingly, the study found that at the maturational age of skeletal maturational index (SMI) 11, only 50% of the MPS was approximated. Since then, authors [6], [7], [8], [9], [10] started to investigate the validity of the chronological age in decision making when it comes to expansion, and a paradigm shift has occurred with an ample of researches [11], [12], [13], [14], [15], [16], [17], [18] aiming at suture separation in young adults through non-surgical RME. These studies tried to make use of the partially fused MPS and performed non-surgical RME in postpubertal subjects. A debate has aroused since then around the success of non-surgical RME in adults when compared to RME in pre-pubertal subjects.

To our knowledge, a well-designed systematic review evaluating the effects of RME in pre- and post-pubertal subjects; according to the skeletal age, hasn't been conducted yet. Such review would help to deliver the status of evidence to solve this debate and provide a guide to future clinical practice, aiming for the optimum treatment option customised to each patient according to his/her skeletal age.

Hence, the aim of the current systematic review was to provide the answer to the following question: in Pre- and Postpubertal subjects with skeletal maxillary constriction, does rapid maxillary expansion (RME) cause widening in the mid-palatal suture (MPS) and/or an increase/change in any skeletal transverse maxillary measurement?

## Material and Methods


This systematic review followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement and the Cochrane handbook for systematic reviews of intervention [19]. The protocol was registered at the Evidence-Based Center, Faculty of Dentistry, Cairo University (online registration was not performed).

Information sources, search strategy, and study selection Pubmed, Cochrane Central Register of Controlled Trials (CENTRAL, The Cochrane Library), Lilacs, Web of Science and Scopus were electronically searched till May 2018 with no language restrictions. Details of the PubMed search are shown in table [1]. Manual Search was performed in the following journals *American Journal of Orthodontics & Dentofacial Orthopedics*, *The Angle Orthodontist*, *European Journal of Orthodontics*, *Progress in Orthodontics*, *Orthodontics & Craniofacial Research* and *Seminars in Orthodontics* (Figure 1).

**Table 1: PubMed search strategy**

|                |   |
|----------------|---|
| #1             | Crossbite OR crossbite OR skeletal crossbite OR skeletal cross bite OR posterior crossbite OR posterior crossbite OR bilateral crossbite OR unilateral crossbite OR constrict* maxilla OR narrow maxilla  |
| #2             | Rapid palatal expan* OR rapid maxillary expan* OR palatal expan* OR maxillary expan* OR bone borne maxillary expan* OR tooth borne maxillary expan* OR miniscrews assisted rapid maxillary expan* OR miniscrews assisted rapid palatal expan* OR maxilla* wide* OR Haas OR hyrax  |
| #3 (#1 AND #2) | Cross bite OR crossbite OR skeletal crossbite OR skeletal cross bite OR posterior crossbite OR posterior cross bite OR bilateral crossbite OR unilateral crossbite OR constrict* maxilla OR narrow maxilla AND Rapid palatal expan* OR rapid maxillary expan* OR palatal expan* OR maxillary expan* OR bone borne maxillary expan* OR tooth borne maxillary expan* OR miniscrews assisted rapid maxillary expan* OR miniscrews assisted rapid palatal expan* OR maxilla* wide* OR Haas OR hyrax |

At the same time, grey literature and unpublished data were electronically searched at the *Central Library of Cairo University*, *Egyptian Universities Libraries*, *Clinical Trials.gov* and *Edinburgh Research Archive* using the following keywords: "posterior crossbite", "hyrax", "rapid maxillary expansion", "rapid palatal expansion" an "maxillary transverse deficiency".



**Figure 1: PRISMA diagram of article retrieval**

Two investigators (N.S) and (S.K) independently performed the search and any points of disagreements or conflicts were finalized through discussion with the third author (M.F), and the final inclusion of the study was after the agreement.

### Eligibility Criteria

The search was conducted upon certain inclusion and exclusion criteria shown in Table 2. An article was considered eligible if it included at least two treatment groups; a study group with rapid palatal expander and an untreated control group. After removal of the internal and external duplicates, the articles obtained from all the databases and manual search were screened by title, abstract and full texts. The full text was obtained when titles and abstracts were insufficient to decide.

### Data Items and Collection

The investigators used the identified articles to develop data extraction sheets to, and data were extracted by each investigator independently. The following data were collected: study design, demographic data including participants' maturational status, number, gender and number of groups, an appliance for RME, its anchorage and design, the frequency of activation, retention protocol and timing and method of records collection.

### Quality Assessment of Individual Studies

Evaluation of the methodological quality was intended to be performed using the Cochrane Risk of Bias Tool [19] to analyse the risk of bias in RCT's. An overall assessment of the risk of bias (high, unclear, low) was made according to seven criteria for assessment: random sequence generation, allocation concealment, blinding of participants and personnel, blinding of assessors, incomplete outcome data, selective reporting of outcomes, and other potential sources of bias.

MINORS [20] (Methodological Index for Non-Randomized Studies) was the tool of choice for non-randomized controlled and retrospective studies due to its simplicity and reliability. For each article, 12 points were to be assessed: a clearly stated aim, inclusion of consecutive patients, prospective collection of data, endpoints appropriate to the aim of the study, unbiased assessment of the study endpoints, follow up period appropriate to the aim of the study, loss to follow up less than 5%, prospective calculation of the study size, adequate control group, contemporary groups, baseline equivalence of groups and adequate statistical analyses.

The items are scored 0 if not reported, 1 if reported but inadequate or 2 if reported and adequate. The global ideal score is 16 for non-comparative

studies and 24 for comparative studies.

**Table 2: Inclusion and exclusion criteria of the studies**

|                 | Inclusion Criteria   | Exclusion Criteria  |
|-----------------|--|---|
| 1. Population   | - Any age group concerning the maturational status.<br>- Both sexes<br>- Transverse Skeletal maxillary constriction  | - Not referring to the maturational status index<br>- Dental & functional cross bites<br>- Skeletal mandibular asymmetry<br>- Cleft lip and palate<br>- Syndromic patients              |
| 2. Intervention | Any rapid palatal expander (tooth-borne or bone-borne)   | - SARPE<br>- Corticotomy or any other surgical intervention/ Distractors<br>- Receiving fixed orthodontic treatment or any conjunctive treatment as headgear, face mask, chin cup...etc |
| 3. Comparator   | Presence of untreated control groups   | If a control group is absent/ with normal occlusion/ receiving any orthodontic treatment  |
| 4. Outcomes     | The primary outcome of the search was to detect any change in maxillary transverse skeletal measurements following RME. Secondary outcomes included detecting changes in:<br>1. Nasal airway<br>2. Maxillary and mandibular dental arches<br>3. Facial and nasal soft tissues. | Any irrelevant outcome  |
| 5. Study Design | 1. Randomised clinical trials (RCT's)<br>2. Quasi-randomised clinical trials.<br>3. Prospective controlled clinical trials (CCT's)<br>4. Retrospective controlled trials.  | - Abstracts<br>- Comments<br>- Case Reports<br>- Narrative Reviews<br>- Case series<br>- Expert opinion<br>- Systematic reviews and meta-analyses<br>- In vitro studies.                |

### Summary Measures and Approach to Synthesis

The final studies included in this review measured different aspects of the effects RME that made the collected data not combinable. Thus, a meta-analysis was not performed due to the great clinical heterogeneity and variation among the measured outcomes, and only a qualitative analysis was performed.

## Results

**Study Selection and Characteristics:** Two thousand and sixty-four articles were obtained from the electronic search of the five databases and eight articles obtained from the manual search. By applying the inclusion and exclusion criteria, 67 articles were found to be eligible and were read as full text.

Most of the excluded studies [11], [21], [22], [23], [24], [25], [26], [27], [28], [29], [30], [31], [32], [33], [34], [35], [36], [37], [38], [39], [40], [41], [42], [43], [44], [45], [46], [47], [48], [49], [50], [51], [52], [53], [54], [55], [56], [57], [58], [59], [60], [61], [62], [63] divided the samples into treated and untreated control groups according to their chronological age and not according to their skeletal age. Seventeen studies [11], [21], [22], [26], [27], [36], [37], [38], [45], [46], [47], [51], [52], [54], [60], [63], [64], [65] had an untreated control group with normal occlusion or with other types of malocclusions other than posterior

crossbites. Other reasons for exclusion included, subjecting the control groups to treatment [11], [32], [44], case reports [66], [67], [68], [69], using interventions other than RME [50], [51], [53], [54], [56], [64], [65], [70] or absence of a control group [31], [71], [72]. Articles which couldn't be retrieved also were excluded [73], [74], [75], [76], [77], [78]. Finally, six trials [12], [79], [80], [81], [82], [83] were included in the qualitative analysis; four prospective controlled studies [79], [80], [82], [83], one retrospective study<sup>(12)</sup> and one prospective with a retrospective control group [81] that evaluated the long term changes of RME. Two researchers studied the effect of RME in pre and post pubertal subjects [12], [79] while the remaining four articles evaluated the effects in pre-pubertal subjects only [80], [81], [82], [83].

### Description of intervention

In the previously mentioned studies, bonded acrylic splints, banded Haas, bonded and banded hyrax was used for expansion without using any bone borne appliances. The studies reported that expansion was found sufficient when observing the palatal cusps of the upper molars occluding with the buccal cusps of the lower ones [79], [80], [81], [83] or when the amount of screw expansion reached 10.5mm [12]. One study<sup>(82)</sup> didn't mention the reference for stopping the activation. Retention protocol varied among the six studies from leaving the appliance in situ for an average of 5 months [81], [83],

65 days [12] and one week then replacement by a removable appliance [79]. The method of retention wasn't mentioned in two studies [80], [82] and only the immediate RME effects were evaluated. The characteristics of the included studies are shown in Table 3.

### Risk of Bias within the Studies

As mentioned previously, no RCTs were included in the qualitative assessment, and so the MINORS tool was used for quality assessment. Upon applying the Minors methodological index for grading of the 6 included articles, the resulting scores were found to be below the ideal global score 24 [20]. The highest score recorded was 21[83], and the last one was 13 [79]. All the studies had lack of blinding in measurements. The studies didn't report any dropouts or change in follow up sample ratio and lacked proper sample size calculations Table 4.

### Results of Individual Studies

Treatment effects of the RME were analyzed according to each study's main outcome: postero-anterior radiographs [12], digital dental casts [81], [83], acoustic Rhinometry [79], 3 dimensional stereo-photogrammetry [80] and direct measurements of the nasal soft tissues were used [82] and shown in Table 5.

**Table 3: Design, patients' demographic characteristics, treatment duration and retention, interventions, and outcomes of studies included in the qualitative synthesis.**

| Study Design                   | Eligibility Criteria                                      | Inclusion/Exclusion Criteria  |   | Total Sample n= | Grouping                               | Appliance Used / Anchorage   | Average Tx Time | Retention Protocol  | Records Taken, Timing   | Reference of Maturation  | Outcomes  |
|--------------------------------|---|---|---|-----------------|--|--|-----------------|---|---|--|---|
|                                |   | Inclusion Criteria  | Exclusion Criteria  |                 |  |  |                 |   |   |  |   |
| Baccetti et al (2001) [13]     | Retrospective Controlled Trial                            | NA  | NA  | 42              | ETG= 29<br>LTG=13<br>ECG=11<br>LCG=9   | Haas<br>-2 turns/day till the expansion screw reaches 10.25mm                                  | About 21 days   | Haas kept in place for an average of 65 days (42-75 days)                                 | Postero-anterior X-rays<br>T0,T1, T6  | CVM<br>ETG:CVS1-3<br>LTG: CVS 4-6<br>ECG:CVS 1-3<br>LCG:CVS4-6 | -Transverse skeletal measurements.<br>-Dental measurements. |
| Bicakci et al. (2005) [16]     | Prospective Controlled Trial                              | -Transverse maxillary deficiency<br>-Bilateral CB<br>-No history of nasal disease<br>-Presence of adequate nasal cavity space.                            | NA  | 58              | ETG=16<br>LTG=13<br>ECG= 16<br>LCG= 13 | Hyrax (fully tooth& tissue borne)<br>-2turns/ day  | 23-27 days      | Hyrax kept in place for 1w, then a new removable app used for retention for 3m            | Acoustic Rhinometry<br>At T0, T1, T2<br>(T0 & T3 only for controls)         | CVM<br>ETG: CVS1-3<br>LTG: CVS4-6<br>ETC: CVS-3<br>LTC: CVS4-6 | Minimal cross-sectional area (MCA) of the nasal cavity.     |
| Dindaroglu et al. (2016) [17]  | Prospective Controlled Trial                              | -Skeletal Maxillary Transverse Deficiency<br>-Skeletal development not exceeding MP3 cap stage<br>-No history of orthodontic treatment.                   | NA  | 50              | S.G= 25<br>C.G= 25                     | Bonded type expansion appliance (Extended onto occlusal surface of post teeth)<br>-2turns/ day | 14-19 days      | NA  | 3D Stereo-photogrammmtric images in NHP (natural head position)<br>At T0,T1 | MP3 cap stage  | Facial soft tissue changes                                  |
| Geran et al (2006) [18]        | Prospective Controlled Trial with a retrospective control | -Crowding<br>-Lingual CB<br>-Esthetics<br>-The tendency towards a CI.II malocclusion  | NA  | 77              | S.G=51<br>C.G= 26                      | Bonded acrylic splint appliances (Bonded acrylic splints covered upper D, E, 6)<br>-1turn/day  | NA              | Appliance left in place for 5m, then stabilisation with a palatal plate full time for 12m | Digital dental casts<br>T0, T6 (after completion of phase II treatment), T7 | CVM (CVS 1-3)  | Transverse dental changes, molar angulation                 |
| Santariello et al. (2014) [19] | Prospective Controlled Clinical Trial                     | NA  | NA  | 102             | S.G= 61<br>C.G= 41                     | Banded Hyrax on upper 6<br>-Protocol of activation NA  | NA              | NA  | Nasal Soft Tissue Width Measurement.<br>T0, T1, T3                          | CVM (CVS 1-3)  | Changes in nasal soft tissue dimensions                     |
| Ugolini et al. (2016) [20]     | Prospective Controlled Clinical Trial                     | -Unilateral posterior CB<br>-Early or mid-mixed dentition stage<br>-Angle CI.I or CI.III<br>- CVS 1-3<br>-No orthodontic treatment in maxilla or mandible | -Angle CI.III<br>-Previous orthodontic treatment.<br>-Hormonal imbalances<br>-TMD, Arthritis<br>-Craniofacial Abnormalities | 48              | S.G= 33<br>C.G= 15                     | -Banded Haas, tooth tissue supported<br>-2turns/ day   | 3w              | Haas was left in situ 5-9 m   | Digital dental casts<br>At T0, T4   | CVM (CVS 1-3)  | Transverse mandibular dental changes.                       |

CB; crossbite, CVS; cervical vertebral stage, ETG; early treated group, ETC: early treated control, LTG; late treated group, LTC; late treated control, SG; study group, CG; control group, T0; pre-expansion, T1; immediate post-expansion, T2; 3 months post-expansion, T3; 6 months post expansion, T4; 1 year post- expansion, T5; 3 years post- expansion, T6; 5 years post-expansion, T7: 10 years post expansion.



Long term treatment changes with RME followed by fixed appliances were assessed by Geran et al., [81] in pre-pubertal subjects.

Their results showed that RME produced greater increments in all variables for maxillary and mandibular arch widths when compared with the controls. Maxillary arch perimeter increased in the study group by (0.9 mm) and decreased in the control group by (1.8 mm). On the other hand, the mandibular arch perimeter decreased less in the study group by (-2.4 mm) versus (- 4.4 mm) in the control group. Long term treatment changes with RME followed by fixed appliances were assessed by Geran et al., [81] in pre-pubertal subjects. Their results showed that RME produced greater increments in all variables for maxillary and mandibular arch widths when compared with the controls. Maxillary arch perimeter increased in the study group by (0.9 mm) and decreased in the control group by (1.8 mm). On the other hand, the mandibular arch perimeter decreased less in the study group by (-2.4 mm) versus (- 4.4 mm) in the control group.

Regarding the effect of RME on soft tissue nasal width in pre-pubertal subjects, Santariello et al., [82] reported an increase in the distance between the widest points of the right and left alae of ( $0.8 \pm 0.2$  mm) in the study group with insignificant changes in the distance between the widest points of the insertion of the nose into the soft tissues of the face. When compared to the control group after a retention period of 6 months, a statistically significant increase of only the GAC width was found in the study group.

Spontaneous mandibular response to RME in pre-pubertal subjects was assessed by Ugolini et al., [83] 15 months post-expansion. The study reported a significant increase in the mandibular intermolar width by 1.9 mm. Control subjects showed a tendency towards contraction of the transverse dimensions and a decrease in a molar, canine, and inferior incisor angulation values.

## Discussion

### **Summary of Evidence**

Since the debate is still ongoing, and there is no exact recommendation of the best timing or age to perform a successful RME, hence the aim of the present review had emerged. The objective of this systematic review was to search the literature for any valid evidence supporting the effects of RME in pre and post-pubertal subjects.

RME has many effects on the nasomaxillary complex other than transverse maxillary expansion and correction of crossbites. Effects of RME were studied on skeletal transverse, vertical and

anteroposterior position of the nasomaxillary structures [84], [85], maxillary and mandibular transverse arch dimensions [86], [87], upper airway dimensions [88], [89], bite force, changes in the masticatory cycle and occlusal force distribution [90], swallowing [25] and condylar response [91], [92], [93], changes in head posture and scapular position, natural head position [94], enamel demineralization and white spot lesion formation [23], [57], hearing loss [95], [96], nocturnal enuresis [21], [97], [98], eruption of 3<sup>rd</sup> molars [55], Class II div1 [99], speech and voice function [58], [100], [101], obstructive sleep apnea [102], tongue posture [103] and Holdaway soft tissue analysis [104]. Effects of RME have also been studied on basal bone changes even in the absence of crossbites [105].

For almost four decades, since Melsen [2] revealed the three maturational stages of the MPS, concepts surrounding RME had greatly changed. Great variability of suture fusion among different age groups had been shown [5], [8], [106] Trials that correlate the MPS maturation with different skeletal maturity methods are increasingly aiming to find the eldest adolescent or youngest adult that could benefit from RME treatment [7], [10], [11], [13], [18], [107]. Hence, the time to shift to using the individual's skeletal age rather than the chronological age to gain the maximum effects out of non-surgical RME. Thus, one of the most important inclusion criteria of the studies in this review was dividing the sample according to their skeletal age. Moreover, including studies with untreated control groups was also important to overcome the confounding effect of the craniofacial growth during the study period [35], [43]. The control group had to have a transverse skeletal deficiency as well because various malocclusions are associated with a distinctive craniofacial pattern [108], so it would make more sense to compare subjects with the same malocclusion.

Upon searching the literature, this systematic review included both prospective and retrospective controlled studies that classify their samples according to their skeletal age. Sixty-seven articles were read by full text, and only 6 articles were finally included, four of which were prospective controlled trials, one was a prospective trial with a retrospective control and the final study was a retrospective trial. No well-designed RCTs were found to be eligible for this search, and most of the trials included in this search had methodological problems. None of the studies included assessed the effects of bone-borne RME and instead, tissue and tooth borne RME were studied. Bone borne RME might have yielded different results in this case. The outcomes of the six studies were quite variable as shown in Table 3. Two trials studied the effects of RME on the dental arches: Geran et al., [81] measured the change in both the upper and lower intermolar width, interpremolar width, intercanine width and molar angulation after 5 years of expansion in pre-pubertal patients.

On the other hand, Ugolini et al., [83] measured the changes mentioned above only in the lower arch 15 months following RME, and so the pooling of the results into a quantitative search was impossible as the study endpoints were extremely different. Hence, a meta-analysis wasn't performed. Although non-comparable to one another, the conclusions of Geran et al., [81] resembled that of an earlier systematic review [109] evaluating the long-term dental changes following RME. Two studies [11], [45] included in the mentioned review evaluated the long-term changes on dental casts showing an increase in both the maxillary and mandibular intermolar width with the increase being greater in the maxillary arch.

Since the demand for perfect esthetics is becoming as important as obtaining comfortable, healthy and stable results after orthodontic treatment, the effect of RME on facial soft tissues was considered as an important secondary outcome in this review. These changes were measured by Dindaroglu et al., [80] using 3 D stereophotogrammetric photographs immediately following RME via 3D deviation analyses on 3D stereophotogrammetric images. Being in close relation to the nasomaxillary complex, the nose changes was of great importance and showed a mean maximum deviation limit of  $0.77 \pm 0.34\text{mm}$  and  $-0.94 \pm 0.41\text{ mm}$ . On the other hand, Santariello et al., [82] targeted the changes in nasal soft tissue widths following RME in pre-pubertal subjects only. Although statistically significant, the increase in the width of the base of the nose was considered clinically insignificant. Nasal measurements were also the primary outcome of interest of Bicakci et al., [79]; the trial measured the change in the MCA of the nasal cavity using acoustic Rhinometry following RME in pre and post-pubertal subjects. It concluded that RME not only expanded the maxilla but decreased the nasal resistance through effectively increasing the nasal MCA, being more effective in pre-pubertal than post-pubertal patients. This finding is emphasised by a recent trial [110] that assessed the short-term nasal changes following RME in mouth breathers using cone beam computed tomography. The study concluded that not only does RME increases the nasal airway volume, but is also capable of changing the nasal form and function, increasing a mouth breather chances to gradually attain normal nasal breathing. In 2016, a systematic review of systematic reviews and meta-analyses [111] was conducted to assess the dental and skeletal effects of palatal expansion techniques among different chronological age groups. It reported that significantly more favourable immediate skeletal changes occurred when RME was performed before the pubertal growth peak, based on low evidence [112], [113]. This systematic review, however, didn't strictly select the articles that used the skeletal age indices in their subject recruitment strategy. Conclusions of that review were mainly based on the study by Bacetti et al., [12] that assessed the

transverse skeletal and dental effects of RME in pre and post-pubertal subjects and reported that maxillary adaptations to RME are more likely to be of a dento-alveolar origin rather than a skeletal one when expansion was performed in post-pubertal subjects. Although being the first study to assess the impact of skeletal age on the results of RME, the trial had many limitations. In addition to being a retrospective study, the inclusion and exclusion criteria of the recruited sample were questionable. It wasn't clear if RME was the sole treatment received by the study group or if expansion was performed in conjunction with other treatment modalities as part of a comprehensive orthodontic treatment. Multiple confounders could have been present at baseline, were the only reported baseline similarity between the study and control groups was the skeletal maturational stage without referring to the type of malocclusion of the control group. Comparing changes following RME in transverse skeletal deficiencies to normal skeletal relationships might be misleading. Also, a great variation among the group sizes was observed where 29 subjects were in the early treated group versus 11 controls, and 13 subjects were in the late treated group versus 9 controls. The study had a high detection bias due to the absence of blinding of outcome assessment. Definitive conclusions can't be withdrawn from such a trial owing to the mentioned methodological limitations. Correctly choosing the most applicable tool for quality assessment of the non-randomized studies included in the review was of great importance. An ideal index should be simple, yet highly sensitive and reliable, and hence MINORS was used [20], [114]. Since the global ideal score is 24 for comparative studies [20], any article scoring less than 24 points will be considered as of 'poor quality'. In our search, the included articles scored 14 [12], 13 [79], 18 [80], 15 [81], 15 [82] and 21[83] points in the MINORS scale and hence, all studies give poor-quality evidence supporting their outcome. A definitive conclusion cannot, therefore, be drawn neither upon the short- or long-term skeletal effects of RME nor upon its effects on the nasal airway, maxillary and mandibular dental arches, facial and nasal soft tissues in pre and post-pubertal subjects.

**Strengths and Limitations:** The current review search strategy had strict inclusion and exclusion criteria, had no language restrictions, included multiple electronic database searches and involved a detailed manual searching process. The evidence retrieved from this systematic review has to be interpreted with caution as retrospective studies were included.

## Conclusions

According to the existing evidence from this review, the following conclusions could be stated regarding the transverse skeletal effects of RME in

pre and post-pubertal subjects:

1. The quality level of the studies included in this review was not sufficient to draw conclusive evidence regarding the transverse skeletal effects of RME in pre and post-pubertal subjects.

2. Studies considering the skeletal age for a successful RME treatment are very scarce in the literature. The only available weak evidence suggests that patients treated by tooth-tissue borne RME before the pubertal peak exhibit more increase in the skeletal transverse dimension than do post-pubertal ones, with the pre-pubertal sample being more stable in their long-term treatment results than the post-pubertal.

3. Regarding the transverse skeletal effects of RME, as well as its effect on maxillary and mandibular dental arches, nasal airway, facial and nasal soft tissues, weak evidence supports positive changes in those outcomes in pre-pubertal subjects.

4. For post-pubertal subjects, the only available low-quality evidence suggests an increase in the nasal cross-sectional area immediately following RME.

*Implications for Research:* Carrying out a well-designed randomised controlled clinical trial is strongly recommended to assess the various effects of RME in pre and post-pubertal subjects. Future research should recruit the samples according to their skeletal age and should ensure the presence of a true skeletal maxillary constriction in both the study and control groups.

## References

- Petren S, Bondemark L, Soderfeldt B. A Systematic Review Concerning Early Orthodontic Treatment of Unilateral Posterior Crossbite. *Angle Orthod.* 2003; 73(5):588-96. PMID:14580028
- Melsen B. Palatal growth studied on human autopsy material. A histologic microradiographic study. *Am J Orthod.* 1975; 68(1):42-54. [https://doi.org/10.1016/0002-9416\(75\)90158-X](https://doi.org/10.1016/0002-9416(75)90158-X)
- Thilander B, Persson M. Palatal Suture Closure in Man from 15 to 35 years of age. *Am J Orthod.* 1977; 72(1):42-52. [https://doi.org/10.1016/0002-9416\(77\)90123-3](https://doi.org/10.1016/0002-9416(77)90123-3)
- Wehrbein H, Yildizhan F. The mid-palatal suture in young adults . A radiological-histological investigation. *Eur Orthod Soc.* 2001; 23:105-14. <https://doi.org/10.1093/ejo/23.2.105>
- Revelo B, Fishman LS. Maturation evaluation of ossification of the midpalatal suture. *Am J Orthod Dentofac Orthop.* 1994; 105(3):288-92. [https://doi.org/10.1016/S0889-5406\(94\)70123-7](https://doi.org/10.1016/S0889-5406(94)70123-7)
- Angelieri F, Franchi L, Cevidanes LHS, McNamara JAJ. Diagnostic performance of skeletal maturity for the assessment of midpalatal suture maturation. *Am J Orthod Dentofac Orthop.* 2015; 148(6):1010-6. <https://doi.org/10.1016/j.ajodo.2015.06.016> PMID:26672707
- Angelieri F, Franchi L, Cevidanes LHS, Bueno-Silva B, McNamara JAJ. Prediction of rapid maxillary expansion by assessing the maturation of the midpalatal suture on cone beam CT. *Dental Press J Orthod.* 2016; 21(6):115-25. <https://doi.org/10.1590/2177-6709.21.6.115-125.sar> PMID:28125147 PMID:PMC5278941
- Knaup B, Yildizhan F, Wehrbein H. Age-Related Changes in the Midpalatal Suture. Altersveränderungen der Sutura palatina mediana. *J Orofac Orthop.* 2004; (6):467-74. <https://doi.org/10.1007/s00056-004-0415-y> PMID:15570405
- Angelieri F, Cevidanes LHS, Franchi L, Goncalves JR, Benavides E, McNamara JAJ. Midpalatal suture maturation: classification method for individual assessment before rapid maxillary expansion. *Am J Orthod Dentofac Orthop.* 2013; 144(5):759-69. <https://doi.org/10.1016/j.ajodo.2013.04.022> PMID:24182592 PMID:PMC4185298
- Thadani M, Shenoy U, Patle B, Kalra A, Goel S TN. Midpalatal Suture Ossification and Skeletal Maturation?: A Comparative Computerized Tomographic Scan and Roentgenographic Study. *J Ind Acad Oral Med Radiol.* 2010; 22:81-7. <https://doi.org/10.5005/jp-journals-10011-1020>
- Handelman CS, Wang L, BeGole EA, Haas AJ. Nonsurgical rapid maxillary expansion in adults: report on 47 cases using the Haas expander. *Angle Orthod.* 2000; 70(2):129-44. PMID:10833001
- Baccetti T, Franchi L, Cameron CG, McNamara JAJ. Treatment timing for rapid maxillary expansion. *Angle Orthod.* 2001; 71(5):343-50. PMID:11605867
- Lin L, Ahn H, Kim S, Moon S, Kim S, Nelson G. Tooth-borne vs bone-borne rapid maxillary expanders in late adolescence. *Angle Orthod.* 2015; 85(2):253-62. <https://doi.org/10.2319/030514-156.1> PMID:25490552
- Carlson C, Sung J, McComb RW, Machado AW, Moon W. Microimplant-assisted rapid palatal expansion appliance to orthopedically correct transverse maxillary deficiency in an adult. *Am J Orthod Dentofac Orthop.* 2016; 149(5):716-28. <https://doi.org/10.1016/j.ajodo.2015.04.043> PMID:27131254
- Brunetto DP, Sant'Anna EF, Machado AW, Moon W. Non-surgical treatment of transverse deficiency in adults using microimplant-assisted rapid palatal expansion (MARPE). *Dental Press J Orthod.* 2017; 22(1):110-25. <https://doi.org/10.1590/2177-6709.22.1.110-125.sar> PMID:28444019 PMID:PMC5398849
- Lee K-J, Park Y-C, Park J-Y, Hwang W-S. Miniscrew-assisted nonsurgical palatal expansion before orthognathic surgery for a patient with severe mandibular prognathism. *Am J Orthod Dentofac Orthop.* 2010; 137(6):830-9. <https://doi.org/10.1016/j.ajodo.2007.10.065> PMID:20685540
- Park JJ, Park Y-C, Lee K-J, Tahk JH, Choi YJ. Skeletal and dentoalveolar changes after miniscrew- assisted rapid palatal expansion in young adults?: A cone-beam computed tomography study. *Korean J Orthod.* 2017; 47(2):77-86. <https://doi.org/10.4041/kjod.2017.47.2.77> PMID:28337417 PMID:PMC5359634
- Choi S-H, Shi K-K, Cha J-Y, Park Y-C, Lee K-J. Nonsurgical miniscrew-assisted rapid maxillary expansion results in acceptable stability in young adults. *Angle Orthod.* 2016; 86(5):713-20. <https://doi.org/10.2319/101415-689.1> PMID:26938955
- Higgins JP, Green S. *Cochrane Handbook for Systematic Reviews of interventions* [version 5.1.0, updated March 2011]. The Cochrane Collaboration, 2011. [www.cochrane-handbook.org](http://www.cochrane-handbook.org)
- Slim K, Nini E, Damien F, Kwiatkowski F, Panis Y, Chipponi J. Methodological Index for non- randomized studies (MINORS) : Development and validation of a new instrument. *ANZ J Surg.* 2003; 73(May):712-6. <https://doi.org/10.1046/j.1445-2197.2003.02748.x> PMID:12956787
- Al-Taai N, Alfatlawi F, Ransjo M, Fakhry S. Effect of rapid maxillary expansion on monosymptomatic primary nocturnal enuresis. *Angle Orthod.* 2015; 85(1):102-8. <https://doi.org/10.2319/031014-172.1> PMID:25075913
- Aziz T, Wheatley FC, Ansari K, Lagravere M, Major M, Flores-Mir C. Nasal septum changes in adolescent patients treated with rapid maxillary expansion. *Dental Press J Orthod.* 2016; 21(1):47-

53. <https://doi.org/10.1590/2177-6709.21.1.047-053.oar> PMID:27007761 PMCID:PMC4816585
23. Baysal A, Ulusoy SN, Uysal T. Evaluation of enamel demineralization in adolescents after rapid maxillary expansion using the quantitative light-induced fluorescence method: A single-center, randomized controlled clinical trial. *Am J Orthod Dentofac Orthop.* 2016; 150(5):731-9. <https://doi.org/10.1016/j.ajodo.2016.06.014> PMID:27871699
24. Baysal A, Ozturk MA, Sahan AO, Uysal T. Facial soft-tissue changes after rapid maxillary expansion analyzed with 3-dimensional stereophotogrammetry: A randomized, controlled clinical trial. *Angle Orthod.* 2016; 86(6):934-42. <https://doi.org/10.2319/111315-766.1> PMID:27058647
25. Cappellette Jr. M, Alves FEMM, Nagai LHY, Fujita RR, Pignatari SSN. Impact of rapid maxillary expansion on nasomaxillary complex volume in mouth-breathers. *Dental Press J Orthod.* 2017; 22(3):79-88. <https://doi.org/10.1590/2177-6709.22.3.079-088.oar> PMID:28746491 PMCID:PMC5525449
26. Chang JY, Mcnamara JA. A longitudinal study of skeletal side effects induced by rapid maxillary expansion. *Am J Orthod Dentofac Orthop.* 1997; 112(3):330-7. [https://doi.org/10.1016/S0889-5406\(97\)70264-6](https://doi.org/10.1016/S0889-5406(97)70264-6)
27. Chiari S, Romsdorfer P, Swoboda H, Bantleon HP, Freudenthaler J. Effects of rapid maxillary expansion on the airways and ears. A pilot study. *Eur J Orthod.* 2009; 31(2):135-41. <https://doi.org/10.1093/ejo/cjn092> PMID:19073961
28. Cozza P, Giancotti A, Petrosino A. Rapid palatal expansion in mixed dentition using a modified expander: a cephalometric investigation. *J Orthod.* 2001; 28(2):129-34. <https://doi.org/10.1093/ortho/28.2.129> PMID:11395527
29. Cozzani M, Guiducci A, Mirengi S, Mutinelli S, Siciliani G. Arch width changes with a rapid maxillary expansion appliance anchored to the primary teeth. *Angle Orthod.* 2007; 77(2):296-302. [https://doi.org/10.2319/0003-3219\(2007\)077\[0296:AWCWAR\]2.0.CO;2](https://doi.org/10.2319/0003-3219(2007)077[0296:AWCWAR]2.0.CO;2)
30. Cross DL, McDonald JP. Effect of rapid maxillary expansion on skeletal, dental, and nasal structures: a postero-anterior cephalometric study. *Eur J Orthod.* 2000; 22(5):519-28. <https://doi.org/10.1093/ejo/22.5.519>
31. da Silva Filho OG, Montes LA, Torelly LF. Rapid maxillary expansion in the deciduous and mixed dentition evaluated through posteroanterior cephalometric analysis. *Am J Orthod Dentofac Orthop.* 1995; 107(3):268-75. [https://doi.org/10.1016/S0889-5406\(95\)70142-7](https://doi.org/10.1016/S0889-5406(95)70142-7)
32. Hakan E, Palomo JM. Three-dimensional evaluation of upper airway following rapid maxillary expansion: a CBCT study. *Angle Orthod.* 2014; 84(2):265-73. <https://doi.org/10.2319/012313-71.1> PMID:23865825
33. Farronato G, Giannini L, Galbiati G, Maspero C. RME: influences on the nasal septum. *Minerva Stomatol.* 2012; 61(4):125-34. PMID:22441415
34. Franchi L, Baccetti T, Cameron CG, Kucipal EA, McNamara Jr. JA. Thin-plate spline analysis of the short- and long-term effects of rapid maxillary expansion. *Eur J Orthod.* 2002; 24(2):143-50. <https://doi.org/10.1093/ejo/24.2.143> PMID:12001550
35. Garib DG, Henriques JFC, Carvalho PEG, Gomes SC. Longitudinal effects of rapid maxillary expansion. *Angle Orthod.* 2007; 77(3):442-8. [https://doi.org/10.2319/0003-3219\(2007\)077\[0442:LEORME\]2.0.CO;2](https://doi.org/10.2319/0003-3219(2007)077[0442:LEORME]2.0.CO;2)
36. Gohl E, Nguyen M, Enciso R. Three-dimensional computed tomography comparison of the maxillary palatal vault between patients with rapid palatal expansion and orthodontically treated controls. *Am J Orthod Dentofac Orthop.* 2010; 138(4):477-85. <https://doi.org/10.1016/j.ajodo.2008.12.025> PMID:20889054
37. Hyla-Klekot L, Truszel M, Paradysz A, Postek-Stefanska L, Zyczkowski M. Influence of Orthodontic Rapid Maxillary Expansion on Nocturnal Enuresis in Children. *Biomed Res Int.* 2015; 2015:201039. <https://doi.org/10.1155/2015/201039> PMID:26351629 PMCID:PMC4553176
38. Ileri Z, Basciftci FA. Asymmetric rapid maxillary expansion in true unilateral crossbite malocclusion: a prospective controlled clinical study. *Angle Orthod.* 2015; 85(2):245-52. <https://doi.org/10.2319/011214-40> PMID:24963912
39. Lagravere MO, Carey J, Heo G, Toogood RW, Major PW. Transverse, vertical, and anteroposterior changes from bone-anchored maxillary expansion vs traditional rapid maxillary expansion: a randomized clinical trial. *Am J Orthod Dentofac Orthop.* 2010; 137(3):304.e1-12.
40. Lagravere MO, Gamble J, Major PW, Heo G. Transverse dental changes after tooth-borne and bone-borne maxillary expansion. *Int Orthod.* 2013; 11(1):21-34. <https://doi.org/10.1016/j.ortho.2012.12.003> PMID:23375921
41. Leonardi R, Caltabiano M, Cavallini C, Sicurezza E, Barbato E, Spampinato C, et al. Condyle fossa relationship associated with functional posterior crossbite, before and after rapid maxillary expansion. *Angle Orthod.* 2012; 82(6):1040-6. <https://doi.org/10.2319/112211-725.1> PMID:22519882
42. Lippold C, Hoppe G, Moiseenko T, Ehmer U, Danesh G. Analysis of condylar differences in functional unilateral posterior crossbite during early treatment--a randomized clinical study. *J Orofac Orthop Fortschritte der Kieferorthopädie Organ/official J Dtsch Gesellschaft für Kieferorthopädie.* 2008; 69(4):283-96. <https://doi.org/10.1007/s00056-008-0803-9> PMID:18797832
43. Lippold C, Stamm T, Meyer U, Vegh A, Moiseenko T, Danesh G. Early treatment of posterior crossbite. A randomised clinical trial. *Trials.* 2013; 14:20. <https://doi.org/10.1186/1745-6215-14-20> PMID:23339736 PMCID:PMC3560255
44. Martins LP. Análise dos resultados do tratamento da mordida cruzada posterior funcional com o expansor fixo Hyrax TT - The use of fixed Hyrax expander on the outcomes of functional posterior crossbite treatment, 2011:53.
45. McNamara JAJ, Baccetti T, Franchi L, Herberger TA. Rapid maxillary expansion followed by fixed appliances: a long-term evaluation of changes in arch dimensions. *Angle Orthod.* 2003; 73(4):344-53. PMID:12940553
46. McNamara JAJ, Sigler LM, Franchi L, Guest SS, Baccetti T. Changes in occlusal relationships in mixed dentition patients treated with rapid maxillary expansion. A prospective clinical study. *Angle Orthod.* 2010; 80(2):230-8. <https://doi.org/10.2319/040309-192.1> PMID:19905846
47. Micheletti K, Iwaki L, Provenzano MG, Cuoghi O, Ramos A. Effects of rapid maxillary expansion with six months of retention and no further orthodontic treatment. *Acta Sci Heal Sci.* 2016; 38:89. <https://doi.org/10.4025/actascihealthsci.v38i1.26431>
48. Mutinelli S, Cozzani M. Rapid maxillary expansion in early-mixed dentition: effectiveness of increasing arch dimension with anchorage on deciduous teeth. *Eur J Paediatr Dent.* 2015; 16(2):115-22. PMID:26147817
49. Mutinelli S, Manfredi M, Guiducci A, Denotti G, Cozzani M. Anchorage onto deciduous teeth: effectiveness of early rapid maxillary expansion in increasing dental arch dimension and improving anterior crowding. *Prog Orthod.* 2015; 16:22. <https://doi.org/10.1186/s40510-015-0093-x> PMID:26154156 PMCID:PMC4495101
50. Phatouros A, Goonewardene MS. Morphologic changes of the palate after rapid maxillary expansion: a 3-dimensional computed tomography evaluation. *Am J Orthod Dentofac Orthop.* 2008; 134(1):117-24. <https://doi.org/10.1016/j.ajodo.2007.05.015> PMID:18617111
51. Piacino MG, Cordero-Ricardo M, Cannavale R, Vallelonga T, Garagiola U, Merlo A. Improvement of masticatory kinematic parameters after correction of unilateral posterior crossbite: Reasons for functional retention. *Angle Orthod.* 2017; 3. <https://doi.org/10.2319/020917-98.1>
52. Pinheiro FH de SL, Garib DG, Janson G, Bombonatti R, Freitas MR de. Longitudinal stability of rapid and slow maxillary expansion. *Dental Press J Orthod.* 2014; 19(6):70-7. <https://doi.org/10.1590/2176-9451.19.6.070-077.oar> PMID:25628082 PMCID:PMC4347413

53. Primozic J, Baccetti T, Franchi L, Richmond S, Farnik F, Ovsenik M. Three-dimensional assessment of palatal change in a controlled study of unilateral posterior crossbite correction in the primary dentition. *Eur J Orthod.* 2013; 35(2):199-204. <https://doi.org/10.1093/ejo/cjr125> PMID:23524586
54. Primozic J, Perinetti G, Contardo L, Ovsenik M. Diagnostic performance of 3-dimensional evaluation of palatal vault changes in assessing successful treatment of constricted maxilla in growing subjects. *Am J Orthod Dentofac Orthop.* 2013; 143(1):42-9. <https://doi.org/10.1016/j.ajodo.2012.08.022> PMID:23273359
55. Sokucu O, Ozturk F, Babacan H, Bicakci AA. Does rapid maxillary expansion affect the eruption of upper third molars? *Angle Orthod.* 2008; 78(2):195-200. <https://doi.org/10.2319/031107-119.1> PMID:18251607
56. Thilander B, Lennartsson B. A study of children with unilateral posterior crossbite, treated and untreated, in the deciduous dentition: occlusal and skeletal characteristics of significance in predicting the long-term outcome. *J Orofac Orthop Fortschritte der Kieferorthopädie Organ/official J Dtsch Gesellschaft für Kieferorthopädie.* 2002; 63(5):371-83. <https://doi.org/10.1007/s00056-002-0210-6> PMID:12297966
57. Yagci A, Korkmaz YN, Buyuk SK, Yagci F, Atilla AO. White spot lesion formation after treatment with full-coverage rapid maxillary expanders. *Am J Orthod Dentofac Orthop.* 2016; 149(3):331-8. <https://doi.org/10.1016/j.ajodo.2015.08.015> PMID:26926020
58. Yurttadur G, Basciftci FA, Ozturk K. The effects of rapid maxillary expansion on voice function. *Angle Orthod.* 2017; 87(1):49-55. <https://doi.org/10.2319/010416-008.1> PMID:27463698
59. Zhao Y, Nguyen M, Gohl E, Mah JK, Sameshima G, Enciso R. Oropharyngeal airway changes after rapid palatal expansion evaluated with cone-beam computed tomography. *Am J Orthod Dentofac Orthop.* 2010; 137(4 Suppl):S71-8. <https://doi.org/10.1016/j.ajodo.2008.08.026> PMID:20381764
60. Mc Donald JP. The Effect of RME on Nasal Airway Resistance, craniofacial morphology and head posture. PhD Thesis, Univ Edinbrugh. 1994; 1-237.
61. Gordon JM. Exploratory Work on the Effect of Rapid Maxillary Expansion on Nasal Airway Dimensions. MSc Thesis, Univ Alberta. 2010; 143.
62. Cross DL. Transverse Dimensional Changes Following Rapid Maxillary Expansion. PhD Thesis, Univ Edinbrugh, 1997.
63. Santos P, Vale T, Moreira J, Braga AC, Costa L, Oliveira P, et al. Assessment of the relationship between the upper and lower arch changes with the opening of the expanding screw after the rapid maxillary expansion. *Bull Group Int Rech Sci Stomatol Odontol.* 2010; 49(1):30-9. PMID:22750264
64. Grassia V, D'Apuzzo F, DiStasio D, Jamilian A, Lucchese A, Perillo L. Upper and lower arch changes after Mixed Palatal Expansion protocol. *Eur J Paediatr Dent.* 2014; 15(4):375-80. PMID:25517583
65. Grassia V, D'Apuzzo F, Ferrulli VE, Matarese G, Femiano F, Perillo L. Dento-skeletal effects of mixed palatal expansion evaluated by postero-anterior cephalometric analysis. *Eur J Paediatr Dent.* 2014; 15(1):59-62. PMID:24745595
66. Aggarwal A, Nandakishore P, Nayak USK. A maxillary expander for treatment of unilateral posterior crossbite with anterior open bite. *J Clin Orthod.* 2009; 43(10):655-63. PMID:20128196
67. Alessandri Bonetti G, Marini I, Rizzi R. Disconnectable rapid palatal expander. *J Clin Orthod.* 1996; 30(6):334-6. PMID:10356526
68. De Mendonca MR, Pereira ALP, Fabre AF, Busato MCA, Cuoghi OA. Immediate reactivation of a modified Haas expander after full screw expansion. *J Clin Orthod.* 2008; 42(1):43-8; quiz 34. PMID:18296793
69. Timms DJ, Preston CB, Daly PF. A computed tomographic assessment of maxillary movement induced by rapid expansion - a pilot study. *Eur J Orthod.* 1982; 4(2):123-7. <https://doi.org/10.1093/ejo/4.2.123> PMID:7049711
70. Erdinc AE, Ugur T, Erbay E. A comparison of different treatment techniques for posterior crossbite in the mixed dentition. *Am J Orthod Dentofac Orthop.* 1999; 116(3):287-300. [https://doi.org/10.1016/S0889-5406\(99\)70240-4](https://doi.org/10.1016/S0889-5406(99)70240-4)
71. Grossi VS. Alteracoes dimensionais da maxila em pacientes com atresia do arco dentario superior tratados com aparelho expensor TT - Dimensional changes of the maxilla in patients with atresia of the upper dental arch treated with rapid maxillary expansion appliance. 2012. p. 94.
72. Winsauer H, Walter A, Jaeschke D, Winsauer C, Muchitsch AP, Wendl B. Pure bone-borne palatal expander anchored on orthodontic mini-implants Micro-4 or Micro-6 Expander. *J Stomatol.* 2015; 68(1):6-18. <https://doi.org/10.5604/00114553.1144370>
73. Bass A. Splint Helix. *Funct Orthod.* 2001; 18(2):5-10. PMID:11577637
74. Brogan WF. Rapid maxillary expansion. A stable procedure for improving the nasal airway. *Med J Aust.* 1977; 1(6):167-72.
75. da Silva Filho OG, Ferrari Junior FM, Aiello CA, Zopone N. Correction of posterior crossbite in the primary dentition. *J Clin Pediatr Dent.* 2000; 24(3):165-80. PMID:11314138
76. Hannuksela A, Laurin A, Lehmus V, Kouri R. Treatment of cross-bite in early mixed dentition. *Proc Finn Dent Soc.* 1988; 84(3):175-82. PMID:3045804
77. Karlowska I. Widening of palatine suture in the treatment of compressed maxilla. *Ann Acad Med Stetin.* 1969; 15:211-8. PMID:5377902
78. Simonsen O. Systematic registration and treatment of cross-bite in the primary dentition. *Tandlaegebladet.* 1973; 77(2):171-80. PMID:4570210
79. Bicakci AA, Agar U, Sokucu O, Babacan H, Doruk C. Nasal airway changes due to rapid maxillary expansion timing. *Angle Orthod.* 2005; 75(1):1-6. PMID:15747808
80. Dindaroglu F, Duran GS, Gorgulu S. Effects of rapid maxillary expansion on facial soft tissues?: Deviation analysis on three-dimensional images. *J Orofac Orthop.* 2016; 77(4):242-50. <https://doi.org/10.1007/s00056-016-0033-5> PMID:27272054
81. Geran RG, McNamara JAJ, Baccetti T, Franchi L, Shapiro LM. A prospective long-term study on the effects of rapid maxillary expansion in the early mixed dentition. *Am J Orthod Dentofac Orthop.* 2006; 129(5):631-40. <https://doi.org/10.1016/j.ajodo.2005.01.020> PMID:16679203
82. Santariello C, Nota A, Baldini A, Ballanti F, Cozza P. Analysis of rapid maxillary expansion effects on soft tissue nasal widths. *Minerva Stomatol.* 2014; 63(9):307-14. PMID:25308568
83. Ugolini A, Doldo T, Ghislanzoni LTH, Mapelli A, Giorgetti R, Sforza C. Rapid palatal expansion effects on mandibular transverse dimensions in unilateral posterior crossbite patients: a three-dimensional digital imaging study. *Prog Orthod.* 2016; 17:1. <https://doi.org/10.1186/s40510-015-0114-9> PMID:26746202 PMID:PMC4706536
84. Lima Filho RMA, Ruellas ACO. Long-term anteroposterior and vertical maxillary changes in skeletal class II patients treated with slow and rapid maxillary expansion. *Angle Orthod.* 2007; 77(5):870-4. <https://doi.org/10.2319/071406-293.1> PMID:17685770
85. Akkaya S, Lorenzon S, Ucem TT. A comparison of sagittal and vertical effects between bonded rapid and slow maxillary expansion procedures. *Eur J Orthod.* 1999; 21(2):175-80. <https://doi.org/10.1093/ejo/21.2.175> PMID:10327741
86. Ugolini A, Cerruto C, Di Vece L, Ghislanzoni LH, Sforza C, Doldo T, et al. Dental arch response to Haas-type rapid maxillary expansion anchored to deciduous vs permanent molars: A multicentric randomized controlled trial. *Angle Orthod DONT.* 2014;
87. Grassia V, d'Apuzzo F, Jamilian A, Femiano F, Favero L, Perillo L. Comparison between rapid and mixed maxillary expansion through an assessment of arch changes on dental casts. *Prog Orthod.* 2015; 16:20. <https://doi.org/10.1186/s40510->

- [015-0089-6](#) PMID:26122076 PMCID:PMC4484519
88. Aloufi F, Preston CB, Zawawi KH. Changes in the upper and lower pharyngeal airway spaces associated with rapid maxillary expansion. *ISRN Dent*. 2012; 2012:290964. <https://doi.org/10.5402/2012/290964>
89. Zeng J, Gao X. A prospective CBCT study of upper airway changes after rapid maxillary expansion. *Int J Pediatr Otorhinolaryngol*. 2013; 77(11):1805-10. <https://doi.org/10.1016/j.ijporl.2013.07.028> PMID:24035072
90. Uzuner FD, Odabasi H, Acar S, Tortop T, Darendeliler N. Evaluation of the effects of modified bonded rapid maxillary expansion on occlusal force distribution: A pilot study. *Eur J Dent*. 2016; 10(1):103-8. <https://doi.org/10.4103/1305-7456.175695> PMID:27011748 PMCID:PMC4784139
91. Arat FE, Arat ZM, Acar M, Beyazova M, Tompson B. Muscular and condylar response to rapid maxillary expansion. Part 1: electromyographic study of anterior temporal and superficial masseter muscles. *Am J Orthod Dentofacial Orthop*. 2008; 133(6):815-22. <https://doi.org/10.1016/j.ajodo.2006.07.028> PMID:18538244
92. Arat FE, Arat ZM, Tompson B, Tanju S, Erden I. Muscular and condylar response to rapid maxillary expansion. Part 2: magnetic resonance imaging study of the temporomandibular joint. *Am J Orthod Dentofacial Orthop*. 2008; 133(6):823-9. <https://doi.org/10.1016/j.ajodo.2006.07.029> PMID:18538245
93. Arat FE, Arat ZM, Tompson B, Tanju S. Muscular and condylar response to rapid maxillary expansion. Part 3: magnetic resonance assessment of condyle-disc relationship. *Am J Orthod Dentofacial Orthop*. 2008; 133(6):830-6. <https://doi.org/10.1016/j.ajodo.2007.03.026> PMID:18538246
94. Cerruto C, Di Vece L, Doldo T, Giovannetti A, Polimeni A, Goracci C. A computerized photographic method to evaluate changes in head posture and scapular position following rapid palatal expansion: a pilot study. *J Clin Pediatr Dent*. 2012; 37(2):213-8. <https://doi.org/10.17796/jcpd.37.2.11q67034v1w000wx> PMID:23534333
95. Ceylan I, Oktay H, Demirci M. The effect of rapid maxillary expansion on conductive hearing loss. *Angle Orthod*. 1996; 66(4):301-7. PMID:8863966
96. Micheletti KR, de Mello JA, de Almeida Barreto Ramos SR, Scheibel PC, Scheibel GG, Ramos AL. Effects of rapid maxillary expansion on middle ear function: one-year follow-up. *Int J Pediatr Otorhinolaryngol*. 2012; 76(8):1184-7. <https://doi.org/10.1016/j.ijporl.2012.05.002> PMID:22652498
97. Kuroi J, Modin H, Bjerkhoel A. Orthodontic maxillary expansion and its effect on nocturnal enuresis. *Angle Orthod*. 1998; 68(3):225-32. PMID:9622759
98. Schutz-Fransson U, Kuroi J. Rapid maxillary expansion effects on nocturnal enuresis in children: a follow-up study. *Angle Orthod*. 2008; 78(2):201-8. <https://doi.org/10.2319/021407-71.1> PMID:18251602
99. Baratieri C, Alves MJ, Bolognese AM, Nojima MCG, Nojima LI. Changes in skeletal and dental relationship in Class II Division I malocclusion after rapid maxillary expansion: a prospective study. *Dental Press J Orthod*. 2014; 19(3):75-81. <https://doi.org/10.1590/2176-9451.19.3.075-081.oar> PMID:25162569 PMCID:PMC4296618
100. Sari E, Kilic MA. The effects of surgical rapid maxillary expansion (SRME) on vowel formants. *Clin Linguist Phon*. 2009; 23(6):393-403. <https://doi.org/10.1080/02699200802716074> PMID:19401873
101. Macari AT, Ziade G, Khandakji M, Tamim H, Hamdan A-L. Effect of Rapid Maxillary Expansion on Voice. *J Voice*. 2016; 30(6):760.e1-760.e6. <https://doi.org/10.1016/j.jvoice.2015.09.013> PMID:26572721
102. Villa MP, Rizzoli A, Miano S, Malagola C. Efficacy of rapid maxillary expansion in children with obstructive sleep apnea syndrome: 36 months of follow-up. *Sleep Breath*. 2011; 15(2):179-84. <https://doi.org/10.1007/s11325-011-0505-1> PMID:21437777
103. Ozbek MM, Memikoglu UTT, Altug-Atac AT, Lowe AA. Stability of maxillary expansion and tongue posture. *Angle Orthod*. 2009; 79(2):214-20. <https://doi.org/10.2319/010908-12.1> PMID:19216585
104. Kilic N, Kiki A, Oktay H, Erdem A. Effects of rapid maxillary expansion on Holdaway soft tissue measurements. *Eur J Orthod*. 2008; 30(3):239-43. <https://doi.org/10.1093/ejo/cjn004> PMID:18540012
105. Rosa M, Lucchi P. Rapid Palatal Expansion in the absence of posterior cross-bite to intercept maxillary incisor crowding in the mixed dentition?: Rapid Palatal Expansion in the absence of posterior cross-bite to intercept maxillary incisor crowding in the mixed denti. *Eur J Paediatr Dent*. 2016; 17(4):286-94. PMID:28045316
106. Korbmacher H, Schilling A, Puschel K, Amling M, Kahl-nieke B. computed Tomography Analysis of the Human Midpalatal Suture [Dreidimensionale mikro-computertomographische Analyse der humanen Sutura palatina mediana in Abhangigkeit vom Alter]. *J Orofac Orthop*. 2007; (5):364-76. <https://doi.org/10.1007/s00056-007-0729-7> PMID:17882364
107. Angelieri F, Cevidanes LHS, Franchi L, Goncalves JR, Benavides E, McNamara JAJ. Midpalatal suture maturation: Classification method for individual assessment before rapid maxillary expansion. *Am J Orthod*. 2013; 144:759-69. <https://doi.org/10.1016/j.ajodo.2013.04.022> PMID:24182592 PMCID:PMC4185298
108. Stahl F, Baccetti T, Franchi L, McNamara JA. Longitudinal growth changes in untreated subjects with Class II Division 1 malocclusion. *Am J Orthod Dentofac Orthop*. 2008; 134(1):125-37. <https://doi.org/10.1016/j.ajodo.2006.06.028> PMID:18617112
109. Lagravere MO, Major PW, Flores-Mir C. Long-term dental arch changes after rapid maxillary expansion treatment: a systematic review. *Angle Orthod*. 2005; 75(2):155-61. PMID:15825776
110. Badreddine FR, Fujita RR, Alves FEMM, Cappellette MJ. Rapid maxillary expansion in mouth breathers: a short-term skeletal and soft-tissue effect on the nose. *Braz J Otorhinolaryngol*. 2017.
111. Bucci R, D'Anto V, Rongo R, Valletta R, Martina R, Michelotti A. Dental and skeletal effects of palatal expansion techniques: a systematic review of the current evidence from systematic reviews and meta-analyses. *J Oral Rehabil*. 2016; 43(7):543-64. <https://doi.org/10.1111/joor.12393> PMID:27004835
112. Lagravere MO, Major PW, Flores-Mir C. Long-term Skeletal Changes with Rapid Maxillary Expansion: *Angle Orthod*. 2005; 75(6):1046-52. PMID:16448254
113. Lione R, Franchi L, Cozza P. Does rapid maxillary expansion induce adverse effects in growing subjects? *Angle Orthod*. 2013; 83(1):172-82. <https://doi.org/10.2319/041012-300.1> PMID:22827478
114. Zeng X, Zhang Y, Kwong JSW, Zhang C, Li S, Sun F, et al. The methodological quality assessment tools for preclinical and clinical studies, systematic review and meta-analysis, and clinical practice guideline: A systematic review. *J Evid Based Med*. 2015; 8(1):2-10. <https://doi.org/10.1111/jebm.12141> PMID:25594108