

Phenotypic Identification and Molecular Characterization of *Malassezia Spp.* Isolated from Pityriasis Versicolor Patients with Special Emphasis to Risk Factors in Diyala Province, Iraq

Ahmed Kamil Awad¹, Ali Ibrahim Ali Al-Ezzy^{2*}, Ghassan H. Jameel¹

¹Department of Microbiology, College of Veterinary Medicine, Diyala University, Iraq; ²Department of Pathology, College of Veterinary Medicine, Diyala University, Iraq

Abstract

Citation: Awad AK, Al-Ezzy AIA, Jameel GH. Phenotypic Identification and Molecular Characterization of *Malassezia Spp.* Isolated from Pityriasis Versicolor Patients with Special Emphasis to Risk Factors in Diyala Province, Iraq. Open Access Maced J Med Sci. 2019 Mar 15; 7(5):707-714.
<https://doi.org/10.3889/oamjms.2019.128>

Keywords: Pityriasis versicolor; *Malassezia spp.*; Molecular diagnosis; Risk factors; Iraq

***Correspondence:** Ali Ibrahim Ali Al-Ezzy, Department of Pathology, College of Veterinary Medicine, Diyala University, Iraq, E-mail: alizzibrahim@gmail.com

Received: 19-Nov-2018; **Revised:** 17-Jan-2019; **Accepted:** 18-Jan-2019; **Online first:** 27-Feb-2019

Copyright: © 2019 Ahmed Kamil Awad, Ali Ibrahim Ali Al-Ezzy, Ghassan H. Jameel. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Funding: This research did not receive any financial support

Competing Interests: The authors have declared that no competing interests exist

AIM: The main objective is isolation and molecular characterisation of *Malassezia spp.* from pityriasis versicolor (PV) patients with special emphasis to risk factors in Diyala province, Iraq.

METHODS: Fifty patients (32 males and 18 females) presented with PV, the age ranged (15-45) years were included. Direct wet mount using KOH 10%, culture of skin scraping and PCR were used for confirmatory diagnosis.

RESULTS: *Malassezia spp.* was isolated from (54%) of skin scraping; *M. furfur* (32%); *M. pachydermatis* (8%) and *M. globosa* (14%). The age group (15-22) years were frequently exposed to *Malassezia* infection. A significant inverse correlation was reported between age and exposure to *Malassezia spp.* Infection. Males were frequently exposed to *Malassezia* infection, (40%). A significant correlation was reported between gender and exposure to *Malassezia spp.* Infection. Females were at risk of getting *Malassezia* infection (2.619) time than males. Patient resident in the urban area frequently exposed to *Malassezia* infection, (34%). Patients resident in the rural area appears to be at risk of getting *Malassezia* infection (1.093) time than those in an urban area. Patient with good economic status was frequently exposed to *Malassezia* infection, (36%). Patients with middle economic status appear to be at risk of getting *Malassezia* infection (0.42) time than those with good economic status. Patients with primary education were frequently exposed to *Malassezia* infection, (22%). A significant correlation was reported between education level and exposure to *Malassezia spp.* Infection. No significant correlation was reported between economic status; type of job; source of water; contact with dogs and birds and *Malassezia spp.* Infection.

CONCLUSION: *M. furfur*, *M. pachydermatis* and *M. globosa* represent the most common *Malassezia spp.* causing PV. Using of PCR is very critical to confirm the diagnosis of *Malassezia spp.* *Malassezia* infection inversely correlated with age and positively correlated with females gender and education. The residency in a rural area and middle economic status increase the possibility of infection. Infection was not affected by the source of water; job and contact with dogs and birds.

Introduction

The family *Malassezia* involves a gathering of superficial dimorphic fungi occurring as normal skin biota on the human body; however, they can also cause infection or are related to certain skin ailments. Rarely, they can become invasive, to cause opportunistic systemic infection in the presence of certain predisposing factors [1]. However, the

complexity of the interaction of a unicellular eukaryotic creature (*Malassezia*) with a tissue of a multicellular living being (skin) makes understanding the communications and advancement of infection a complex process [2].

Malassezia yeasts inhabit various body sites including scalp, forehead, shoulder, abdomen, lower axilla, groin and forearm, due to an increased thickness of sebaceous organs at these locales [3]. It

has been reported that *Malassezia* yeasts are related to dermatoses such as pityriasis versicolor, seborrheic dermatitis and *Malassezia* folliculitis, recently, reported is the implication of *Malassezia* yeasts in atopic dermatitis and psoriasis [4]. Pityriasis versicolor (PV) is a superficial fungal infection, characterized by changes in skin pigment due to colonization of the stratum corneum by a dimorphic lipophilic fungus of the normal flora of the skin, and its characteristic by multiple macules and/or patches of variable appearance (hypopigmented, hyperpigmented, dark brown or erythematous) surrounded by normal skin [5]. Pityriasis versicolor is a worldwide in distribution, particularly in tropical areas. Infection by pityriasis versicolor has varied according to age and gender. It usually infects adults due to increased sebum secretion after puberty.

The main objective of this research is isolation and molecular characterisation of *Malassezia spp.* from pityriasis versicolor (PV) patients with special emphasis to risk factors in Diyala province, Iraq

Material and Methods

Study Area and Study Population

This study was performed in Baqubah city - Diyala province 33°45'34.71" N; 44°36'23.97" E, Northeast. The study included 50 pityriasis versicolor patients, age ranged (15-45) years; (32 males and 18 females) attended to the dermatology outpatient's clinic, Baqubah teaching hospital from 14 of October 2017 to 13 of January 2018.

Collection of Samples

Samples used were hair and skin scrapings. Forceps and surgical blades were used for collecting hair and skin samples. The scales were collected in sterile empty Petri dishes. [6].

Wet mount KOH

Scales specimens were subjected to direct examination by placing on a clean slide mounted with a drop of 10% KOH, covered with a coverslip. The slide was warmed gently and examined under a microscope (40X). Characteristic spaghetti and meatballs appearance (shortly curved hyphae and round yeasts) confirmed the diagnosis of PV [7].

Culture

Scales were inoculated onto Sabouraud's dextrose agar (SDA), containing 0.05 gm/L chloramphenicol, Penicillin at a concentration of 0.4

ml/L and Streptomycin at a concentration of 2 ml/L with olive oil and without olive oil. The vials were incubated at 37°C for 1-2 weeks. All the slants examined on days 3, 7 and then at weekly intervals up to three weeks for any developing colonies [8].

Staining

For examination, 5 ml of sterile distilled water was mixed with a loopful of actively growing yeast, and the concentration was adjusted to about 10^5 cell/ml Smears from the colonies were stained with Gram's stain and examined under a microscope for identification of *Malassezia* [9]. Lactophenol cotton blue stain used for microscopic observation of yeast cells, the suspension of yeast cells were prepared, loopful of culture were stained with lactophenol cotton blue on a sterile glass slide.

Catalase

Catalase test was applied by using a drop of 3% hydrogen peroxide, and the production of gas bubbles was considered a positive reaction [10].

Tween assimilation tests

According to the method reported by [11], yeast cells of (2×10 to 3×10 CFU/ml) was suspended in 1ml sterile distilled water and poured into plate containing SDA with 0.05 gm/L chloramphenicol, Penicillin at concentration of 0.4 ml/L and Streptomycin at concentration of 2 ml/L cooled at about 50°C. The inoculum was then spread evenly. After solidification, four holes were made using a 2 mm diameter punch and filled with 5 μ l of Tween 20, 40, 60 and 80, respectively. The plates were incubated for 1 week at 32°C. The utilisation of Tween was assessed by the degree of growth and/or reaction (precipitation) of the lipophilic yeasts around the wells [11].

Assimilation of Castor oil

Assimilation of castor oil was examined using the agar diffusion test. Ten ml sterile Brain Heart Agar (BHI), was melted and allowed to cool to about 50°C. Yeast cells of (2×10 to 3×10 CFU/ml) were suspended in 1ml sterilised distilled water and poured into a plate containing the medium was then spread evenly. Once the medium had solidified, one hole was made using a sterile 6mm punch and filled with 50 μ l of castor oil. The plates were incubated at 32°C for 10 days and assessed for growth around the individual wells after 2, 4, 6, 8 and 10 days [6].

Splitting of Esculin

Glucosidase activity was assayed by using

the esculin agar tube. Using a loop, the yeast inoculum was deeply inoculated into the agar and incubated at 32°C for 5 days. The splitting of esculin into esculetin and glucose is revealed by the darkening of the medium with the liberation of soluble ferric salt incorporated in the medium [10].

Tryptophan test (Pigment Induction medium)

Yeast cells were cultured on m Dixons medium which was prepared earlier addition of 0.6% of tryptophan instead of peptone to the original medium. After sterilisation and cooling at room temperature, the suspension was smeared on the agar medium using a sterile swab. The plates were incubated at 32°C for 2 to 4 weeks [12].

Preservation of the *Malassezia* isolates

Malassezia spp. isolates were grown in the universal bottle containing 5ml of SDA used as slants, and they were enveloped with parafilm and stocked at 4°C for 3 months [13].

DNA Extraction

DNA was extracted from *Malassezia* spp. By using the QIAamp DNA Mini Kit (Qiagen, Germany) according to the protocol stated by the kit manufacturer [14].

Concentration and purity of DNA

DNA was extracted from a hundred isolated of *Malassezia* spp. And they were concentrated in one tube. The concentration and the purity of the DNA samples were determined by Quantus Fluorometer at (9.9 ng/μl and 57 ng/μl) was used to detect the concentration of extracted DNA to detect the goodness of samples for downstream applications. For 1 μl of DNA, 199 μl of diluted QuantyFlour Dye was mixed. After 5min incubation at room temperature, DNA concentration values were detected, according to the protocol stated by the kit manufacturer (Promega, U.S.A).

Primers selection and preparation

Universal primers ITS3 (GCATCGATGAAGAACGCAGC) and ITS4 (TCC TCC GCT TATTGA TAT GC). The 5.8S rDNA and the it's 2 region, amplified from type, neotype, reference and clinical isolates of *Malassezia* species by using the ITS 3 and 4 primers (Gaitanis, G et al., 2002) [2]. We're synthesised by (QIAGEN, Germany).

PCR working solution

Optimisation of PCR was accomplished after several trials. Thus the following mixture was adopted amplification reactions were produced in the 25μl final volume containing 12.5 μl Go Taq® master mix (Promega, USA), 2μl of the primers and 2μl DNA template and complete the volume by 8.2 ul nuclease-free water

Programmable thermal controller

Program for amplifying the 5.8S rDNA and the it's 2 region, amplified from the type of ITS3 and ITS4 for *Malassezia* spp. For identification of *Malassezia* spp., an initial denaturation step at 95°C for five minutes was followed by thirty cycles of denaturation at 95°C for thirty seconds, annealing at 55°C for thirty seconds, and extension at 72°C for thirty seconds, with a final extension step of 72°C for seven minutes [15].

Agarose Gel Electrophoresis

After PCR amplification, agarose gel electrophoresis was adopted to confirm the presence of amplification. PCR was completely dependable on the extracted DNA criteria, according to the protocol stated by the kit manufacturer (Promega, U.S.A).




Figure 1: Agarose gel electrophoresis (2%) for 100 v/mAmp for 80 min of *Malassezia* spp. DNA products generated through ITS 3 (GCATCGATGAAGAACGCAGC), and ITS4 (TCCTCCGCTTATTGATATGC) primers, stained with Ethidium bromide; M: Molecular marker (100bp); lanes 1H, 2H, 3H: *M. furfur* (509bp); lanes H4: *M. globosa* (430bp); lanes H5: *M. pachydermatis* (483bp)

Statistical Data Analysis

Patients demography and cross tabulation were calculated by Statistical Package for the Social Sciences for Windows version 17 (SPSS, Armonk, NY: IBM Corp.). Pearson's chi-square and Pearson's correlation coefficient were used for the correlation between the variables of the two tests. A p value of ≤ 0.05 and ≤ 0.01 (two-tailed) was set to be statistically significant

Results

Table 1 shows the morphological characteristics and physiological characteristics of *Malassezia* spp. Isolated from human samples. According to these features, the total number of *Malassezia* spp. Isolated from a human was 27/50, (54%), *M. furfur* isolated from 16/50, (32%); *M. pachydermatis* 4/50, (8%) and *M. globosa* were isolated from 7/50, (14%). These results were confirmed by PCR as shown in Figure 1.

Table 1: Diagnosis of *Malassezia* spp. from human samples by morphological and physiological characteristics compared with conventional PCR using ITS3&ITS4 primers

Characteristics	<i>Malassezia</i> Isolates			
	<i>M. furfur</i>	<i>M. pachydermatis</i>	<i>M. globosa</i>	
Morphological characteristics	Colony morphology	Mat, dull, smooth, umbonate or slightly folded with convex elevation	Mat, convex, umbonate (sometimes)	Raised, folded, rough
	Pigmentation	+	+	+
	Cell shape	Oval, cylindrical spherical	Oval	Spherical
Physiological characteristics	Growth on SDA			
	Growth on SDA-castor oil	+	+	+
	Growth on Esculin agar	+	+	+
	Growth on mDA 32°C	+	+	+
	Growth on mDA 37°C	+	+	+
	Growth on mDA 41°C	+	+	+
	Catalase reaction	+	+	+
	Utilization of Tween 20	+	+	+
	Utilization of Tween 40	+	+	+
	Utilization of Tween 60	+	+	+
	Utilization of Tween 80	+	+	+
	Total number of <i>Malassezia</i> isolates	16 (32%)	4 (8%)	7 (14%)
	Diagnosis by Conventional PCR using ITS3&ITS4 primers	16 (32%)	4 (8%)	7 (14%)

As shown in Table 2, the age group (15-22) years was more frequently exposed to *Malassezia* infection, followed by (23-30) year's age group. The age group (39-42) years was the minimally exposed to infection compared with other groups. *M. furfur* was the predominant species recovered from PV cases while *M. pachydermatis* was the least once. The isolation rate was drastically decreased while the age of patients approximates to forties. No significant difference was reported between age groups exposed to *Malassezia* spp. Infection. An inverse significant correlation was reported between age and exposure to *Malassezia* spp. Infection.

As shown in Table 3, males were more frequently exposed to *Malassezia* infection, 20/50, (40%). *M. furfur* was the predominant species recovered from males with PV cases, 11/50, (22%) while *M. globosa* recovered from 5/50, (10%) and *M. pachydermatis* was the least once, isolated from 4/50, (8%).

Table 2: Age as a possible risk factor associated with *Malassezia* infection in human

Descriptive statistic of patients age					
Minimum (years)	15				
Maximum (years)	45				
Mean ± Std. Deviation	27.82 ± 8.39				
Age group (years)	<i>Malassezia</i> spp. infection			Total No. (%) of positive cases	Negative No. (%)
	<i>M. furfur</i>	<i>M. pachydermatis</i>	<i>M. globosa</i>		
15-22	7 (10%)	1 (2%)	3 (6%)	11 (22%)	5 (10%)
23-30	4 (8%)	2 (4%)	3 (6%)	9 (18%)	7 (14%)
31-38	3 (6%)	1 (2%)	0 (0%)	4 (8%)	8 (16%)
39-42	1 (2%)	0 (0%)	0 (0%)	1 (2%)	2 (4%)
43-46	1 (2%)	0 (0%)	1 (2%)	2 (4%)	1 (2%)
Total No. (%)	16 (32%)	4 (8%)	7 (14%)	27 (54%)	23 (46%)
χ ²	27.523				
P value	0.154				
R	-0.284				
P value	0.046				

No significant difference was reported between genders exposed to *Malassezia* spp. Infection. A significant correlation was reported between gender and exposure to *Malassezia* spp. Infection. Females appear to be at risk of getting *Malassezia* infection at (2.619) time than males.

Table 3: Gender as a possible risk factor associated with *Malassezia* infection human

Gender	Positive No.(%)			Total No. of positive cases	Negative No.(%)
	<i>M. furfur</i>	<i>M. pachydermatis</i>	<i>M. globosa</i>		
Male	11 (22%)	4 (8%)	5 (10%)	20 (40%)	12 (24%)
Female	5 (10%)	0 (0%)	2 (4%)	7 (14%)	11 (22%)
Total No. (%)	16 (32%)	4 (8%)	7 (14%)	27 (54%)	23 (46%)
χ ²	3.970				
P value	0.265				
R	0.270				
P value	0.05				
Risk Estimate	95% Confidence Interval				
Odds Ratio for gender (female / male)	Value	Lower	Upper		
	2.619	0.799	8.588		

As shown in Table 4, patient resident in urban area were more frequently exposed to *malassezia* infection, 17/50, (34%). *M. furfur* was the predominant species recovered from patients with PV cases, 11/50, (22%) while *M. globosa* recovered from 4/50, (8%) and *M. pachydermatis* was the least once, isolated from 2/50, (4%).

Neither significant difference nor correlation was reported between residency and exposure to *malassezia* spp. infection. Patients residency in rural area appear to be at risk of getting *malassezia* infection at (1.093) time than those in urban area.

Table 4: Residence as a possible risk factor associated with *Malassezia* infection in human

Residence	Positive NO.(%)			Total No. of positive cases	Negative No.(%)
	<i>M. furfur</i>	<i>M. pachydermatis</i>	<i>M. globosa</i>		
Rural	5 (10%)	2 (4%)	3 (6%)	10 (20%)	9 (18%)
Urban	11 (22%)	2 (4%)	4 (8%)	17 (34%)	14 (28%)
Total No.(%)	16 (32%)	4 (8%)	7 (14%)	27 (54%)	23 (46%)
χ ²	0.023				
P value	0.879				
R	-0.021				
P value	0.882				
Risk Estimate	95% Confidence Interval				
Odds Ratio for Residence (rural / urban)	Value	Lower	Upper		
	1.093	0.348	3.435		

As shown in Table 5, patient with good economic status were more frequently exposed to *malassezia* infection, 18/50, (36%). *M. furfur* was the

predominant species recovered from patients with PV cases, 6/50, (12%) while *M. globosa* recovered from 4/50, (8%). Neither significant difference nor correlation was reported between economic status and exposure to *Malassezia* spp. Infection. Patients with middle economic status appear to be at risk of getting *Malassezia* infection at (0.42) time than those with good economic status.

Table 5: Family economic status as a possible risk factor associated with *Malassezia* infection in human

Economic Status	Positive No.(%)			Total No. of positive cases	Negative No.(%)
	<i>M. furfur</i>	<i>M. pachydermatis</i>	<i>M. globosa</i>		
Middle	6 (12%)	0 (0%)	3 (6%)	9 (18%)	4 (8%)
Good	10 (20%)	4 (8%)	4 (8%)	18 (36%)	18 (36%)
Very good	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (2%)
Total No.(%)	16 (32%)	4 (8%)	7 (14%)	27 (54%)	23 (46%)
χ^2	5.424				
P value	0.491				
R	-0.086				
Risk Estimate	95% Confidence Interval				
Odds Ratio for economic status (middle/good)	Value	Lower	Upper		
	0.421	0.110	1.612		

As shown in Table 6, patients with Primary education were more frequently exposed to *Malassezia* infection, 11/50, (22%). *M. furfur* was the predominant species recovered from patients with PV cases, 8/50, (16%) while *M. globosa* recovered from 3/50, (6%). Illiterate patients were less frequently exposed to *Malassezia* infection, 2/50, (4%). *M. pachydermatis* was the predominant species recovered from patients with PV cases, 2/50, (4%).

No significant differences were reported between education status and *Malassezia* spp. Infection. Significant correlation was reported between education level and exposure to *Malassezia* spp. Infection.

Table 6: Patients education status as possible risk factors associated with *Malassezia* infection in human

Education level	Malassezia spp			Total No. (%) of positive cases	Total No. (%) of negative cases
	<i>M. furfur</i>	<i>M. pachydermatis</i>	<i>M. globosa</i>		
Illiterate	0 (0%)	2 (4%)	0 (0%)	2 (4%)	4 (8%)
Primary	8 (16%)	0 (0%)	3 (6%)	11 (22%)	12 (24%)
Secondary	6 (12%)	1 (2%)	2 (4%)	9 (18%)	6 (12%)
Higher Education	2 (4%)	1 (2%)	2 (4%)	5 (10%)	1 (2%)
Total	16 (32%)	4 (8%)	7 (14%)	27 (54%)	23 (46%)
χ^2	3.785				
P value	0.286				
R	0.280				
P value	0.05				

As shown in Table 7, Students were more frequently exposed to *Malassezia* infection, 10/50, (20%). *M. furfur* was the predominant species recovered from patients with PV cases, 8/50, (16%) while *M. globosa* and *M. pachydermatis* recovered from 1/50, (2%). Patients with the academic job were less frequently exposed to *Malassezia* infection, 2/50, (4%). *M. pachydermatis* and *M. globosa* were the predominant species recovered from patients with PV cases, 2/50, (4%). Neither significant difference nor correlations were reported between the type of job

and *Malassezia* spp. Infection.

Table 7: Patient's job as possible risk factors associated with *Malassezia* infection in human

Job	Malassezia spp			Total No. (%) of positive cases	Total No. (%) of negative cases
	<i>M. furfur</i>	<i>M. pachydermatis</i>	<i>M. globosa</i>		
Free Work (Self-employer)	2 (4%)	0 (0%)	2 (4%)	4 (8%)	3 (6%)
Farmer	2 (4%)	2 (4%)	1 (2%)	5 (10%)	3 (6%)
Housewife	2 (4%)	0 (0%)	1 (2%)	3 (6%)	7 (14%)
Military	2 (4%)	0 (0%)	1 (2%)	3 (6%)	6 (12%)
Student	8 (16%)	1 (2%)	1 (2%)	10 (20%)	3 (6%)
Education	0 (0%)	1 (2%)	1 (2%)	2 (4%)	1 (2%)
Total	16 (32%)	4 (8%)	7 (14%)	27 (54%)	23 (46%)
χ^2	18.981				
P value	0.215				
R	0.063				
P value	0.663				

As shown in Table 8, patient utilizing filtrated water were more frequently exposed to *Malassezia* infection, 21/50, (42%). *M. furfur* was the predominant species recovered from patients with PV cases, 14/50, (28%) while *M. globosa* 4/50, (8%) and *M. pachydermatis* recovered from 3/50, (6%). patients utilizing river water were less frequently exposed to *malassezia* infection, mainly *M. furfur*, 1/50, (2%). Neither significant difference nor correlations were reported between source of water and *Malassezia* spp. infection.

Table 8: Source of water as possible risk factors associated with *Malassezia* infection in human

Source of water	Malassezia spp			Total No. (%) of positive cases	Total No. (%) of negative cases
	<i>M. furfur</i>	<i>M. pachydermatis</i>	<i>M. globosa</i>		
Tap Water	1 (2%)	1 (2%)	3 (6%)	5 (10%)	2 (4%)
Filtrated	14 (28%)	3 (6%)	4 (8%)	21 (42%)	21 (42%)
River Water	1 (2%)	0 (0%)	0 (0%)	1 (2%)	0 (0%)
Total	16 (32%)	4 (8%)	7 (14%)	27 (54%)	23 (46%)
χ^2	8.592				
P value	0.198				
R	-0.219				
P value	0.126				

As shown in Table 9, patients who do not have any type of contact with animals were more frequently exposed to *malassezia* infection, 15/50, (30%). *M. furfur* was the predominant species recovered from patients with PV cases, 10/50, (20%) while *M. globosa* 4/50, (8%) and *M. pachydermatis* recovered from 1/50, (2%). patients who have contact with dogs only were exposed equally to *M. furfur* 1/50, (2 %) and *M. pachydermatis* recovered from 1/50, (2%) infection. Neither significant difference nor correlations were reported between contact with animals and *Malassezia* spp. infection.

Table 9: contact with animals as a possible risk factors associated with *Malassezia* infection in human

contact with Animals	Malassezia spp			Total No. (%) of positive cases	Total No. (%) of negative cases
	<i>M. furfur</i>	<i>M. pachydermatis</i>	<i>M. globosa</i>		
No contact	10 (20%)	1 (2%)	4 (8%)	15 (30%)	13 (26%)
bird	3 (6%)	1 (2%)	1 (2%)	5 (10%)	6 (12%)
dogs	1 (2%)	1 (2%)	0 (0%)	2 (4%)	2 (4%)
Birds and dogs	2 (4%)	1 (2%)	2 (4%)	5 (10%)	2 (4%)
Total	16 (32%)	4 (8%)	7 (14%)	27 (54%)	23 (46%)
χ^2	5.243				
P value	0.813				
R	0.185				
P value	0.199				

Discussion

Malassezia species are part of the resident skin flora of humans and other warm-blooded vertebrates that are discovered in 75-80 % of healthy subjects [16]. These yeasts are associated with a wide spectrum of clinical manifestations from benign skin conditions, such as pityriasis versicolor to fungemia in the immunocompromised host [17]. A molecular analysis-based nonculture method appears to be the most reliable and appropriate method by which to analyse *Malassezia* spp since the isolation media and procedures used does not influence such a method [18]. Therefore molecular methods have revolutionised the study of diseases caused by *Malassezia* species because the organisms are fastidious and difficult to identify. Therefore molecular methods overcome the limits of the conventional methods [19].

The current study showed that *M.furfur* was isolated from (32%) which was more common than *M. globosa* which isolated from (14%) and *M. pachydermatis* which was isolated from (8%).

This study comes in line with that reported by Al-ammari *et al.*, [20], in Baghdad, with different clinical presentations including steroids acne, folliculitis, seborrheic dermatitis and atopic dermatitis. In another study, in Diyala province-Iraq; Al-Ezzy *et al.*, [21], found that *Malassezia* was isolated from (61.7%) of patients with PV specimens with similar age group distribution during 2017 in Baqubah teaching hospital. Current result comes by Shah in India [17], who recovered *Malassezia* spp from (50.35%) of PV patients. Current result disagrees with Shah and Chaudhary *et al.* from India [17], [22], reported the recovery of *M. globosa* from the majority of PV cases and stated that *M. globosa* is more pathogenic than other *Malassezia* species as it has greater enzymatic activity, involving lipase and esterase production.

Current isolation rate come in contrary with that reported by Chaudhary *et al.* and Archana *et al.* from India [22], [23], who recorded that the isolation rate of *Malassezia* species in Indian patients with PV was (70% and 96.66%) respectively. Also, current result comes in contrary with that reported by Shokohi from Iran [8], who recorded (88.4%) as isolation rate from PV cases.

The current study revealed that the age group (15-22) years was frequently exposed to *Malassezia* infection, followed by (23-30) years age group. The age group (39-42) years was the minimally exposed to infection compared with other groups. This result come in accordance with that reported in India and Iraq by [17], [23], [24], [25], [26], [27], [28]. *M. furfur* was the predominant species recovered from PV cases while *M. pachydermatis* was the least once. The isolation rate was drastically decreased while the

age of patients approximates to forties which come in line with Al-ammari from Iraq and Akaza from Japan [20], [28] and disagree with Shah from India, [17] who stated that *M. globosa* was the predominant yeast recovered from PV cases. Also disagree with Archana from India, [23] stated that *M. furfur* represent the second cause of PV after *M. sympodialis*. An inverse significant correlation was reported between age and exposure to *Malassezia* spp. Infection, which agrees with Al-ammari from Iraq and Thayikkannu *et al.*, from India [20], [25].

The current study revealed that males were more frequently exposed to malassezia infection, (40%) . *M. furfur* was the predominant species recovered from males with PV cases, (22%) which come in accordance with studies from Iraq and India [20], [21], [23], [24], [26], [27], [29]. While *M. globosa* recovered from, (10%) and *M. pachydermatis* was the least once, isolated from, (8%). Significant correlation was reported between gender and exposure to malassezia spp infection which come in accordance with Al-Ezzy from Iraq [21]. On the other hand, Hasan *et al.*, [24], stated that military personnel, athletes, and those doing hard works that usually associated with hyper sweating were more vulnerable to pityriasis versicolor infection. Furthermore, sebaceous glands secretion is increased in males 15 years and older compared to females, and that may also promote pityriasis versicolor infection in males.

Females appear to be at risk of getting malassezia infection at (2.619) time than males which come in accordance with Al-Ammari from Iraq [20], stated that many factors play role in *Malassezia* pathogenicity such as increased sebum production , hormonal fluctuations, stress, illness, infrequent shampooing, food allergies, vitamin B deficiency, hair curlers and blow dryers, cold weather (winter), use of hair sprays, gels and hair coloring chemicals which more probably resented in females and hence they become more susceptible to *Malassezia* infections than males.

Current study revealed that patient resident in urban area were more frequently exposed to malassezia infection, (34%) which come in line with Devendrappa and Javed from India [26]. *M. furfur* was the predominant species, (22%) while *M. globosa* recovered from, (8%) and *M. pachydermatis* was the least once, isolated from, (4%). This result disagree with Hasan *et al.*, from Iraq, [24], who reported no significant correlation. On the other hand current study revealed that patients residency in rural area appear to be at risk of getting malassezia infection at (1.093) time than those in urban area. This result come in accordance with Sharma from India [30], stated that geographical variations have been observed in the densities of different *Malassezia* species on the skin. *Malassezia* infections more commonly reported in the warm and humid tropical and subtropical climates which is more suited for its growth. The higher incidence in urban people was mainly due to the

location of the hospital in the city. As the health facilities are also available in rural areas, the attendance of the patients from these areas to this hospital was reduced; probably this could be the reason for the low incidence from rural areas. Another factor could be that the rural population is not bothered about the cosmetic aspect of this disease.

The current study revealed that no correlation was reported between economic status and exposure to *Malassezia* spp. Infection. This may be attributed to factors other than economy play a vital role in infection such as psychological and immunological stress.

The current study revealed that patients with primary education as well as students were more frequently exposed to *Malassezia* infection. Illiterate patients and patients with an academic job were less frequently exposed to *Malassezia* infection, (4%). A significant correlation was reported between education level and exposure to *Malassezia* spp while *Malassezia* spp. Infection does not correlate with the type of job. This result may be attributed to the limited number of illiterate patients seeking medical consultations as well as investigations related to skin infections possibly due to they consider this is a type of luxury checkup. On the other hand, the limited number of patients with higher education reflects their knowledge about the complications of cutaneous infections as well as their attention to check any skin abnormalities such as hyperpigmentation or hypopigmentation or even scales. Type of patient's job may facilitate the flaring of infection indirectly by acting as a stress factor which acts with other genetic, hormonal, and environmental factors such as humidity and temperature to establishing a clinical presentation of PV.

The current study revealed those patients utilising filtrated water were more frequently exposed to *Malassezia* infection. *M. furfur* was the predominant species recovered from patients with PV. Patients utilizing river water were less frequently exposed to *Malassezia* infection, mainly *M. furfur*, (2%) although, no significant correlation was reported between the source of water and *Malassezia* spp. Infection. Current result comes in agreement with Babic *et al.*, from Slovenia [31], who stated that fungi including genus *Malassezia* can form fungal biofilms within 48 h in tap systems in private homes, hospitals or industrial network was confirmed for opportunistic and pathogenic species from the genera *Malassezia*. Once established, biofilms are difficult to be fully removed from the pipe system, which on the long-term leads to altered taste and odour of water, production of allergenic or irritating compounds, and mycotoxins with an effect on human health [31].

The current study revealed that patients who do not have any type of contact with animals were more frequently exposed to *Malassezia* infection and *M. furfur* was the predominant species which explain

that the source of infection not restricted to birds or dogs. Patients who have contact with dogs only were equally exposed to *M. furfur*, and *M. pachydermatis* which reflect that dogs not only the source of infection but also other factors such as this may be attributed to several factors includes but not limited to, humidity and high temperature, hyperhidrosis, familial susceptibility, and immunosuppression [32]. A range of skin microenvironmental factors, such as the bacterial microbiota present, pH, salts, immune responses, cutaneous biochemistry, and physiology, may play a role in adherence and growth of *Malassezia* species, favouring distinct genotypes depending on the geographical area and/or the skin sites [33]. No significant correlation was reported between contact with animals and *Malassezia* spp — infection which comes in line with Al-Ezzy from Iraq [21].

In conclusion, *M. furfur*, *M. pachydermatis* and *M. globosa* represent the most common *Malassezia* spp. causing PV. Using of PCR is very critical to confirm the diagnosis of *Malassezia* spp. *Malassezia* infection inversely correlated with age and positively correlated with females gender and education. The residency in a rural area and middle economic status increase the possibility of infection. Infection was not affected by the source of water; job and contact with dogs and birds.

References

- Inamadar AC, Palit A. The genus *Malassezia* and human disease. *Indian Journal of Dermatology, Venereology, and Leprology*. 2003; 69(4):265.
- Gaitanis G, Magiatis P, Hantschke M, Bassukas ID, Velegriaki A. The *Malassezia* genus in the skin and systemic diseases. *Clinical microbiology reviews*. 2012; 25(1):106-41. <https://doi.org/10.1128/CMR.00021-11>
- Varada S, Dabade T, Loo DS. Uncommon presentations of tinea versicolor. *Dermatology practical & conceptual*. 2014; 4(3):93. <https://doi.org/10.5826/dpc.0403a21>
- Dawson T. The human skin microbiome: cause or effect? The role of *Malassezia* in human skin health and disease. *Medical Mycology Oxford Univ Press Great Clarendon St, Oxford OX2 6dp, England*, 2018.
- Rai M, Wankhade S. Tinea versicolor—an epidemiology. *J Microbial Biochem Technol*. 2009; 1(1):51-6. <https://doi.org/10.4172/1948-5948.1000010>
- Khosravi A, Eidi S, Katiraei F, Ziglari T, Bayat M, Nissiani M. Identification of different *Malassezia* species isolated from patients with *Malassezia* infections. *World Journal of Zoology*. 2009; 4(2):85-9.
- ElShabrawy WO, Saady N, Sallam M. Molecular and Phenotypic Identification and Speciation of *Malassezia* Yeasts Isolated from Egyptian Patients with Pityriasis Versicolor. *Journal of clinical and diagnostic research: JCDR*. 2017; 11(8):DC12. <https://doi.org/10.7860/JCDR/2017/27747.10416>
- Shokohi T, Afshar P, Barzgar A. Distribution of *Malassezia* species in patients with pityriasis versicolor in Northern Iran. *Indian journal of medical microbiology*. 2009; 27(4):321.

<https://doi.org/10.4103/0255-0857.55445>

9. Tu WT, Chin SY, Chou CL, Hsu CY, Chen YT, Liu D, et al. Utility of Gram staining for diagnosis of Malassezia folliculitis. *The Journal of dermatology*. 2018; 45(2):228-31. <https://doi.org/10.1111/1346-8138.14120>
10. Karhoot J, Noaimi A, Ahmad W. Isolation and identification of Malassezia species in patients with pityriasis versicolor. *The Iraqi Postgraduate Medical Journal*. 2012; 11(6):724-30.
11. Darling MJ, Lambiasi MC, Young RJ. Tinea versicolor mimicking pityriasis rubra pilaris. *CUTIS-NEW YORK*. 2005; 75(5):265.
12. Gouda A. Malassezia species isolated from lesional and non lesional skin in patients with pityriasis versicolor: M. Sc. thesis. College of Medicine, University of Ain Shams, 2008.
13. Petti CA, Carroll KC. Procedures for the Storage of Microorganisms. In *Manual of Clinical Microbiology*, 10th Edition, 2011:124-131. <https://doi.org/10.1128/9781555816728.ch9>
14. Qiagen. QIAamp® DNA Mini and Blood Mini Handbook. Germany Qiagen, 2016.
15. Mirhendi H, Makimura K, Khoramzadeh M, Yamaguchi H. A one-enzyme PCR-RFLP assay for identification of six medically important Candida species. *Nippon Ishinkin Gakkai Zasshi*. 2006; 47(3):225-9. <https://doi.org/10.3314/ijmm.47.225>
16. Jang S-J, Lim S-H, Ko J-H, Oh B-H, Kim S-M, Song Y-C, et al. The investigation on the distribution of Malassezia yeasts on the normal Korean skin by 26S rDNA PCR-RFLP. *Annals of dermatology*. 2009; 21(1):18-26. <https://doi.org/10.5021/ad.2009.21.1.18>
17. Shah A, Koticha A, Ubale M, Wanjare S, Mehta P, Khopkar U. Identification and speciation of Malassezia in patients clinically suspected of having pityriasis versicolor. *Indian journal of dermatology*. 2013; 58(3):239. <https://doi.org/10.4103/0019-5154.110841>
18. El-Hadidy G, Gomaa N, AboBakar R, Metwally L. Direct molecular identification of Malassezia species from skin scales of patients with seborrheic dermatitis by nested terminal fragment length polymorphism analysis Egypt. *J Med Microbiol*. 2007; 16:437-44.
19. Talaee R, Katirae F, Ghaderi M, Erami M, Alavi AK, Nazeri M. Molecular identification and prevalence of Malassezia species in pityriasis versicolor patients from Kashan, Iran. *Jundishapur journal of microbiology*. 2014; 7(8). <https://doi.org/10.5812/ijm.11561>
20. Al-Ammari AMM. Relation of Malassezia species with some skin diseases. Baghdad University: Baghdad University, 2012.
21. Al-Ezzy AIA, Jameel GH, Minnat TR. Isolation of Malassezia Furfur and Evaluation of Ivermectin and Cal-vatia Craniiformis as A Novel Antifungal Agents for Pityriasis Versicolor with Special Refer to Risk Factors in Iraqi Patients. *International Journal of Current Pharmaceutical Review and Research*. 2017; 8(4):311-9.
22. Chaudhary R, Singh S, Banerjee T, Tilak R. Prevalence of different Malassezia species in pityriasis versicolor in central India. *Indian Journal of Dermatology, Venereology, and Leprology*. 2010; 76(2):159. <https://doi.org/10.4103/0378-6323.60566>
23. Archana BR, Beena PM, Kumar S. Study of the distribution of malassezia species in patients with pityriasis versicolor in Kolar Region, Karnataka. *Indian journal of dermatology*. 2015; 60(3):321. <https://doi.org/10.4103/0019-5154.156436>
24. Hasan A-RS, Abass AA, Khudier MA-K. Clinical and Fungal Study of Pityriasis Versicolor Infection among Patients with Skin Mycoses in Baquba. *Iraqi Journal of Community Medicine*. 2009; 22(1):30-3.
25. Thayikkannu AB, Kindo AJ, Veeraraghavan M. Malassezia— Can it be Ignored? *Indian journal of dermatology*. 2015; 60(4):332. <https://doi.org/10.4103/0019-5154.160475>
26. Devendrappa K, Javed MW. Clinical profile of patients with tinea versicolor. *International Journal of Research in Dermatology*. 2018; 4(1):33-7. <https://doi.org/10.18203/issn.2455-4529.IntJResDermatol20180017>
27. Abdul-Hussein AA. Clinical and Pigmentary Variation of Pityriasis Versicolor in Al-Muthana Government's Patients. *Medical Journal of Babylon*. 2010; 7(3-4):383-8.
28. Akaza N, Akamatsu H, Takeoka S, Sasaki Y, Mizutani H, Nakata S, et al. Malassezia globosa tends to grow actively in summer conditions more than other cutaneous Malassezia species. *The Journal of dermatology*. 2012; 39(7):613-6. <https://doi.org/10.1111/j.1346-8138.2011.01477.x>
29. Laschinsky T. Extreme Dandruff and Seborrheic Dermatitis Hair Conditions. 2010.
30. Sharma M, Sharma R. Profile of dermatophytic and other fungal infections in Jaipur. *Indian journal of microbiology*. 2012; 52(2):270-4. <https://doi.org/10.1007/s12088-011-0217-z>
31. Babič MN, Gunde-Cimerman N, Vargha M, Tischner Z, Magyar D, Verissimo C, et al. Fungal contaminants in drinking water regulation? a tale of ecology, exposure, purification and clinical relevance. *International journal of environmental research and public health*. 2017; 14(6):636. <https://doi.org/10.3390/ijerph14060636>
32. Gupta A, Foley K. Antifungal Treatment for Pityriasis Versicolor. *Journal of Fungi* 2015; 1: 13-29. <https://doi.org/10.3390/jof1010013>
33. Cafarchia C, Gasser R, Figueredo L, Latrofa M, Otranto D. Advances in the identification of Malassezia. *Molecular and cellular probes Molecular and cellular probes*. 2011; 25: 1-7. <https://doi.org/10.1016/j.mcp.2010.12.003>