

# The Correlation between Hemoglobin Concentration during Pregnancy with the Maternal and Neonatal Outcome

Sarma Nursani Lumbanraja<sup>1\*</sup>, Muhammad Rizki Yaznil<sup>1</sup>, Dewi Indah Sari Siregar<sup>2</sup>, Adriani Sakina<sup>3</sup>

<sup>1</sup>Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia;

<sup>2</sup>Department of Clinical Pathology, Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia; <sup>3</sup>Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia

## Abstract

**Citation:** Lumbanraja SN, Yaznil MR, Siregar DIS, Sakina A. The Correlation between Hemoglobin Concentration during Pregnancy with the Maternal and Neonatal Outcome. Open Access Maced J Med Sci. 2019 Feb 28; 7(4):594-598. <https://doi.org/10.3889/oamjms.2019.150>

**Keywords:** Hemoglobins concentration; Anemia during pregnancy; Maternal; Neonatal outcome

**\*Correspondence:** Sarma Nursani Lumbanraja, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia. E-mail: [sarmalumbanraja@yahoo.com](mailto:sarmalumbanraja@yahoo.com)

**Received:** 19-Nov-2018; **Revised:** 06-Feb-2019; **Accepted:** 07-Feb-2019; **Online first:** 27-Feb-2019

**Copyright:** © 2019 Sarma Nursani Lumbanraja, Muhammad Rizki Yaznil, Dewi Indah Sari Siregar, Adriani Sakina. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

**Funding:** This project was supported by Basic Research Award from the TALENTA Universitas Sumatera Utara, Indonesia

**Competing Interests:** The authors have declared that no competing interests exist

**BACKGROUND:** The prevalence of anaemia is higher among women, including pregnant women. The estimation was about 24.8% of the population in the world suffering anaemia. Anaemia during pregnancy is a big problem because it can contribute morbidity and mortality, either in mother or newborn. The impacts of anaemia during pregnancy included post-partum haemorrhage, low birth weight (LBW), preterm delivery, and low Appearance, Pulse, Grimace, Activity, Respiration (APGAR) score.

**AIM:** This study aimed to determine the correlation between haemoglobin concentration during pregnancy and the outcome of mothers and newborns.

**METHODS:** It was a cohort study that included 200 pregnant women in second or third trimester at antenatal care of Sundari General Hospital Outpatient Clinic on February until September 2018. The participants were interviewed using a questionnaire, and their blood was checked to measure haemoglobin concentration using portable Easy Touch Hemoglobinometer. In the next three until six months, the following investigation was conducted to assess the maternal and neonatal outcome.

**RESULTS:** The result of this study showed among the maternal outcome, only antepartum haemoglobin concentration had a statistically significant correlation with the haemoglobin concentration during pregnancy ( $p < 0.05$ ), meanwhile, among the neonatal outcome. LBW was the only factor that statistically significantly correlated to the haemoglobin concentration during pregnancy ( $p < 0.05$ ).

**CONCLUSION:** We can conclude that once anaemia occurs in pregnant women, then the women kept suffering from anaemia with its correlation was statistically significant.

## Introduction

In all over the world, anaemia has been a global health burden since it can affect anyone without considering age or gender group [1]. However, the prevalence of this nutritional disorder is higher among women, including pregnant women. About 24.8% of the population in the world suffers from anaemia [2], [3].

Anaemia during pregnancy has become a widespread nutritional disorder either in developing or even developed countries. According to the World Health Organization (WHO), the proportion of the population suffering from anaemia in pregnancy was

14% in developed countries and 51% in developing countries. WHO also estimated that among the population, the cases were the most frequently found in Africa and Southeast Asia. Indonesia is a part of Southeast Asia. Thus, anaemia in pregnant women is frequently found here. It is proven by the data from Basic Health Research Ministry of Health of the Republic of Indonesia in 2013 showing that the prevalence was about 37.1% [1], [2], [4].

Anaemia during pregnancy is defined as the haemoglobin criteria is less than 11 g/dL. Once pregnant women suffer from anaemia, iron deficiency becomes the most common cause that should be considered [3]. Other risk factors related to maternal anemia are unhealthy lifestyle, poor socio-economic

status, malnutrition, hemoglobinopathies, age (under 20 years or above 35 years), early marriage or teenage pregnancy, decreasing period of pregnancy interval, smoking or alcohol use, history of menstrual disorder or infection, and gemelli or multiple pregnancies [5].

Anaemia is a big problem especially when it occurs during pregnancy because it may contribute to morbidity and mortality, either in mothers or newborns. Anaemia (regardless the severity) accounts 12.8% maternal death which is the second leading cause of the death. In Indonesia, the national Maternal Mortality Rate (MMR) is still so high with 307/100.000 live births. Anaemia during pregnancy becomes an indirect cause of postpartum haemorrhage and results in maternal mortality in the latter [4], [5]. Meanwhile, the national Neonatal Mortality Rate is much higher with 987/100.000 live births. Moreover, the fetal or neonatal complication can include prematurity, low birth weight, and low APGAR Score. All complications were significantly ended with maternal and neonatal mortality [4], [6], [7].

Although anaemia during pregnancy may lead to many adverse effects, actually anaemia can be the most preventable cause of maternal and neonatal mortality. In the upcoming years, anaemia in pregnant women should be eradicated to improve maternal and neonatal health status. Hence, this study was aimed to investigate the correlation between haemoglobin concentration during pregnancy and the outcome of mothers and newborns. In the future, hopefully, this study can be the reference to counsel pregnant women, especially in Indonesia, to be more aware of how threatening anaemia is and to decrease the prevalence of anaemia itself.

## Material and Methods

This study was conducted at antenatal care of Sundari General Hospital Outpatient Clinic. It was a prospective randomised study that included 200 pregnant women in the second or third trimester who met the inclusion and exclusion criteria.

This study took a period of 7 months — February until September 2018. After obtaining the consent, we interviewed the pregnant women who consumed Fe tablets, consist of ferrous sulfate 200 mg which contains 60 mg elemental iron for 90 days after the first trimester using a questionnaire to know their characteristics including age, gestational age, education background, and occupation. Then, the haemoglobin measurement was conducted by using Easy Touch portable hemoglobinometer. The data was collected during their antenatal care in the second or third trimester.

In the next three until six months, following investigation was conducted by asking the pregnant women to come again before giving birth. Then, we assessed the second haemoglobin measurement during their antepartum by the same portable measurement.

A few weeks later, the data about the outcome of maternal and neonatal was collected from the medical record if the subjects are giving birth in Sundari General Hospital. Unless they did not give birth in the same hospital, follow up was done by the phone. The maternal outcome includes the estimation of haemoglobin antepartum, bleeding volume while giving birth, and the initial breastfeeding. The neonatal outcome includes the data whether the newborn alive or not. Among the alive and healthy newborn, other data was collected, including preterm birth, low birth weight, and APGAR score. In this study, low birth weight defined as the infant birth weight which less than 2500 gram and preterm birth was considered as gestational age under 37 weeks. Because of our limitation, APGAR score was measured by only asking whether the newborn cried spontaneously or not. The newborn crying spontaneously was considered as good APGAR Score, but if they did not, it probably showed poor APGAR Score

### Statistical Analysis

All data collected and recorded using the Statistic Product and Service Solution (SPSS) program 21<sup>st</sup> version. Values were expressed with the mean  $\pm$  SD or percentage as appropriate. The analysis of the correlation between haemoglobin concentration during pregnancy and the outcome of mother and newborn were computed statistically using Chi-Square Test. A difference was considered significant at the p-value  $< 0.05$ .

## Results

There were 200 pregnant women involved with their newborns that evaluated in this study. There was found no maternal mortality. In this study, the mean of haemoglobin (Hb) concentration was  $10.73 \pm 2$  g/dl, the median was 10.6 g/dl, the maximum was 15.8 g/dl, and the minimum was 7.8 g/dl. The mean birth weight was  $3015 \pm 584$  gram.

The subject characteristics were shown in Table 1. The mean age of all participants was  $29 \pm 5.5$  years. For all characteristics, except birth method, either anaemia or normal group, had the similar result of majority variable. The most common age was 20 – 35 years. The majority of participants were in the third trimester of gestational age. Most of the participants were housewives. The majority education background

was senior high school. Most of the participants gave birth in a hospital. Otherwise, for the birth method, the anaemia group had different result compared to the normal group. The majority birth method in anaemia group was spontaneous birth. Meanwhile, cesarean birth was most commonly conducted in the normal group. By using Chi-Square Test, there were no characteristics which had a significant correlation with the hemoglobin during pregnancy (p-value > 0.05).

**Table 1: Demographic of subject characteristics**

Characteristics	Haemoglobin concentration during pregnancy		Total	P value
	Anemia (52.5%)	Normal (47.5%)		
Age				
	Mean 29 ± 5.5 years			0.218
< 20 years	0 (0%)	2 (2.1%)	2 (1%)	
20 – 35 years	88 (83.8%)	79 (83.2%)	167 (83.5%)	
> 35 years	17 (16.2%)	14 (14.7%)	31 (15.5%)	
Gestational Age				
2 <sup>nd</sup> Trimester	28 (26.7%)	19 (20%)	47 (23.5%)	0.267
3 <sup>rd</sup> Trimester	77 (73.3%)	76 (80%)	153 (76.5%)	
Occupation				
Housewife	77 (73.3%)	72 (75.8%)	149 (74.5%)	0.974
Civil servant	9 (8.6%)	7 (7.4%)	16 (8%)	
Private servant	9 (8.6%)	7 (7.4%)	16 (8%)	
Entrepreneur	10 (9.5%)	9 (9.5%)	19 (9.5%)	
Education background				
Elementary school	4 (3.8%)	2 (2.1%)	6 (3%)	0.453
Junior high school	13 (12.4%)	14 (14.7%)	27 (13.5%)	
Senior high school	59 (56.2%)	57 (60%)	116 (58%)	
Diploma	6 (5.7%)	9 (9.5%)	15 (7.5%)	
Undergraduate	23 (21.9%)	13 (13.7%)	36 (18%)	
Mode of Delivery				
Normal vaginal	76 (72.4%)	36 (37.9%)	112 (56%)	0.121
Cesarean	29 (27.6%)	59 (62.1%)	88(44%)	
Birth location				
Hospital	56 (53.3%)	59 (62.1%)	115 (57.5%)	0.210
Midwifery unit	49 (46.7%)	36 (37.9%)	85 (42.5%)	

Chi-Square Test. p-value < 0.05 were considered as statistically significant.

The indication of cesarean birth can be seen in Table 2. The most common indication was the previous cesarean delivery. The other indications related to pregnancy problems, such as obstructed labour, placenta previa, abnormal fetal presentation, and cephalopelvic disproportion.

**Table 2: The indications of cesarean birth**

Factors	Anaemia	Non-anaemia
Previous cesarean birth	21 (72.4%)	20 (58.8%)
Obstructed labor	3 (10.3%)	5 (14.7%)
Placenta previa	2 (6.9%)	1 (2.9%)
Cephalopelvic disproportion	1 (3.4%)	2 (5.8%)
Abnormal fetal presentation	2 (6.9%)	5 (14.7%)
Others	0 (0%)	1 (2.9%)

Table 3 showed how maternal outcome related to haemoglobin concentration during pregnancy. Among the three variables, only antepartum haemoglobin concentration that significantly associated with haemoglobin concentration during pregnancy (p < 0.05).

**Table 3: The correlation between haemoglobin concentration during pregnancy and maternal outcome**

Maternal outcome	Haemoglobin concentration during pregnancy		P value
	Anaemia	Normal	
Antepartum hemoglobin concentration	Anemia 89 (84.8%) Non-anemia 16 (15.2%)	57 (60%) 38 (40%)	0.000*
Postpartum hemorrhage (PPH)	< 500 cc 85 (81%) ≥ 500 cc 20 (19%)	78 (82.1%) 17 (17.9%)	0.834
Initial breastfeeding	Yes 27 (25.7%) No 74 (74.3%)	27 (28.4%) 68 (71.6%)	0.791

Chi-Square Test \*p-value < 0.05 were considered as statistically significant.

It means that the pregnant women in either second or third trimester kept suffering from anaemia until the antepartum period.

Table 4 showed that among all participants, unfortunately, stillbirth (3.8%) was found. All stillbirth cases occurred in the anaemia group. There was no significant correlation between haemoglobin concentration during pregnancy and neonatal condition (p > 0.05).

**Table 4: The correlation between hemoglobin concentration during pregnancy and neonatal condition**

Neonatal condition	Hemoglobin concentration during pregnancy		P value
	Anemia	Normal	
Alive newborn	101 (96.2%)	95 (100%)	0.157
Stillbirth	4 (3.8%)	0 (0%)	

Chi Square Test.

To analyse the correlation between haemoglobin concentration during pregnancy and fetal outcome, participants with stillbirth were excluded. We found four cases of stillbirth. Thus, there were only 196 participants involved in this analysis. The result showed in Table 5. Among all variables of neonatal outcome, low birth weight was the only outcome that significantly related to the haemoglobin concentration during pregnancy (p < 0.05).

**Table 5: The Correlation between haemoglobin concentration during pregnancy and fetal outcome**

Neonatal outcome	Haemoglobin during pregnancy		P value
	Anaemia	Non-anaemia	
Spontaneous cry	Yes 92(96.8%) No 9(3.2%)	86 (90.5%) 9 (9.5%)	0.892
Preterm delivery	Yes 4 (4%) No 97 (96%)	10 (10.5%) 85 (89.5%)	0.074
Low birth weight	Yes (< 2500 gram) 4 (81%) No (≥ 2500 gram) 97 (19%)	11 (82.1%) 84 (17.9%)	0.045*

Chi-Square Test \*p-value < 0.05 were considered as statistically significant.

## Discussion

Out of 200 pregnant women in this study, more than half (52.5%) of them were suffering from anaemia. It is relevant to the previous study which stated that half population of pregnant women in the world affected by anemia. Meanwhile, the proportion of anemic pregnant women varies in each countries, such as 58% in China, 50% in Southeast Asia, and 40% in Istanbul [8].

Based on Chi-Square Test, this study showed that no significant correlation between hemoglobin concentration during pregnancy with any characteristic variables, included maternal age (p-value = 0.218), gestational age (p-value = 0.267), education background (p-value = 0.974), occupation (p-value = 0.453), mode of delivery (p-value = 0.121) and place of birth (p-value = 0.210). It was relevant to

the previous study conducted by Vural *et al.*, showing that there was a statistically significant correlation between anaemia prevalence and mode of delivery. Otherwise, in Vural *et al.*, study, the correlation between maternal age and anaemia prevalence was statistically significant [8].

This study found that the mean of haemoglobin concentration was about  $10.73 \pm 2$  g/dl. Based on both the World Health Organization (WHO) and the Center for Disease Controls and Preventions (CDC), the mean is defined as anaemia since the haemoglobin concentration less than 11 g/dl. [9]. Although it was categorised as anaemia according to criteria of WHO and CDC, the concentration of 10 g/dl in the mid-trimester of gestational age seems to reflect the adequate expansion of plasma volume [10].

Although more than half of the population were anaemic, there was found no maternal mortality in this study. It was suggested that the haemoglobin concentration still able to compensate for optimal plasma volume expansion. It was relevant to the previous study stated that the cutoff of extremely low haemoglobin concentration was less than 6.5 g/dl. This condition with other factors can contribute to maternal mortality. Even, the other cutoff with 8.9 g/dl associated with twice risk of maternal mortality [11].

In this study, out of 115 anaemic pregnant women, 20 participants (19%) had bleeding after birth  $\geq 500$  cc without significant correlation between haemoglobin level during pregnancy and the bleeding volume. The previous study of Frass had similar result showing 29.1% of anaemic pregnant women developed postpartum haemorrhage during cesarean delivery because of the uterine atony [12]. Even, the study of Kayle *et al.* showed that there was a strongly significant correlation between moderate-to-severe anaemia and blood loss severity [13]. Despite the widespread postpartum haemorrhage in all over the world, unfortunately, there is still a lack of data in the literature about contributing factors of it, especially in developing countries where many PPH and maternal death occur. David study stated that weak uterine muscular strength and lower resistance to infectious disease possibly occur due to severe anaemia. Meanwhile, the higher risk of PPH experience related to severity anaemia still needs further studies [13], [14].

This study investigated whether there is a correlation between initial breastfeeding and the haemoglobin concentration during pregnancy. Our study showed that most of the initial breastfeeding failure found in the anaemia group (74.3%). Despite the fact, the correlation was statistically not significant ( $p$ -value = 0.791). Another complication of anaemia during pregnancy that still not familiar is breastfeeding failure. However, an article review mentioned breastfeeding failure becoming the impact of anaemia during pregnancy, after puerperal sepsis and sub-involution [15].

Investigation of neonatal outcome was done by analysing three factors, including spontaneous crying to assess APGAR Score, preterm delivery, and low birth weight. Among the factors, only low birth weight had a significant correlation with the haemoglobin level during pregnancy ( $p = 0.045$ ).

Birth weight is a good indicator to evaluate whether the mother supports the fetus adequately or not. Besides, it is the only determinant factor of newborn mortality in the first year of life. Birth weight less than 2500 gram is defined as low birth weight that is most commonly caused by anaemia during pregnancy. Previous meta-analysis literature showed that there was a significant correlation between anaemia during pregnancy and low birth weight in the 3<sup>rd</sup> trimester. Otherwise, the same study showed no significant correlation between both variables in 1<sup>st</sup> and 2<sup>nd</sup> trimester [9], [16]. Haemoglobin concentration of less than 10.5 g/dl was reported to increase the sevenfold risk of low birth weight [17]. Abnormally a previous literature stated that several studies reported low birth weight in anaemic pregnant women, but high haemoglobin concentration in 1<sup>st</sup> and 3<sup>rd</sup> trimester also correlated with the risk of low birth weight due to similarly poor plasma volume expansion [10], [17].

Besides low birth weight, preterm delivery and low APGAR score were the other impacts of anaemia during pregnancy in neonatal outcome. This study showed that there is no significant correlation between haemoglobin concentration during pregnancy with any factor ( $p > 0.05$ ). It seems relevant with the previous study conducted in Moshi Municipality showing a similar result about no correlation between anaemia and low birth weight and preterm delivery. However, a different study showed that maternal anaemia and preterm delivery had significant correlation statistically. Even,  $Hb < 10.5$  g/dl can increase fivefold risk of preterm delivery [17], [18]. Our study can give more information about how the impact of haemoglobin concentration on the outcome of maternal and neonatal. However, our study had a limitation that did not bring any information about the dietary pattern of the pregnant woman, especially during pregnancy. This study also did not give any intervention for the anaemia correction before the childbirth.

In conclusion, from this study, we can conclude that once anaemia occurs in pregnant women, then the women kept suffering from anaemia with its correlation was statistically significant. Several studies showed that there was a significant correlation between anaemia during pregnancy and postpartum haemorrhage. However, this study found a different result with no significant correlation between haemoglobin concentration with PPH and initial breastfeeding. Therefore, further studies are needed to investigate more about the maternal and neonatal outcome due to anaemia during pregnancy. Moreover, this kind of study is still limited developed in Indonesia.

## Reference

1. Vanamala VG, Rachel A, Pakyanadhan S. Incidence and outcome of anemia in pregnant women: a study in a tertiary care centre. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*. 2018; 7(2):462-6. <https://doi.org/10.18203/2320-1770.ijrcog20180155>
2. Noronha JA, Al Khasawneh E, Seshan V, Ramasubramaniam S, Raman S. Anemia in pregnancy-consequences and challenges: a review of literature. *Journal of South Asian Federation of Obstetrics and Gynecology*. 2012; 4(1):64-70. <https://doi.org/10.5005/jp-journals-10006-1177>
3. Maka SS, Tondare SB, Tondare MB. Study of impact of anemia on pregnancy. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*. 2017; 6(11):4847-50. <https://doi.org/10.18203/2320-1770.ijrcog20174692>
4. Lestari S, Fujiati II, Keumalasari D, Daulay M, Martina SJ, Syarifah S. The prevalence of anemia in pregnant women and its associated risk factors in North Sumatera, Indonesia. *IOP Conference Series: Earth and Environmental Science*. 2018; 125(1): 012195). <https://doi.org/10.1088/1755-1315/125/1/012195>
5. Kumar KJ, Asha N, Murthy DS, Sujatha MS, Manjunath VG. Maternal anemia in various trimesters and its effect on newborn weight and maturity: an observational study. *International journal of preventive medicine*. 2013; 4(2):193-9. PMID:23543625  
PMCID:PMC3604852
6. Abu-Ouf NM, Jan MM. The impact of maternal iron deficiency and iron deficiency anemia on child's health. *Saudi medical journal*. 2015; 36(2):146-9. <https://doi.org/10.15537/smj.2015.2.10289>  
PMid:25719576 PMCID:PMC4375689
7. Sukrat B, Wilasrusmee C, Siribumrungwong B, McEvoy M, Okascharoen C, Attia J, Thakkinstian A. Hemoglobin concentration and pregnancy outcomes: a systematic review and meta-analysis. *BioMed research international*. 2013; 2013.
8. Sharma JB, Shankar M. Anemia in pregnancy. *JIMSA*. 2010; 23(4):253-60.
9. Steer P, Alam MA, Wadsworth J, Welch A. Relation between maternal haemoglobin concentration and birth weight in different ethnic groups. *BmJ*. 1995; 310(6978):489-91. <https://doi.org/10.1136/bmj.310.6978.489> PMID:7888886  
PMCID:PMC2548871
10. Allen LH. Anemia and iron deficiency: effects on pregnancy outcome-. *The American journal of clinical nutrition*. 2000; 71(5):1280S-4S. <https://doi.org/10.1093/ajcn/71.5.1280s>  
PMid:10799402
11. Vural T, Toz E, Ozcan A, Biler A, Ileri A, Inan AH. Can anemia predict perinatal outcomes in different stages of pregnancy? *Pakistan journal of medical sciences*. 2016; 32(6):1354-1359. <https://doi.org/10.12669/pjms.326.11199> PMID:28083025  
PMCID:PMC5216281
12. Frass KA. Postpartum hemorrhage is related to the hemoglobin levels at labor: Observational study. *Alexandria Journal of Medicine*. 2015; 51(4):333-7. <https://doi.org/10.1016/j.ajme.2014.12.002>
13. Kavle JA, Stoltzfus RJ, Witter F, Tielsch JM, Khalfan SS, Caulfield LE. Association between anaemia during pregnancy and blood loss at and after delivery among women with vaginal births in Pemba Island, Zanzibar, Tanzania. *Journal of health, population, and nutrition*. 2008; 26(2):232-240. PMID:18686556  
PMCID:PMC2740668
14. Rush D. Nutrition and maternal mortality in the developing world-. *The American journal of clinical nutrition*. 2000; 72(1):212S-40S. <https://doi.org/10.1093/ajcn/72.1.212S>  
PMid:10871588
15. Chowdhury S, Rahman M, Moniruddin AB. Anemia in pregnancy. *Medicine Today*. 2014; 26(1):49-52. <https://doi.org/10.3329/medtoday.v26i1.21314>
16. Rahmati S, Delpishe A, Azami M, Ahmadi MR, Sayehmiri K. Maternal Anemia during pregnancy and infant low birth weight: A systematic review and Meta-analysis. *International Journal of Reproductive BioMedicine*. 2017; 15(3):125-134. <https://doi.org/10.29252/ijrm.15.3.125> PMID:28580444  
PMCID:PMC5447828
17. Scholl TO, Reilly T. Anemia, iron and pregnancy outcome. *The Journal of nutrition*. 2000; 130(2):443S-7S. <https://doi.org/10.1093/jn/130.2.443S> PMID:10721924
18. Stephen G, Mgongo M, Hussein Hashim T, Katanga J, Stray-Pedersen B, Msuya SE. Anaemia in Pregnancy: Prevalence, Risk Factors, and Adverse Perinatal Outcomes in Northern Tanzania. *Anemia*. 2018:1-9. <https://doi.org/10.1155/2018/1846280>  
PMid:29854446 PMCID:PMC5954959