

# Hepatoprotective Activity and Nephroprotective Activity of Peel Extract from Three Varieties of the Passion Fruit (*Passiflora Sp.*) in the Albino Rat

Nerdy Nerdy<sup>1\*</sup>, Kiking Ritarwan<sup>2</sup>

<sup>1</sup>Department of Pharmacy, Academy of Pharmacy Yayasan Tenaga Pembangunan Arjuna, Pintubosi, Laguboti, Toba Samosir, Sumatera Utara, 22381, Indonesia; <sup>2</sup>Department of Neurology, Faculty of Medicine, University of Sumatera Utara, Padang Bulan, Medan Baru, Medan, Sumatera Utara, 20155, Indonesia

## Abstract

**Citation:** Nerdy N, Ritarwan K. Hepatoprotective Activity and Nephroprotective Activity of Peel Extract from Three Varieties of the Passion Fruit (*Passiflora Sp.*) in the Albino Rat. Open Access Maced J Med Sci. 2019 Feb 28; 7(4):536-542. <https://doi.org/10.3889/oamjms.2019.153>

**Keywords:** Hepatoprotective; Nephroprotective; Passion Fruit; Biochemical Analysis

**\*Correspondence:** Nerdy Nerdy, Department of Pharmacy, Academy of Pharmacy Yayasan Tenaga Pembangunan Arjuna, Pintubosi, Laguboti, Toba Samosir, Sumatera Utara, 22381, Indonesia. E-mail: nerdy190690@gmail.com

**Received:** 22-Nov-2018; **Revised:** 30-Jan-2019; **Accepted:** 01-Feb-2019; **Online first:** 27-Feb-2019

**Copyright:** © 2019 Nerdy Nerdy, Kiking Ritarwan. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

**Funding:** This research did not receive any financial support

**Competing Interests:** The authors have declared that no competing interests exist

**BACKGROUND:** The Passion Fruit (*Passiflora sp.*) that grows in the Indonesian region generally has three varieties, namely purple passion fruit (*Passiflora edulis Sims.*), red passion fruit (*Passiflora ligularis Juss.*), and yellow passion fruit (*Passiflora verrucifera Lindl.*). The passion fruit peel is an economic waste that has not been utilised optimally, but has many efficacious phytochemical contents.

**AIM:** The objectives of this research are to examine scientifically hepatoprotective activity (with paracetamol-induced hepatotoxic) and nephroprotective activity (with gentamicin-induced nephrotoxic) from three varieties of the passion fruit (purple passion fruit peel extract, red passion fruit peel extract and yellow passion fruit peel extract) in the albino rat (*Rattus norvegicus*).

**METHODS:** Three varieties of passion fruit peels were extracted by maceration method. The experimental animals used were the albino rat (*Rattus norvegicus*). Hepatoprotective activity was done by the liver biochemical (alanine transaminase and aspartate transaminase) analysis with paracetamol (hepatotoxic compound) induced after 10 days of treatment with extract. Nephroprotective activity was done by the kidney biochemical (urea and creatinine) analysis with gentamicin (nephrotoxic compound) induced after 10 days of treatment with extract.

**RESULTS:** The hepatoprotective activity for positive control was similar to the 250 mg of purple passion fruit peel extract per kg of body weight, 250 mg of red passion fruit peel extract per kg of body weight, and 500 mg of yellow passion fruit peel extract per kg of body weight. The nephroprotective activity for positive control (50 mg of silymarin per kg of body weight) was similar to the 250 mg of purple passion fruit peel extract per kg of body weight, 500 mg of red passion fruit peel extract per kg of body weight, and 500 mg of yellow passion fruit peel extract per kg of body weight.

**CONCLUSIONS:** The extracts were shown hepatoprotective activity and nephroprotective activity with a dose-dependent activity. The hepatoprotective activity and nephroprotective activity of purple passion fruit peel extract were the best compared to red passion fruit peel extract and yellow passion fruit peel extract.

## Introduction

The liver is an important organ in the body that has an important function, which is: bile secretion, bilirubin metabolism, nutritional metabolism, producing immune agents to control infections, and metabolism of foreign molecules (exogenous chemical and endogenous chemical). The liver is the metabolic centre in the body, especially those given orally. Drug

metabolism in the liver occurs in microsome cells through an enzyme system that is very complex through two phases. The first phase includes oxidation reactions, reduction reactions, and hydrolysis reactions. The second phase includes conjugation reactions. Liver damage can be caused by a viral of hepatitis or cirrhosis; drugs can cause toxic effects on the liver resulting in liver damage [1].

The kidney is an important organ that has an important role in the body for blood filtration to excrete

waste products, balance electrolytes in the body, control blood pressure, stimulate the production of red blood cells, and regulate the balance of water and metabolites in the body and maintain acid-base balance in the blood. Drugs and metabolic results of drugs that are in the blood and are no longer used in the body will be excreted by the kidneys through urine. Urine will leave the kidneys to the urinary tract to be removed from the body. Most drugs are excreted by the kidneys, so the use of drugs that exceed the therapeutic dosage can damage the kidneys [2].

Paracetamol is a drug with analgesic and antipyretic effects that is widely used by the wider community. Paracetamol is an over the counter class drug that can be traded freely without supervision, but this drug causes liver damage in the use of high doses. Paracetamol is predicted to be a major factor in the cause of acute liver damage [3], [4]. Gentamicin is an antibiotic derived from aminoglycosides, which is generally an option in the treatment of infections that are needed immediately if a condition that can endanger life is found. Gentamicin is a broad-spectrum antibiotic that has bactericidal activity against gram-positive bacteria and gram-negative bacteria. Gentamicin can cause kidney damage because it rapidly induces extensive renal cortical necrosis with renal dysfunction [5], [6].

The passion fruit is a plant that originates from Brazil and spreads to all countries of the world. The passion fruit can grow in subtropical regions or highland in tropical regions [7]. The passion fruit can be found in several regions in Indonesia; one region in Indonesia that can cultivate the passion fruit is North Sumatera region (Karo district, Simalungun district, Dairi district, and North Tapanuli district). The passion fruit has many varieties that cultivate in all countries of the world, but the passion fruit that grows and develops in Indonesia are three varieties, namely: purple passion fruit, red passion fruit, and yellow passion fruit [8], [9]. Figure 1 shows the physical appearance of purple passion fruit, red passion fruit, and yellow passion fruit.



Figure 1: Physical appearance of purple passion fruit, red passion fruit, and yellow passion fruit

The passion fruit is generally consumed on the inside of the fruit, because the inside of the fruit is the edible portion, and the skin is a waste that has not been utilised optimally [10]. The passion fruit peel has many different phytochemical contents, including

alkaloids, flavonoids, steroids, triterpenoids, saponins, tannins, glycosides, and phenolic [11], [12], [13]. The passion fruit peel has also been tested for pharmacological activity as an antioxidant [12], [14], [15], [16], an antimicrobial [13], [15], [16], a neuroprotective [15], [17], a cardioprotective [18], a gastroprotective [19], an analgesic [20], an anti-inflammation [21], an antihypertriglyceridemic [21], and an antihyperglycemic [21], [22].

The hepatoprotective activity and nephroprotective activity of the passion fruit peel extract have never been reported in the research article. But with the abundance of phytochemical content found in the passion fruit peel extract, an extract that is likely to provide hepatoprotective activity [23] and nephroprotective activity [24].

This study aims to examine scientifically hepatoprotective activity (with paracetamol-induced hepatotoxic) and nephroprotective activity (with gentamicin-induced nephrotoxic) from three varieties of the passion fruit (*Passiflora sp.*) (purple passion fruit peel extract, red passion fruit peel extract and yellow passion fruit peel extract) in the albino rat (*Rattus norvegicus*).

## Methods

### Research

This research was conducted by experimental research. The independent variables were type of the extract (purple passion fruit peel extract, red passion fruit peel extract, and yellow passion fruit peel extract) and dose of the extract (100 mg extract per kg of body weight, 200 mg extract per kg of body weight, and 400 mg extract per kg of body weight). The dependent variables were level of the hepatotoxic or hepatoprotective biochemical marker (alanine transaminase and aspartate transaminase) in the serum and nephrotoxic or nephroprotective biochemical marker (urea and creatinine) in the serum. These research samples were purple passion fruit peel and yellow passion fruit peel obtained from the passion fruit farmer. The passion fruit growth in Gundaling, Berastagi, Karo, Sumatera Utara, 22152, Indonesia. The growth condition was a suitable condition for the passion fruit growth with 1,375 meters above sea level for the average altitude, 16°C to 26°C for the temperature, 60% to 100% for the relative humidity, 2100 mm to 3200 mm for the rainfall, 5 to 6 for the soil pH, 1009 HPa to 1015 HPa for the atmospheric pressure, 4 km/hr to 11 km/hr for the wind speed. Harvest of purple passion fruit, red passion fruit, and yellow passion fruit was done in 85<sup>th</sup> days to 90<sup>th</sup> days after the flowers bloom.

### Materials and Tools

The materials used in this research were purple passion fruit peel, red passion fruit peel, yellow passion fruit peel, ethanol (E-Merck), ethylene diamine tetraacetic acid (E-Merck), alanine transaminase activity assay kit (Sigma Aldrich), aspartate transaminase activity assay kit (Sigma Aldrich), urea assay kit (Sigma Aldrich), creatinine assay kit (Sigma Aldrich), sodium carboxymethyl cellulose (Tokyo Chemical Industry), and demineralized water (Brataco).

The tools used in this research were cutter (Kenko), drying cabinet (Alumex), blender (Miyako), balance (Mettler Toledo), analytical balance (Mettler Toledo), filter paper (Whatman), dropper (Iwaki), evaporating dish (Iwaki), maceration vessel (Iwaki), beaker glass (Iwaki), measuring glass (Iwaki), volumetric flask (Iwaki), test tube (Iwaki), hot plate stirrer (Thermo), water bath (Mettler), incubator (Mettler), rotary evaporator (Buchi), thermometer (Lutron), test sieve 10 (Retsch), stopwatch (Casio), spectrophotometer (Agilent).

### Samples Identification and Samples Extraction

Samples used in this research were collected and identified at Herbarium Medanense (MEDA), Faculty of Mathematics and Natural Sciences, University of Sumatera Utara, Padang Bulan, Medan Baru, Medan, Sumatera Utara, 20155, Indonesia. The extraction process was based on a modification of the maceration method from Irawan et al., 2018 and Zhang et al., 2018. The passion fruit peel is washed, cut, dried, and powdered. One kilogram of dried passion fruit peel powder was weighed, soaked with 10 L ethanol for 5 days (stirred every day), and filtered. The residue was squeezed, soaked with 5 L ethanol for 3 days (stirred every day), and filtered (the maceration procedure was repeated several times until clear extract was obtained). The aqueous extract was collected and evaporated with a rotary evaporator until a viscous extract was obtained [25], [26]. The extracts obtained were used for hepatoprotective activity test and nephroprotective activity test.

### Hepatoprotective Activity Test and Nephroprotective Activity Test

The method used for nephroprotective activity and hepatoprotective activity is a modification of the method from Abel-Hady et al., 2018; Fatima and Sultana, 2018; Tung et al., 2017; Okokon et al., 2017; and Thuwaini et al., 2016. The animals used were the Albino Rat (*Rattus norvegicus*) with Wistar strain which was 8 weeks old-12 weeks and weighed 250 g  $\pm$  10 g. The animals were kept for 5 days before the experiments. The animals were kept in standard conditions (12 hours light and 12 hours dark cycle),

with temperatures of 23°C  $\pm$  2°C, and with relative humidity 50%  $\pm$  10%. Experimental animals were divided into 12 groups for testing hepatoprotective activity and 12 groups for testing nephroprotective activity with each group consisting of 10 animals. Table 1 shows the treatment given to each group of experimental animals for 10 days and after 10 days [27], [28], [29], [30], [31].

**Table 1: Treatment given to each group of experimental animals for 10 days and after 10 days**

Number	Given	Group											
		1	2	3	4	5	6	7	8	9	10	11	12
1	Protection with Standard Protector for 10 Days (mg of Curcumin or mg of Silymarin per kg of body weight) by Oral Purple Passion Fruit Peel Extract for 10 Days	-	-	✓	-	-	-	-	-	-	-	-	-
2	(mg of Purple Passion Fruit Peel Extract per kg of body weight) by Oral Red Passion Fruit Peel Extract for 10 Days	-	-	-	✓	✓	✓	-	-	-	-	-	-
3	(mg of Red Passion Fruit Peel Extract per kg of body weight) by Oral Yellow Passion Fruit Peel Extract for 10 Days	-	-	-	-	-	-	✓	✓	✓	-	-	-
4	(mg of Yellow Passion Fruit Peel Extract per kg of body weight) by Oral Induction with Toxic Inducer after 10 Days	-	-	-	-	-	-	-	-	-	✓	✓	✓
5	(mg of Paracetamol or mg of Gentamicin per kg of body weight) by Intraperitoneal	-	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

All groups (group 1 to group 12) were given food and drink ad libitum. The protector used were 250 mg of curcumin per kg of body weight (for hepatoprotective activity) and 50 mg of silymarin per kg of body weight (for nephroprotective activity). The inducer used were 500 mg of paracetamol per kg of body weight (for hepatotoxic activity) and 125 mg of gentamicin per kg of body weight (for nephrotoxic activity). The samples used were purple passion fruit peel extract, red passion fruit peel extract and yellow passion fruit peel extract. Each extract was used with various concentrations (125 mg of extract per kg of body weight, 250 mg of extract per kg of body weight, and 500 mg of extract per kg of body weight). The animals from all groups were sacrificed under diethyl ether anaesthesia, and then the blood samples were collected after 24-hour induction with toxic inducer by cardiac puncture by using 21 gauge needles mounted on a 5 ml syringe into ethylene diamine tetraacetic acid.

The blood samples obtained were centrifuged at 3000 rotations per minute for 15 minutes. Serum obtained were used for the biochemical analysis with hepatoprotective activity parameters (alanine transaminase and aspartate transaminase) and nephroprotective activity parameters (urea and creatinine) [27], [28], [29], [30], [31]. The biochemical analysis of hepatoprotective activity parameters (alanine transaminase and aspartate transaminase) and the biochemical analysis of nephroprotective activity parameters (urea and creatinine) were carried out by the standard method from Sigma Aldrich.

## Results

Increasing the dose of administration will increase the hepatoprotective activity and the nephroprotective activity. This can be seen from the results that increase treatment dose will improve the liver function through a decrease in the alanine transaminase value and aspartate transaminase value and improve the kidney function through a decrease in the urea value and creatinine value so that it can be seen that purple passion fruit peel extract, red passion fruit peel extract, and yellow passion fruit peel extract have a dose-dependent hepatoprotective activity and dose-dependent nephroprotective activity. Table 2 shows the results of treatment of hepatoprotective activity test (alanine transaminase and aspartate transaminase) and treatment of nephroprotective activity test (urea and creatinine).

**Table 2: Results of treatment of hepatoprotective activity test (alanine transaminase and aspartate transaminase) and treatment of nephroprotective activity test (urea and creatinine)**

Number	Treatment	Alanine Transaminase (IU/L)	Aspartate Transaminase (IU/L)	Urea (mg/dL)	Creatinine (mg/dL)
1	Normal	27.45 ± 2.84 <sup>#†</sup>	33.19 ± 3.11 <sup>#†</sup>	13.12 ± 1.51 <sup>#†</sup>	0.441 ± 0.024 <sup>#†</sup>
2	Control	95.84 ± 9.87 <sup>**‡</sup>	116.17 ± 10.94 <sup>**‡</sup>	77.64 ± 9.12 <sup>**‡</sup>	7.419 ± 0.409 <sup>**‡</sup>
3	Positive	32.34 ± 3.41 <sup>*#</sup>	38.09 ± 3.61 <sup>*#</sup>	17.57 ± 2.02 <sup>*#</sup>	0.521 ± 0.029 <sup>*#</sup>
4	Purple Passion Fruit Peel Extract (mg of Purple Passion Fruit Peel Extract per kg of body weight)	40.97 ± 4.18 <sup>*#‡</sup>	54.07 ± 4.94 <sup>*#‡</sup>	24.14 ± 3.24 <sup>*#‡</sup>	0.559 ± 0.031 <sup>*#‡</sup>
5	250	32.59 ± 3.32 <sup>*#</sup>	38.33 ± 3.71 <sup>*#</sup>	17.67 ± 2.25 <sup>*#</sup>	0.518 ± 0.028 <sup>*#</sup>
6	500	30.84 ± 3.22 <sup>*#†</sup>	33.55 ± 3.09 <sup>*#†</sup>	13.19 ± 1.95 <sup>*#†</sup>	0.500 ± 0.026 <sup>*#†</sup>
7	Red Passion Fruit Peel Extract (mg of Red Passion Fruit Peel Extract per kg of body weight)	41.15 ± 4.19 <sup>*#‡</sup>	53.57 ± 4.91 <sup>*#‡</sup>	31.99 ± 4.08 <sup>*#‡</sup>	0.675 ± 0.042 <sup>*#‡</sup>
8	250	32.13 ± 3.30 <sup>*#</sup>	38.69 ± 3.76 <sup>*#</sup>	24.56 ± 3.13 <sup>*#‡</sup>	0.595 ± 0.036 <sup>*#‡</sup>
9	500	30.34 ± 3.18 <sup>*#†</sup>	33.99 ± 3.12 <sup>*#†</sup>	17.69 ± 2.52 <sup>*#</sup>	0.534 ± 0.032 <sup>*#</sup>
10	Yellow Passion Fruit Peel Extract (mg of Yellow Passion Fruit Peel Extract per kg of body weight)	66.21 ± 6.94 <sup>*#‡</sup>	75.04 ± 7.14 <sup>*#‡</sup>	32.24 ± 4.10 <sup>*#‡</sup>	0.671 ± 0.041 <sup>*#‡</sup>
11	250	43.75 ± 4.49 <sup>*#‡</sup>	49.74 ± 4.71 <sup>*#‡</sup>	24.22 ± 3.07 <sup>*#‡</sup>	0.591 ± 0.035 <sup>*#‡</sup>
12	500	32.94 ± 3.53 <sup>*#</sup>	37.88 ± 3.48 <sup>*#</sup>	18.01 ± 2.59 <sup>*#</sup>	0.530 ± 0.030 <sup>*#</sup>

Note: value represents as mean ± standard deviation; <sup>#</sup>p < 0.05, a significant difference compared with the normal control group (n = 10); <sup>†</sup>p < 0.05, a significant difference compared with the negative control group (n = 10); <sup>‡</sup>p < 0.05, a significant difference (lower) compared with the positive control group (n = 10) <sup>\*</sup>p < 0.05, a significant difference (higher) compared with the positive control group (n = 10).

The normal control group that was not given a protector and was not given an inducer showed normal liver biochemical value (alanine transaminase value < 35.0 IU/L and aspartate transaminase value < 45.0 IU/L) and normal kidney biochemical value (urea value < 24.0 mg/dL and creatinine value < 1.20 mg/dL). The negative control group that was not given a protector and was given an inducer showed significantly increased in liver biochemical value (alanine transaminase value > 35.0 IU/L and aspartate transaminase value > 45.0 IU/L) and significantly increased in kidney biochemical value (urea value > 24.0 mg/dL and creatinine value > 1.20 mg/dL) compared to the normal control group. This indicates that the hepatotoxic inducer given and nephrotoxic inducer given have been able to cause hepatotoxic in

the liver organ and nephrotoxic in the kidney organ.

The positive control group that was given a protector and was given an inducer showed significantly decreased in liver biochemical value (alanine transaminase value and aspartate transaminase value) and significantly decreased in kidney biochemical value (urea value and creatinine value) compared to the normal control group and the negative control group. Although the positive control group results showed significantly different from the normal control group, but the results obtained were within the normal range of the biochemical value (alanine transaminase value < 35.0 IU/L and aspartate transaminase value < 45.0 IU/L) and normal range of kidney biochemical value (urea value < 24.0 mg/dL and creatinine value < 1.20 mg/dL).

The test group that was given a passion fruit extract and was given an inducer showed significantly decreased in liver biochemical value (alanine transaminase value and aspartate transaminase value) and significantly decreased in kidney biochemical value (urea value and creatinine value) compared to the negative control group. This result means that the passion fruit peel extract has hepatoprotective activity and has nephroprotective activity. But only several treatment doses of the passion fruit peel extract were given a similar (not significantly different) to the positive control group.

The hepatoprotective activity for positive control (250 mg of curcumin per kg of body weight) was similar (not significantly different) to the 250 mg of purple passion fruit peel extract per kg of body weight, 250 mg of red passion fruit peel extract per kg of body weight, and 500 mg of yellow passion fruit peel extract per kg of body weight. The nephroprotective activity for positive control (50 mg of silymarin per kg of body weight) was similar (not significantly different) to the 250 mg of purple passion fruit peel extract per kg of body weight, 500 mg of red passion fruit peel extract per kg of body weight, and 500 mg of yellow passion fruit peel extract per kg of body weight.

## Discussions

Transaminase is a type of intracellular enzyme that is involved in nutrition metabolism. Transaminase enzymes are present in the cells of several organs such as the heart, liver, kidneys and pancreas. The alanine transaminase enzyme and the aspartate transaminase enzyme are present in hepatocytes cell and are released into the bloodstream in response to hepatocyte injury or death (hepatitis). Elevation of alanine transaminase enzyme or aspartate transaminase enzyme on the blood test for liver profile indicates the abnormality in the liver

organ. The good liver function will give a low value for alanine transaminase parameters and aspartate transaminase parameters [32], [33], [34], [35].

Both urea and creatinine are metabolic waste products that are excreted by the kidneys through urine, and only a small amount is left in the blood. If there is a disruption in kidney function, then there is an increase in these two parameters. Certain medical conditions can cause an increase in these two parameters, for example, chronic uncontrolled hypertension, uncontrolled diabetes mellitus, kidney stones, kidney inflammation, kidney infection, dehydration, and others [36], [37], [38], [39].

The hepatoprotective activity and the nephroprotective activity of an extract, in general, is a dose-dependent activity [40]. Purple passion fruit peel extract has the best hepatoprotective activity and the best nephroprotective activity compared to red passion fruit peel extract and yellow passion fruit peel extract. Although the purple passion fruit has the same genus with the red passion fruit and has the same species with the yellow passion fruit, due to genetic differences (differences in species or differences in varieties) there are morphological differences (colors and sizes) and genetical differences (sequences) [41], [42]. Also, due to differences in species or differences in varieties, it can cause qualitative differences in phytochemical contents and/or quantitative differences in phytochemical contents, also pharmacological differences and toxicological differences [43], [44].

The hepatoprotective activity possessed by the passion fruit peel extract may be due to the phytochemical content in the passion fruit peel. The phytochemical content in an extract is a good prospect for the development of phytotherapy [45], [46], [47]. The greater the content of alkaloids, flavonoids, and saponins in an extract, the higher the hepatoprotective activity possessed by the extract. Alkaloids are found in abundance in almost all parts of plants and have activities in scavenging the reactive oxygen species which can damage hepatocytes and are often useful compounds in medicinal chemistry for the development of new drugs. Flavonoids are polyphenol compounds that have been proven for hepatocytes protection from free radical scavenging activity. Saponins can directly protect hepatocytes from apoptosis through a mechanism of inhibition of the production of Tumor Nuclear Factor Alpha (TNF $\alpha$ ) [48], [49], [50].

The nephroprotective activity of the passion fruit peel extract is produced by the complex phytochemical content and varied phytochemical content contained in the passion fruit peel. An extract is thought to have nephroprotective activity through the mechanism of its antioxidant activity [51]. High flavonoid content extracts are reported to have high nephroprotective activity through an antioxidant mechanism [52]. The nephroprotective mechanism is

also shown by tannins compounds and which have good antioxidant activity [53]. Antioxidant compounds can protect the kidneys from alteration of kidney proteins and prevented the medical histopathological change of kidney tissue [54].

In conclusion, the extracts were shown a dose-dependent activity in hepatoprotective activity and nephroprotective activity. Purple passion fruit peel extract and red passion fruit peel extract have better hepatoprotective activity than yellow passion fruit peel extract. Purple passion fruit peel extract has better nephroprotective activity than red passion fruit peel extract and yellow passion fruit peel extract. The best hepatoprotective activity and nephroprotective activity showed by purple passion fruit peel extract compared to red passion fruit peel extract and yellow passion fruit peel extract.

## References

- Ozougwu J. Physiology of the Liver. *International Journal of Research in Pharmacy and Biosciences*. 2017; 4(8):13-24.
- Hoenig MP, Zeidel ML. Renal Physiology Homeostasis, the Milieu Interieur, and the Wisdom of the Nephron. *Clinical Journal of the American Society of Nephrology*. 2014; 9:1-13. <https://doi.org/10.2215/CJN.08860813> PMID:24789550 PMCid:PMC4078957
- Ramachandran A, and Jaeschke H. Mechanisms of Acetaminophen Hepatotoxicity and their Translation to the Human Pathophysiology. *Journal of Clinical and Translational Research*. 2017; 3(S1):157-69. <https://doi.org/10.18053/jctres.03.2017S1.002>
- Feng D, Wang Y, Wang H, Weng H, Kong X, Martin-Murphy BV, Li Y, Park O, Dooley S, Ju C, and Gao B. Acute and Chronic Effects of IL-22 on Acetaminophen-Induced Liver Injury. *Journal of Immunology*. 2014; 193:2512-8. <https://doi.org/10.4049/jimmunol.1400588> PMID:25063867 PMCid:PMC4135042
- Srivastava P, Rao RR, Shenoy PJ, Manjrekar PA, Teerthanath S, Bhuvaneshwari S. Nephroprotective Effect of Anethum graveolens in a Murine Model of Gentamicin Induced Nephrotoxicity. *Journal of Young Pharmacists*. 2018; 10(2):155-8. <https://doi.org/10.5530/jyp.2018.10.35>
- Aiswarya N, Rashmi RR, Preethi JS, Chandran V, Teerthanath S, Sunil PB, and Rakesh KB. Nephroprotective Effect of Aqueous Extract of Pimpinella anisum in Gentamicin Induced Nephrotoxicity in Wistar Rats. *Pharmacognosy Journal*. 2018; 10(3):403-7. <https://doi.org/10.5530/pj.2018.3.66>
- de Toledo NMV, de Camargo AC, Ramos PBM, Button DC, Granato D, and Canniatti-Brazaca SG. Potentials and Pitfalls on the Use of Passion Fruit by Products in Drinkable Yogurt - Physicochemical, Technological, Microbiological, and Sensory Aspects. *Beverages*. 2018; 4(47):1-15. <https://doi.org/10.3390/beverages4030047>
- Karsinah K, Hutabarat RC, and Manshur A. Passion Fruit Acid (*Passiflora edulis* Sims) Exotic Fruit Rich in Benefits. *Iptek Hortikultura*. 2010; (6):30-35.
- Marpaung AE, Karsinah K, and Br Karo B. Characterization and Evaluation of Passion Fruit Acid Hybrid from Purple and Red Passion Fruit Acid (*Passiflora* sp.). *Journal Hortikultura*. 2016; (26)2:163-70. <https://doi.org/10.21082/jhort.v26n2.2016.p163-170>
- Thokchom R, and Mandal G. Production Preference and Importance of Passion Fruit (*Passiflora edulis*). *Journal of*

- Agricultural Engineering and Food Technology. 2017; 4(1):27-30.
11. Anurangi CR, and Shamina S. Preliminary Phytochemical Screening and Acute & Subacute Toxicity Study on Different Concentrations of Unripen Fruit Peel Flour of *Passiflora edulis* in Male Albino Rats. *World Journal of Pharmacy and Pharmaceutical Sciences*. 2018; 7(3):828-34.
  12. Sihombing JR, Dharma A, Chaidir Z, Almahdy A, Fachrial E, and Munaf E. Phytochemical Screening and Antioxidant Activities of 31 Fruit Peel Extract from Sumatera, Indonesia. *Journal of Chemical and Pharmaceutical Research*. 2015; 7(11):190-6.
  13. Wasagu RSU, Lawal M, Amedu AM, Sabir AA, Kabir S, Tukur UG, and Zaharadeen A. Comparative Chemical Analysis, Phytochemical Screening and Antimicrobial Activities of the Rinds, Seeds and Juice of (*Passiflora edulis* var. *Flavicarpa*) Passion Fruit. *Journal of Natural Sciences Research*. 2016; 6(19):138-43.
  14. Wong YS, Sia CH, Khoo HE, Ang YK, Chang SK, and Yim HS. Influence of Extraction Conditions on Antioxidant Properties of Passion Fruit (*Passiflora edulis*) Peel. *Acta Scientiarum Polonorum Technologia Alimentaria*. 2014; 13(3):257-65. <https://doi.org/10.17306/J.AFS.2014.3.4> PMID:24887941
  15. Lima DS, Duarte NBA, Barreto DLC, de Oliveira GP, Takahashi JA, Fabrini SP, and Sande D. Passion Fruit and Apple - from Residues to Antioxidant, Antimicrobial and Anti-Alzheimer's Potential. *Ciência Rural Santa Maria*. 2018; 48(9):e20180076. <https://doi.org/10.1590/0103-8478cr20180076>
  16. Ramaiya SD, Bujang JS, and Zakaria MH. Assessment of Total Phenolic, Antioxidant, and Antibacterial Activities of *Passiflora* Species. *The Scientific World Journal*. 2014; 2014:ID167309.
  17. Doungue HT, Kengne APN, and Kuete D. Neuroprotective Effect and Antioxidant Activity of *Passiflora edulis* Fruit Flavonoid Fraction, Aqueous Extract, and Juice in Aluminum Chloride Induced Alzheimer's Disease Rats. *Nutrire*. 2018; 43:23-34. <https://doi.org/10.1186/s41110-018-0082-1>
  18. Raju IN, Reddy KK, Kumari CK, Reddy EB, Rao SD, Reddy CD, and Watson RR. Efficacy of Purple Passion Fruit Peel Extract in Lowering Cardiovascular Risk Factors in Type 2 Diabetic Subjects. *Journal of Evidence Based Complementary & Alternative Medicine*. 2013; 18(3):183-90. <https://doi.org/10.1177/2156587213475627>
  19. Cazarin CBB, da Silva JK, Colomeu TC, Batista AG, Vilella CA, Ferreira AL, Bogusz Junior S, Fukuda K, Augusto F, de Meirelles LR, de L Zollner E, and Marostica Junior MR. *Passiflora edulis* Peel Intake and Ulcerative Colitis - Approaches for Prevention and Treatment. *Experimental Biology and Medicine*. 2014; 239:542-51. <https://doi.org/10.1177/1535370214525306> PMID:24623393
  20. Farid R, Rezaieyazdi Z, Mirfeizi Z, Hatef MR, Mirheidari M, Mansouri H, Esmaeli H, Bentley G, Lu Y, Foo Y, Watson RR. Oral Intake of Purple Passion Fruit Peel Extract Reduces Pain and Stiffness and Improves Physical Function in Adult Patients with Knee Osteoarthritis. *Nutrition Research*. 2010; 30(9):601-6. <https://doi.org/10.1016/j.nutres.2010.08.010> PMID:20934601
  21. Silva DC, Freitas ALP, Pessoa CDS, Paula RCM, Mesquita JX, Leal LKAM, Brito GAC, Gonçalves DO, and Viana GSB. Pectin from *Passiflora edulis* Shows Antiinflammatory Action as well as Hypoglycemic and Hypotriglyceridemic Properties in Diabetic Rats. *Journal of Medicinal Food*. 2011; 14(10):1118-26. <https://doi.org/10.1089/jmf.2010.0220> PMID:21554121
  22. Anusooriya P, Malarvizhi D, Gopalakrishnan VK, and Devaki K. Antioxidant and Antidiabetic Effect of Aqueous Fruit Extract of *Passiflora ligularis* Juss. on Streptozotocin Induced Diabetic Rats. *International Scholarly Research Notices*. 2014; 2014:ID130342.
  23. Srivastava R, and Srivastava P. Hepatotoxicity and the Role of Some Herbal Hepatoprotective Plants in Present Scenario. *Global Journal of Digestive Diseases*. 2018; 3(2):1-4. <https://doi.org/10.4172/2472-1891.100034>
  24. Torres-González L, Cienfuegos-Pecina E, Perales-Quintana MM, Alarcon-Galvan G, Mu-oz-Espinosa LE, Pérez-Rodríguez E, Cordero-Pérez P. Nephroprotective Effect of *Sonchus oleraceus* Extract against Kidney Injury Induced by Ischemia Reperfusion in Wistar Rats. *Oxidative Medicine and Cellular Longevity*. 2018; 2018:ID9572803.
  25. Irawan C, Foliatini F, Hanafi H, Sulistiawaty L, Sukiman M. Volatile Compound Analysis Using GC-MS, Phytochemical Screening and Antioxidant Activities of the Husk of "Julang-Jaling" (*Archidendron bubalinum* (Jack) I.C Nielsen) from Lampung, Indonesia. *Pharmacognosy Journal*. 2018; 10(1):92-8. <https://doi.org/10.5530/pi.2018.1.17>
  26. Zhang QW, Lin LG, and Ye WC. Techniques for Extraction and Isolation of Natural Products - a Comprehensive Review. *Chinese Medicine*. 2018; 13:20-45. <https://doi.org/10.1186/s13020-018-0177-x> PMID:29692864 PMCid:PMC5905184
  27. Abdel-Hady H, El-Sayed M, Abdel-Hady AA, Hashash MM, Abdel-Hady AM, Aboushousha T, Abdel-Hameed ES, Abdel-Lateef EE, and Morsi EA. Nephroprotective Activity of Methanolic Extract of *Lantana camara* and Squash (*Cucurbita pepo*) on Cisplatin Induced Nephrotoxicity in Rats and Identification of Certain Chemical Constituents of *Lantana camara* by HPLC-ESI-MS. *Pharmacognosy Journal*. 2018; 10(1):136-47. <https://doi.org/10.5530/pi.2018.1.24>
  28. Fatima N, and Sultana A. Renoprotective and Antioxidant Effects of *Coleus forskohlii* against Gentamicin Induced Nephrotoxicity in Albino Wistar Rats. *Acta Pharmaceutica Scientia*. 2018; 56(2):67-84. <https://doi.org/10.23893/1307-2080.APS.05612>
  29. Tung BT, Hai NT, and Son PK. Hepatoprotective Effect of Phytosome Curcumin against Paracetamol Induced Liver Toxicity in Mice. *Brazilian Journal of Pharmaceutical Sciences*. 2017; 53(1):e16136. <https://doi.org/10.1590/s2175-97902017000116136>
  30. Okokon JE, Simeon JO, and Umoh EE. Hepatoprotective Activity of the Extract of *Homalium letestui* Stem against Paracetamol Induced Liver Injury. *Avicenna Journal of Phytomedicine*. 2017; 7(1):27-36. PMID:28265544 PMCid:PMC5329174
  31. Thuwaini MM, Abdul-Mounther M, and Kadhem HS. Hepatoprotective Effects of the Aqueous Extract of Clove (*Syzygium aromaticum*) against Paracetamol Induced Hepatotoxicity and Oxidative Stress in Rats. *European Journal of Pharmaceutical and Medical Research*. 2016; 3(8):36-42.
  32. Deb S, Puthanveetil P, and Sakharkar P. Population Based Cross Sectional Study of the Association between Liver Enzymes and Lipid Levels. *International Journal of Hepatology*. 2018; 2018:ID1286170.
  33. Ahmed Z, Ahmed U, Walayat S, Ren J, Martin DK, Moole H, Koppe S, Yong S, Dhillion S. Liver Function Tests in Identifying Patients with Liver Disease. *Clinical and Experimental Gastroenterology*. 2018; 11:301-7. <https://doi.org/10.2147/CEG.S160537> PMID:30197529 PMCid:PMC6112813
  34. Kim HR, and Han MA. Association between Serum Liver Enzymes and Metabolic Syndrome in Korean Adults. *International Journal of Environmental Research and Public Health*. 2018; 15:1658-64. <https://doi.org/10.3390/ijerph15081658> PMID:30081587 PMCid:PMC6121325
  35. Newsome PN, Cramb R, Davison SM, Dillon JF, Foulerton M, Godfrey EM, Hall R, Harrower U, Hudson M, Langford A, Mackie A, Mitchell-Thain R, Sennett K, Sheron NC, Verne J, Walmsley M, and Yeoman A. Guidelines on the Management of Abnormal Liver Blood Tests. *Gut*. 2018; 67:6-19. <https://doi.org/10.1136/gutjnl-2017-314924> PMID:29122851 PMCid:PMC5754852
  36. Tvarijonavičute A, Pardo-Marin L, Tecles F, Carrillo JD, Garcia-Martinez JD, Bernal L, Pastor J, Cerón JJ, and Martínez-Subiela S. Measurement of Urea and Creatinine in Saliva of Dogs - Pilot Study. *BMC Veterinary Research*. 2018; 14:223-8. <https://doi.org/10.1186/s12917-018-1546-5> PMID:30029662 PMCid:PMC6053741
  37. Romero-Aroca P, Baget-Bernaldiz M, Navarro-Gil R, Moreno-Ribas A, Valls-Mateu A, Sagarra-Alamo R, Barrot-De La Puente JF, and Mundet-Tuduri X. Glomerular Filtration Rate and/or Ratio of Urine Albumin to Creatinine as Markers for Diabetic Retinopathy - Ten-Year Follow Up Study. *Journal of Diabetes Research*. 2018; 2018:ID5637130.

38. Arihan O, Wernly B, Lichtenauer M, Franz M, Kabisch B, Muessig J, Masyuk M, Lauten A, Schulze PC, Hoppe UC, Kelm M, and Jung C. Blood Urea Nitrogen (BUN) is Independently Associated with Mortality in Critically Ill Patients Admitted to Intensive Care Unit (ICU). *Plos One*. 2018; 13(1):e0191697. <https://doi.org/10.1371/journal.pone.0191697> PMID:29370259 PMCID:PMC5784990
39. Lopez-Giacoman S, and Madero M. Biomarkers in Chronic Kidney Disease, from Kidney Function to Kidney Damage. *World Journal of Nephrology*. 2015; 4(1):57-73. <https://doi.org/10.5527/wjn.v4.i1.57> PMID:25664247 PMCID:PMC4317628
40. Chakraborty M, Bagchi B, Das S, Basu R, and Nandy P. A Dose Dependent Hepatoprotective and Nephroprotective Activity of Eucalyptus Oil on Streptozotocin Induced Diabetic Mice Model. *Clinical Phytoscience International Journal of Phytomedicine and Phytotherapy*. 2018; 4:10-9. <https://doi.org/10.1186/s40816-018-0067-x>
41. Ramaiya SD, Bujang JS, and Zakaria MH. Genetic Diversity in *Passiflora* Species Assessed by Morphological and Internal Transcribed Spacer (ITS) Sequence Analysis. *The Scientific World Journal*. 2014; 2014:ID598313.
42. Şesan TE, Sârbu A, Smarandache D, Oancea F, Oancea A, Savin A, Toma A, Ştefan L, Negru G, Bira AF, Vlăsceanu G, Ghiurea M, Jecu L, Vasilescu G, and Pomohaci CM. Botanical and Phytochemical Approach on *Passiflora* spp. - New Nutraceutical Crop in Romania. *Journal of Plant Development*. 2016; 23:97-126.
43. Sameh S, Al-Sayed E, Labib RM, and Singab AN. Genus *Spondias* - a Phytochemical and Pharmacological Review. *Evidence Based Complementary and Alternative Medicine*. 2018; 2018:ID5382904.
44. Abarca-Vargas R, and Petricevich VL. *Bougainvillea* Genus - a Review on Phytochemistry, Pharmacology, and Toxicology. *Evidence Based Complementary and Alternative Medicine*. 2018; 2018:ID9070927.
45. Abdelhafez OH, Fawzy MA, Fahim JR, Desoukey SY, Krischke M, Mueller MJ, and Abdelmohsen UR. Hepatoprotective Potential of *Malvaviscus arboreus* against Carbon Tetrachloride Induced Liver Injury in Rats. *Plos One*. 2018; 13(8):e0202362. <https://doi.org/10.1371/journal.pone.0202362> PMID:30138328 PMCID:PMC6107176
46. Agbafor KN, Ogbanshi ME, and Akubugwo EI. Phytochemical Screening, Hepatoprotective and Antioxidant Effects of Leaf Extracts of *Zapoteca portoricensis*. *Advances in Biological Chemistry*. 2014. 4:35-9. <https://doi.org/10.4236/abc.2014.41006>
47. Fan YC, Yue SJ, Guo ZL, Xin LT, Wang CY, Zhao DL, Guan HS, and Wang CY. Phytochemical Composition, Hepatoprotective, and Antioxidant Activities of *Phyllodium pulchellum* (L.) Desv. *Molecules*. 2018; 23:1361-75. <https://doi.org/10.3390/molecules23061361> PMID:29874868 PMCID:PMC6100508
48. Talluri MR, Gummadi VP, Battu GR. Chemical Composition and Hepatoprotective Activity of *Saponaria officinalis* on Paracetamol Induced Liver Toxicity in Rats. *Pharmacognosy Journal*. 2018; 10(6):1196-201. <https://doi.org/10.5530/pj.2018.6.205>
49. Rehman S, and Khan H. Advances in Antioxidant Potential of Natural Alkaloids. *Current Bioactive Compounds*. 2017; 13(2):101-8. <https://doi.org/10.2174/1573407212666160614075157>
50. Mohapatra SS, Roy RK, Mohan P, Upadhyaya TN, and Sarma J. Phytochemical Analysis and Hepatoprotective Effect of Hydroethanolic Extract of Stem Bark of *Oroxylum indicum*. *International Journal of Current Microbiology and Applied Sciences*. 2018; 7(1):1000-6. <https://doi.org/10.20546/ijcmas.2018.701.120>
51. Ahmad F, Al-Subaie AM, Al-Ohali AI, and Mohammed AS. Phytochemical and Nephroprotective Activity of *Eclipta prostrata* against Gentamicin Induced Nephrotoxicity in Wistar Rats. *International Journal of Pharma Research and Health Sciences*. 2018; 6(2):2559-64.
52. Mahipal P, and Pawar RS. Nephroprotective Effect of *Murraya koenigii* on Cyclophosphamide Induced Nephrotoxicity in Rats. *Asian Pacific Journal of Tropical Medicine*. 2017; 10(8):808-12. <https://doi.org/10.1016/j.apjtm.2017.08.005> PMID:28942830
53. Jedage HD, and Manjunath KP. Phytochemical, Pharmacological Evaluation of *Morinda pubescens* J.E.Sm. Bark Extract for Nephroprotective Activity. *Ayu*. 2016; 37(3-4):244-9. [https://doi.org/10.4103/ayu.AYU\\_196\\_15](https://doi.org/10.4103/ayu.AYU_196_15) PMID:29491678 PMCID:PMC5822978
54. Shah NA, Khan MR, and Nigussie D. Phytochemical Investigation and Nephroprotective Potential of *Sida cordata* in Rat. *BMC Complementary and Alternative Medicine*. 2017; 17:388-97. <https://doi.org/10.1186/s12906-017-1896-8> PMID:28778164 PMCID:PMC5545008