

Alarming Eating Behaviours among Adolescents in Egypt

Azza Abd El-Shaheed, Nermine N. Mahfouz^{*}, Rehab S.I. Moustafa, Mona A. Elabd

Department of Child Health, National Research Centre (33rd El Bohouth st, former El Tahrir St, Dokki, PO 12622), Egypt;
Medical Research Centre of Excellence (MRCE), Giza, Egypt

Abstract

Citation: Abd El-Shaheed A, Mahfouz NN, Moustafa RSI, Elabd, MA. Alarming Eating Behaviours among Adolescents in Egypt. Open Access Maced J Med Sci. <https://doi.org/10.3889/oamjms.2019.583>

Keywords: Eating behaviours; Adolescents; BMI; Obese; Non-obese

***Correspondence:** Nermine N. Mahfouz. Department of Child Health, National Research Centre (33rd El Bohouth st, former El Tahrir St, Dokki, PO 12622) and Medical Research Centre of Excellence (MRCE), Giza, Egypt. E-mail: nerninabil@yahoo.com

Received: 21-Apr-2019; **Revised:** 12-Jun-2019;
Accepted: 23-Jun-2019; **Online first:** 13-Jul-2019

Copyright: © 2019 Azza Abd El-Shaheed, Nermine N. Mahfouz, Rehab S.I. Moustafa, Mona A. Elabd. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Funding: This research was financially supported by the National Research Centre (NRC), Giza, Egypt

Competing Interests: The authors have declared that no competing interests exist

BACKGROUND: Adolescence is a phase of rapid growth and increased nutritional needs. It includes the stressful burden of pubertal changes, both physically and psychologically. Moreover, it is associated with the utmost need for independence and identity formation. An adolescent finds a great way to practice taking their own decisions by making personal food choices. But unfortunately, wrong dietary choices lead to unsatisfactory nutritional status.

AIM: To investigate the prevalence of six eating behaviours among adolescents.

SUBJECTS AND METHODS: A case-control study was conducted on ninety Egyptian adolescents from 10 to 18 years old. Anthropometric measurements were taken. Body Mass Index (BMI) was calculated. The cases were forty-five children with body mass index \geq 85th percentile. The control group involved forty-five of matched peers with body mass index $<$ 85th centile. A questionnaire form was constructed according to local customs in Egypt.

RESULTS: Two unhealthy behaviours were mostly found in our study group. The first and the predominant one was multitasking while eating practised by 92.1% of candidates and showing the equal distribution in both groups. The second was skipping breakfast and was adopted by 51.7% of the study group with a significantly higher distribution in the $<$ 85th centile group.

CONCLUSION: Faulty eating is a behaviour encountered in adolescence irrespective to BMI category. Thus, a normal BMI does not reflect healthy dietary behaviours.

Introduction

As a result of globalisation, heterogeneous western food habits were intruded in developing countries. This went in parallel with an increment in the frequency of chronic non-transmissible diseases. Thus, it became crucial to reassess the eating patterns in these countries and to evaluate their possible contribution as risk factors for these diseases [1].

Chronic non-transmissible diseases include overweight and obesity. The pervasiveness of high body mass index in the Middle East has been ranked the second worldwide, next to North America. Obesity is considered a serious health hazard, universally [2].

A sedentary lifestyle and faulty eating habits are two key threats that predispose to chronic non-transmissible diseases. Fortunately, these two risk factors are adjustable [3].

The World Health Organization (WHO) announced that almost 66.7% of early deaths, and 33.3% of grown-up illnesses, are consequent to undesirable behaviours that started in youth [4].

Thus the negative behaviours in adolescence are real challenges to face and overcome as early as possible before jeopardising lifetime health status [5]. That is why we should do this research.

Subjects and methods

The current study was conducted in the Nutrition Immunotherapy Clinic at the Medical Research Centre of Excellence (MRCE), National Research Centre (NRC). As a part of the in-house project entitled "Early Renal injury markers in obese adolescents".

Ninety Egyptian adolescents of both sexes were enrolled in this case-control study. The cases were forty-five children with BMI above or equal to 85th centile. The control group involved forty-five of matched peers with BMI below the 85th centile.

Inclusion criteria: adolescents of both sexes who are 10 to 18 years old.

Exclusion criteria: syndromic (e.g. Prader-Willi syndrome) and endocrinal (e.g. hypothyroidism) causes of obesity

Anthropometric measures: were assessed with a record of the height and the weight of each candidate. The height was measured to the nearest 0.5 cm on a Holtain portable anthropometer. The weight was determined to the nearest 0.1 kg on a Seca scale Balance with the subject dressed minimum clothes and no shoes. BMI was calculated as Weight (kg)/Height (m²). Data were plotted on WHO curves through the software AnthroCalc v1.66 Home

A questionnaire form was constructed in the light of literature and modified according to local customs in Egypt. A "yes", or "no" or "sometimes" answers were obtained for each question by a personal interview with every child. The six questions were about: breakfast meal, fast food, multitask eating, dinner meal content, dinner meal timing and intake of sweetened beverages.

Ethical approval

The protocol of the study was approved by the "Ethical Committee" of the NRC. Written informed consent was signed by the legal guardian of each participant before enrollment in the study by the code of ethics of the world medical association (Declaration of Helsinki) approval number 16130.

Statistical analysis

Analysis of data was performed by using Statistical Package for the Social Science SPSS version 16.G. Data were presented as a mean and standard deviation. Chi-square test was conducted for the variables and was used to detect the significant difference in the distribution between groups at P - value < 0.05 (SPSS version 16.G.)

Results

The study group comprised of ninety adolescents. They were 31 (34.4%) males and 59 (65.6%) females. Their mean age was (12.62 ± 2.6) and (13.05 ± 2.61) for case and control groups

respectively. According to BMI percentiles, we stratified the candidates into two equal groups of 45 children each. The cutoff BMI for the case group was ≥ 85th centile. While the cutoff BMI for the control group was < 85th centile, this was done according to the WHO growth charts. Detailed anthropometric data are presented as mean and standard deviation in Table 1.

Table 1: Anthropometric data

Variable	Case group (n = 45)	Control group (n = 45)
	Mean ± SD	
WT	73.41 ± 18.26	37.39 ± 10.68
WT Centile	96.26 ± 4.80	27.43 ± 25.05
HT	154.16 ± 10.65	146.05 ± 13.04
HT Centile	51.50 ± 28.89	32.60 ± 28.74
BMI	30.55 ± 5.61	17.22 ± 2.71
BMI Centile	98.30 ± 2.68	34.40 ± 28.16

Every child was personally interviewed to answer a form of questionnaires. This form was composed of six questions to monitor one desirable and five undesirable behaviours. The desirable one was tackled in question (Q1) and was about commitment to the breakfast meal. More than half of the participants (51.7%) skipped their breakfast either daily or sometimes with a per cent of 34.8% and 16.9% respectively.

The most prominent undesirable behaviour found was multitask eating (eating + watching TV) in Q3, it was adopted by a vast majority of 92.1% participants whether daily versus sometimes with a per cent of 91% and 1.1% respectively.

Otherwise, less than half of the study group members practised the four-remaining unhealthy behaviours, either daily or sometimes, as follows in a decrement order:

Ingestion of fast food > 2 times\week (Q2) and eating a heavy meal at dinner (Q4) were both encountered in a comparable per cent of participants (46.1%).

To a lesser extent, unhealthy intake of sugary beverages (Q6) was found in a per cent of 38.2%. A slightly lower per cent of 35.9% of candidates slept post-dinner by less than two hours (Q5), as illustrated in Table 2.

Table 2: Questions and per cent of different answers in the whole study group

Nine questions	Per cent		
	Yes	No	Sometimes
Q1. A daily commitment to breakfast intake	48.3	34.8	16.9
Q2. Eat ready-made fast food more than twice a week	27.0	53.9	19.1
Q3. Eating in front of the television	91.0	7.9	1.1
Q4. Eat a heavy meal at dinner	28.1	53.9	18.0
Q5. Sleep after dinner by less than two hours	25.8	64.0	10.1
Q6. Three sugary drinks per day (soft drink/juices reserved / warm drinks + 3 teaspoonful extra sugar per cup)	32.6	61.8	5.6

Then a comparison between the two groups was made as regards the frequency of each behaviour (as shown in Table 3).

A significantly higher number among obese

adolescents consumed breakfast everyday (n = 27) compared to non-obese group (n = 16) (P-value = 0.042).

The fast-food consumption was more frequent in obese (13 daily + 12 sometimes) than non-obese (11 daily+ 5 sometimes) adolescents but with insignificant P-value.

The majority of candidates in both groups were eating while watching TV (93.3% of the obese and 88.6% of the non-obese).

Almost similar per cent of candidates from the two groups was drinking three sugary beverages daily (33.3% of the obese and 31.8% of the non-obese).

Unexpectedly, the two unhealthy behaviours of ingesting a heavy dinner and late dinner time were both encountered in a greater number of the non-obese (n = 15 and n = 14) versus the obese ones (n = 10 and n = 9).

Table 3: Comparison of the dietary habit's responses of the two groups

Parameters	Groups	Obese	Non-Obese	Chi-Square	P-value
Q1. Regular Breakfast intake	Yes	N 27	16	6.361	0.042*
		% 60.00%	36.40%		
	No	N 14	17		
		% 31.10%	38.60%		
	Some-times	N 4	11		
		% 8.90%	25.00%		
Q2. Fast Food > twice/week	Yes	N 13	11	4.372	0.112
		% 28.90%	25.00%		
	No	N 20	28		
		% 44.40%	63.60%		
	Some-times	N 12	5		
		% 26.70%	11.40%		
Q3. Eating in front of the TV	Yes	N 42	39	1.243	0.537
		% 93.30%	88.60%		
	No	N 3	4		
		% 6.70%	9.10%		
	Some-times	N 0	1		
		% 0.00%	2.30%		
Q4. Heavy Dinner meal	Yes	N 10	15	1.739	0.419
		% 22.20%	34.10%		
	No	N 27	21		
		% 60.00%	47.70%		
	Some-times	N 8	8		
		% 17.80%	18.20%		
Q5. < 2 hours dinner/bed-time	Yes	N 9	14	2.047	0.359
		% 20.00%	31.80%		
	No	N 32	25		
		% 71.10%	56.80%		
	Some-times	N 4	5		
		% 8.90%	11.40%		
Q6. Three Sugary Beverages/day	Yes	N 15	14	0.241	0.886
		% 33.30%	31.80%		
	No	N 27	28		
		% 60.00%	63.60%		
	Some-times	N 3	2		
		% 6.70%	4.50%		

*P-value < 0.05 considered significant.

Discussion

The WHO described adolescence as a highly vulnerable stage of life. Because, children at this stage, have a deceiving grown-up a physique that hides an immature psychosocial aspect. Therefore, the concern of the current study was directed towards children at this critical age. Same age group was also chosen to investigate dietary habits in Syria, Sudan, Arabian Gulf countries, China, India, Britain and many

other countries [6], [7], [5], [8], [9], and [10].

The participants were grouped according to BMI percentiles. As in numerous studies and various fields of research, the BMI categorisation was the criteria of stratifying the candidates into case and control groups. This "BMI-based selection" in Egyptian studies was adopted by many authors [11], [12], [13], [14].

We classified the studied population into a case group with a BMI \geq 85th centile and a control group with a BMI < 85th centile. The same cutoff for BMI percentile was chosen by bin Zaal and co-workers in their study about adolescents' dietary habits [15].

A personal face-to-face interviewing was chosen to ensure complete data collection by avoiding questions' misunderstanding and missing answers. The structured questionnaire used was quick, easy, reproducible, coded and interpreted. Six straightforward, closed-ended questions were the tool to retrieve information. These were questions with a limited fixed set of responses (yes/no/sometimes). In other studies, the self-administered questionnaires were used instead of interviewing technique. For example, the Arab Teens Lifestyle Study (ATLS) in which the large study group rendered personal interview impossible. Also, in the ATLS survey, a set of scaled questions was used [1].

In the current study, Q1 monitored the healthy behaviour of regular breakfast intake. More than half of the participants (51.7%) were breakfast skippers either daily or sometimes. Comparably, in the ATLS survey, 52% to 82% of Arab adolescents did not have breakfast every day [16]. Smaller per cent was reported in a Sudanese study where only 244 out of 945 adolescents (25.8%) were skipping breakfast meal [16]. While, in the survey done by Li et al., in China, the majority (93%) of the adolescents were having breakfast regularly (1661 from a total of 1774) [8].

A significantly higher number of breakfast consumers was noticed among the case group compared to the control group (P-value 0.042). On the contrary, several studies linked omitting breakfast with high BMI in childhood and adolescence [17], [18], [19], [20], [21]. Although Albertson et al., and Berkey et al., and recently Casazza et al., and Dhurandhar et al., did not support this "breakfast skippers\high BMI link" in their researches [22], [23], [24], [25]. This contradictory is due to the variability of meal content. If the consumed meal were rich in fibres and micronutrients, then it would have been an addition to the healthy eating index score. Also, it would have contributed to a lower intake of sweets and Trans-fat. But if the consumed meal was of poor quality high in fat and sugar, then skipping, it would not be harmful [26] and [27].

According to responses to Q2, candidates

with BMI < 85th centiles were consuming less fast food than those with BMI ≥ 85th centiles. This finding matched with the conclusion deduced by Bhadoran et al. They stated that junk food intake more than once a week was linked to a higher risk of obesity while eating fast food more than twice a week was associated with a higher risk of metabolic syndrome [28]. Similarly, Bhattacharjee et al. found a positive correlation between the frequency of junk food intake and high BMI among Indian adolescents [9].

As regards the answers to Q3, multitask eating was the most prominent undesirable behaviour found in the study group. It was practised daily by 91% of participants irrespective to BMI categories. This bad habit results in overeating, improper chewing; distraction of the brain from appropriate processing of eating and deprive the child of family interaction during mealtime. In an Egyptian study conducted in Cairo, a comparable per cent of 87.5% of adolescents were taking snacks while watching TV [29]. In Canadian research, eating in front of the TV was positively associated with unhealthy food choices [30].

Through the unpredictable answers to Q4 and Q5, it was noticed that the combined two unhealthy behaviours of late and heavy dinner meal were commoners in non-obese candidates and not their counterparts. The reverse was detected by Ong et al., who noted an association between high BMI and ingesting calorie-dense food late at night [31]. This unexpected finding may be due to the widespread Egyptian habit of skipping lunch and getting the main meal at dinner time. Thus, the total caloric intake may be adequate or even below the requirement of the day. Also, awareness of the overweight/obesity problem may be the reason for those with high BMI to get a light early dinner meal as a way of caloric restriction.

The unhealthy behaviour of drinking three sugary beverages per day, in Q6, showed an equivocal distribution in both groups (33.3% of the obese and 31.8% of the non-obese). This comparable per cent may be due to adolescents' tendency to imitate their friends in drinking soft drinks and preserved juices. Moreover, in Egypt, dark sweetened tea is quite popular, and it is considered the everyday beverage in most Egyptian homes. So, it is an Egyptian society issue rather than a high BMI related one. In an Egyptian study conducted in both rural and urban sectors, by Abdel-Hady et al., eighty per cent of adolescents had more than 3 small glasses of sweetened black tea daily [32]. In Oman and the United Arab Emirates, more than 50% of the teenagers' males and about 50% of the teenagers' females consumed ≥ 1 soft drink/day in the month before the study. While in Qatar, 60%/65% of adolescent's boys/girls respectively drank soft drinks every day [5].

In conclusion, many undesirable dietary

behaviours are acquired in adolescence. Such behaviours are encountered in all BMI categories. Thus, a normal BMI does not always reflect healthful dietary intake. The rectification of faulty eating "behaviours" in youth is mandatory before becoming "habits" in adulthood.

Recommendations; The most powerful influencers on eating behaviours in adolescence are family, friends and media. Thus, good parental support and understanding have a major imprint on reforming such harmful conduct. Also, scholastic healthy eating programs are extremely important as a source of easy access to widespread information. The college's canteen must be an ideal exemplar of how healthy eating should be. The advertisements about food products should be under strict supervision. Further research studies are needed to explore more about eating behaviors on a larger number of adolescents and in different countries.

Limitations: The study was limited by the small number of participants.

Acknowledgement

The authors are grateful to the administrators, the laboratory team and the nurses of the Medical Research Centre of Excellence (MRCE). We are thankful to the National Research Centre (NRC), Giza, Egypt, for funding this research. Also, we express our appreciation to all the children who participated in the study and their parents.

References

1. Al-Hazaa Hazzaa M, Musaiger Abdulrahman O, ATLS Research Group. Arab Teens Lifestyle Study (ATLS): objectives, design, methodology and implications. Dove Press journal: Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy. 2011; 4:417-426. <https://doi.org/10.2147/DMSO.S26676> PMID:22253540 PMCID:PMC3257970
2. Mahfouz NN, Fahmy RF, Nassar MS, Wahba SA. Body Weight Concern and Belief among Adolescent Egyptian Girls. Open Access Maced J Med Sci. 2018; 6(3):582-587. <https://doi.org/10.3889/oamjms.2018.145> PMID:29610625 PMCID:PMC5874390
3. Donnelly TT, Fung TS, Al-Thani A-AbM. Fostering active living and healthy eating through understanding physical activity and dietary behaviours of Arabic-speaking adults: a cross-sectional study from the Middle East. BMJ. 2018; 8:e019980. <https://doi.org/10.1136/bmjopen-2017-019980> PMID:29678976 PMCID:PMC5914903
4. Makhlof Obermeyer C. Adolescents in Arab countries: Health statistics and social context, DIFI Family Research and Proceedings. 2015:1-17. <https://doi.org/10.5339/difi.2015.1>
5. Al Makadma AS. Adolescent health and health care in the Arab

- Gulf countries: today's needs and tomorrow's challenges. *International Journal of Pediatrics and Adolescent Medicine*. 2017; 4(1):1-8. <https://doi.org/10.1016/j.ijpam.2016.12.006> PMID:30805493 PMCID:PMC6372452
6. Bashour HN. Survey of dietary habits of in-school adolescents in Damascus, Syrian Arab Republic. *EMHJ-Eastern Mediterranean Health Journal*. 2004; 10(6):853-862.
7. Musaiger AO, Nabag FO, Al-Mannai M. Obesity, dietary habits, and sedentary behaviors among adolescents in Sudan: alarming risk factors for chronic diseases in a poor country. *Food and nutrition bulletin*. 2016; 37(1):65-72. <https://doi.org/10.1177/0379572116629244> PMID:26880662
8. Li M, Dibley MJ, Sibbritt DW, Yan H. Dietary habits and overweight/obesity in adolescents in Xi'an City, China. *Asia Pac J Clin Nutr* 2010; 19(1):76-82.
9. Prasun B, Sujaya M, Payas J, Sahibjeet S. Food habits and obesity: a study in adolescents. *Int J Contemp Pediatr*. 2017; 4(2):336-340. <https://doi.org/10.18203/2349-3291.ijcp20170526>
10. Al-Hazaa H, Al-Nakeeb Y, Duncan M, Al-Sobayel H, Abahussain N, Musaiger A, Lyons M, Collins P, Nevill A. A cross-cultural comparison of health behaviors between Saudi and British adolescents living in urban areas: gender by country analyses. *International journal of environmental research and public health*. 2013; 10(12):6701-20. <https://doi.org/10.3390/ijerph10126701> PMID:24300072 PMCID:PMC3881136
11. Ibrahim OM, Gabre AA, Sallam SF, El-Alameey IR, Sabry RN, Galal EM, Tawfik SM, Zarouk WA, Mosaad RM, Ramadan A. Influence of Interleukin-6 (174G/C) Gene Polymorphism on Obesity in Egyptian Children. *Open Access Maced J Med Sci*. 2017; 5(7):831-835. <https://doi.org/10.3889/oamjms.2017.175> PMID:29362605 PMCID:PMC5771281
12. Mahfouz Nermine N., Fahmy Reham F., Nassar Maysa S., Wahba Saneya A. Body Weight Concern and Belief among Adolescent Egyptian Girls. *Open Access Macedonian Journal of Medical Sciences*. 2018; 6(3):582-587. <https://doi.org/10.3889/oamjms.2018.145> PMID:29610625 PMCID:PMC5874390
13. El Wakeel MA, El-Kassas GM, Kamhawy AH, Galal EM, Nassar MS, Hammad EM, El-Zayat SR. Serum Apelin and Obesity-Related Complications in Egyptian Children. *Open Access Maced J Med Sci*. 2018; 6(8):1354-1358. <https://doi.org/10.3889/oamjms.2018.312> PMID:30159056 PMCID:PMC6108807
14. El Kassas GM, Shehata MA, El Wakeel MA, Amer AF, Elzaree FA, Darwish MK, Amer MF. Role of Procalcitonin As an Inflammatory Marker in a Sample of Egyptian Children with Simple Obesity. *Open Access Maced J Med Sci*. 2018; 6(8):1349-1353. <https://doi.org/10.3889/oamjms.2018.323> PMID:30159055 PMCID:PMC6108804
15. Bin Zaal AA, Musaiger AO, D'Souza R. Dietary habits associated with obesity among adolescents in Dubai, United Arab Emirates. *Nutricion hospitalaria*. 2009; 24(4):437-44.
16. Abdulrahman M. Unhealthy Eating Habits, Physical Inactivity and Sedentary Behaviours among Arab Adolescents the Main Risk Factors for Chronic Disease. *Aspetar Sports Medicine Journal*. 2016; 5(2):384-388.
17. Alexander KE, Ventura EE, Spruijt-Metz D, Weigensberg MJ, Goran MI, Davis JN. Association of breakfast skipping with visceral fat and insulin indices in overweight Latino youth. *Obesity*. 2009; 17(8):1528-33. <https://doi.org/10.1038/oby.2009.127> PMID:19424166 PMCID:PMC2836758
18. Fiore H, Travis S, Whalen A, Auinger P, Ryan S. Potentially protective factors associated with healthful body mass index in adolescents with obese and nonobese parents: a secondary data analysis of the third national health and nutrition examination survey, 1988-1994. *Journal of the American Dietetic Association*. 2006; 106(1):55-64. <https://doi.org/10.1016/j.jada.2005.09.046> PMID:16390667
19. Shafiee G, Kelishadi R, Qorbani M, Motlagh ME, Taheri M, Ardalan G, Taslimi M, Poursafa P, Heshmat R, Larijani B. Association of breakfast intake with cardiometabolic risk factors. *Jornal de pediatria*. 2013; 89(6):575-82. <https://doi.org/10.1016/j.jped.2013.03.020> PMID:24029551
20. Vanelli M, Iovane B, Bernardini A. Breakfast habits of 1,202 northern Italian children admitted to a summer sport school. Breakfast skipping is associated with overweight and obesity. *Acta Bio Medica Atenei Parmensis*. 2005; 76(2):79-85.
21. Szajewska H, Ruszczyński M. Systematic review demonstrating that breakfast consumption influences body weight outcomes in children and adolescents in Europe. *Critical reviews in food science and nutrition*. 2010; 50(2):113-9. <https://doi.org/10.1080/10408390903467514> PMID:20112153
22. Albertson AM, Franko DL, Thompson D, Eldridge AL, Holschuh N, Affenito SG, Bauserman R, Striegel-Moore RH. Longitudinal patterns of breakfast eating in black and white adolescent girls. *Obesity*. 2007; 15(9):2282-92. <https://doi.org/10.1038/oby.2007.271> PMID:17890497
23. Berkey CS, Rockett HR, Gillman MW, Field AE, Colditz GA. Longitudinal study of skipping breakfast and weight change in adolescents. *International journal of obesity*. 2003; 27(10):1258-1266. <https://doi.org/10.1038/sj.ijo.0802402> PMID:14513075
24. Casazza K, Brown A, Astrup A, Bertz F, Baum C, Brown MB, Dawson J, Durant N, Dutton G, Fields DA, Fontaine KR. Weighing the evidence of common beliefs in obesity research. *Critical reviews in food science and nutrition*. 2015; 55(14):2014-53. <https://doi.org/10.1080/10408398.2014.922044> PMID:24950157 PMCID:PMC4272668
25. Dhurandhar EJ. True, true, unrelated? A review of recent evidence for a causal influence of breakfast on obesity. *Current Opinion in Endocrinology, Diabetes and Obesity*. 2016; 23(5):384-8. <https://doi.org/10.1097/MED.0000000000000281> PMID:27584010
26. Medin AC, Myhre JB, Diep LM, Andersen LF. Diet quality on days without breakfast or lunch-Identifying targets to improve adolescents' diet. *Appetite*. 2019; 135:123-30. <https://doi.org/10.1016/j.appet.2019.01.001> PMID:30639294
27. Panizza C, Shvetsov Y, Harmon B, Wilkens L, Le Marchand L, Haiman C, Reedy J, Boushey C. Testing the predictive validity of the healthy eating index-2015 in the multiethnic cohort: is the score associated with a reduced risk of all-cause and cause-specific mortality? *Nutrients*. 2018; 10(4):452. <https://doi.org/10.3390/nu10040452> PMID:29621192 PMCID:PMC5946237
28. Bahadoran Z, Mirmiran P, Azizi F. Fast food pattern and cardiometabolic disorders: a review of current studies. *Health promotion perspectives*. 2015; 5(4):231-240. <https://doi.org/10.15171/hpp.2015.028> PMID:26933642 PMCID:PMC4772793
29. Hussien Hanan A. Effect of Food Habits on the Nutritional Status of Children in Cairo. *Biomedical Statistics and Informatics* 2017; 2(1):10-17.
30. Borghese MM, Tremblay MS, Leduc G, Boyer C, Bélanger P, LeBlanc AG, Francis C, Chaput JP. Independent and combined associations of total sedentary time and television viewing time with food intake patterns of 9-to 11-year-old Canadian children. *Applied Physiology, Nutrition, and Metabolism*. 2014; 39(8):937-43. <https://doi.org/10.1139/apnm-2013-0551> PMID:24892903
31. Halib H, Qian OY, Suan WB. Breakfast Intake and its Association with Body Mass Index among Pre-schoolers in Taska Permata Keluarga Kuala Nerus, Terengganu. *Malaysian Journal of Applied Sciences*. 2018; 3(2):57-70.
32. Abdel-Hady D, El-Gilany AH, Sarraf B. Dietary habits of adolescent students in Mansoura, Egypt. *International Journal of Collaborative Research on Internal Medicine & Public Health*. 2014; 6(6):132. <https://doi.org/10.1155/2014/258470> PMID:24895560 PMCID:PMC4033417