

Misidentification of *S. suis* as a Zoonotic Agent

Ni Made Adi Tarini¹, Marta Setiabudy^{1*}, NM Susilawathi², NND Fatmawati¹, IPB Mayura¹, EA Darwinata¹, NKA Sudiariani²

¹Department of Microbiology, Faculty of Medical and Health Sciences Udayana University, Denpasar, Bali, Indonesia;

²Department of Neurology, Faculty of Medical and Health Sciences Udayana University, Denpasar, Bali, Indonesia

Abstract

Citation: Tarini NMA, Setiabudy M, Susilawathi NM, Fatmawati NND, Mayura IPB, Darwinata EA, Sudiariani NKA. Misidentification of *S. suis* as a Zoonotic Agent. Open Access Maced J Med Sci. 2019 Jul 30; 7(14):2309-2312. https://doi.org/10.3889/oamjms.2019.619

Keywords: Meningitis; Misdiagnosis; PCR; Streptococcus suis; Zoonosis

***Correspondence:** Marta Setiabudy. Department of Microbiology, Faculty of Medical and Health Sciences Udayana University, Denpasar, Bali, Indonesia. E-mail: martasetiabudy@gmail.com

Received: 06-May-2019; **Revised:** 05-Jun-2019; **Accepted:** 06-Jun-2019; **Online first:** 25-Jul-2019

Copyright: © 2019 Ni Made Adi Tarini, Marta Setiabudy, NM Susilawathi, NND Fatmawati, IPB Mayura, EA Darwinata, NKA Sudiariani. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Funding: This research was financially supported by the Research and Community Services Unit, Faculty of Medicine, Udayana University, Indonesia by grant number 230.5/UN14.2.2/PD/2017

Competing Interests: The authors have declared that no competing interests exist

BACKGROUND: *Streptococcus suis* is an emerging zoonotic pathogen. This bacterium commonly causes meningitis in human and is often associated with hearing and vestibular dysfunction. *S. suis* tends to be misidentified, leading to under-diagnosis.

CASE PRESENTATION: A previously healthy 50-year-old man was admitted to one of the district hospitals in Bali Province, Indonesia, due to meningitis. He had a history of consuming homemade raw pork product two days before the onset of illness. *Streptococcus mitis* was identified from the cerebrospinal fluid culture by using VITEX 2 COMPACT (BiomeriueX) with a 99% probability score. This patient had clinical symptoms and risk factor identical to *S. suis* infection. Therefore, we performed confirmation tests for the cerebrospinal fluid by PCR (using primer specific for *gdh* and *recN*) and sequencing of those PCR products. Both of the confirmation tests showed a positive result for *S. suis*.

CONCLUSION: There are few reports of *S. suis* infections in Indonesia, but we believe that the cases in Indonesia, especially Bali, are not uncommon. The under-reported cases are perhaps due to the difficulties in differentiating *S. suis* from other *Streptococcus* species by culture method, particularly *Streptococcus mitis*. Therefore, confirmation by PCR is necessary.

Introduction

Streptococcus suis (*S. suis*) is the main pathogen in pigs that can cause meningitis, endocarditis, and sepsis in humans [1], [2], [3]. *S. suis* infection is an emerging infectious disease whose incidence is reported to increase and causes public health problems [4], [5], [6]. There are two outbreaks of *S. suis* infection in China that affect more than 200 people with a mortality rate of almost 20%, which have since changed the views of experts on the threat of this pathogen to human health [1]. The first case of *S. suis* infection was reported in 1954 after an

outbreak of meningitis, septicemia, and arthritis in piglets. The first case of *S. suis* meningitis in a human was reported in Denmark in 1968 [7], and it has occurred in Bali in 2014, which was followed by an increase in cases of infection in Bali [8].

S. suis is a gram-positive coccus, facultatively anaerobic, and based on Lancefield's classification it has a cell wall structure by group D *Streptococcus* [9]. These bacteria are often reported as *Streptococcus viridans* in their initial identification from CSF culture due to the similarity [10], [11].

Misidentification has an impact on the case finding and to control the transmission, which is highly important in the effort to comprehensively manage *S.*

suis as one of the zoonotic agents. The discussion of the following case is expected to enhance the understanding of the characteristics of *S. suis* and the possibility of identification errors that might occur.

Case Illustration

A 50-year-old male was brought to the emergency department in one of the regional hospitals in Bali with a decrease of consciousness accompanied by headache, fever, nausea, vomiting, and anorexia for 2 days before hospitalisation. He had a history of consuming homemade red *lawar* (a traditional Balinese food made by mixing chopped pork with fresh blood) three days before admission. He had no history of other acute infections. His past medical history was not significant.

On examination, his condition was confused with a Glasgow Coma Scale (GCS) E3V3M5, fever (38°C), and neck stiffness. Routine laboratory test showed leucocytosis ($15.9 \times 10^3/\mu\text{L}$) with neutrophil predominant (93.6%) and thrombocytopenia ($125 \times 10^3/\mu\text{L}$). A brain computed tomography scan (CT scan) demonstrated cerebral oedema. The Cerebrospinal fluid (CSF) had a turbid color, an increased leukocyte count ($1080 \text{ cell}/\text{mm}^3$) predominating polymorphs (71%), an increased protein level (226 mg/dL) and a decreased glucose level (65 mg/dL) with a decreased CSF / serum glucose ratio (0.38). The CSF culture showed Gram-positive cocci, negative catalase, and colony on blood agar showed hemolytic alpha. Identification with VITEK 2 COMPACT (bioMérieux) shows the results of *Streptococcus mitis* (*S. mitis*) with a probability of 99%.

The diagnosis of acute bacterial meningitis was established, and the patient was treated with 2 gram of ceftriaxone as empirical antibiotic at 12-hour intervals until the culture's result was done. After 3 days of treatment, the patient experienced hearing impairment. Based on those clinical features and history of consuming food containing raw pork products, *S. suis* meningitis was suspected. Furthermore, *S. mitis* as the culture result was known as a commensal bacteria and had never been reported as a cause of meningitis. We suspected there was a misidentification of the causative agent in this case. Hence, a confirmation examination was performed by PCR using the specific primers of the glutamate dehydrogenase (Gdh) and the recombination/repair protein (RecN) encoding gene in *S. suis*. The sequencing of those PCR products was also carried out to confirm whether the causative agent in these cases was *S. mitis* or *S. suis* [12], [13]. From the PCR results, it was found that the isolates showed the positive result in both amplified *S. suis* specific genes.



Figure 1: PCR *gdh* and *recN* gene detection. The amplicon was electrophoresed on 1 % agarose gel. The sample was positive for *gdh* gene (688bp) and *recN* gene (336bp). (M = marker 100 bp (Invitrogen); lane 1 = control positive for *gdh* gene; lane 2 = negative control; lane 3 = sample (*gdh* gene was positive); Lane 4 = Marker 100 bp (Invitrogen); Lane 5 = control positive for *recN* gene and lane 6 = sample (*recN* gene was positive)

The BLAST result of PCR product sequences showed 100% identity to *gdh* and 99% identity to *recN* of *S. suis* strains (CP020863.1). According to the performed confirmation tests, the patient was diagnosed with *S. suis* meningitis. After the patient had been hospitalized for 14 days, the patient was discharged from the hospital with the complications of hearing impairment.

Discussion

Regarding the case illustration above, our patient exhibits typical symptoms as acute bacterial meningitis including fever, headache, anorexia, hearing loss, and a history of consuming raw pork food products. *S. suis* can cause systemic infections in humans in the form of meningitis, sepsis, endocarditis, arthritis, endophthalmitis, uveitis, spondylodiscitis, ophthalmoplegia, and epidural abscess [7]. The most common manifestation of *S. suis* infection is meningitis that occurs in about two-thirds of patients [1], [2], [14]. One of the typical symptoms of *S. suis* meningitis is hearing impairment, or loss, which occurs in more than 50% of the cases [1], [2], [14]. Various risk factors associated with acquiring *S. suis* infection included the consumption of raw pork products, pig-related occupation, pigs, or pork exposure, alcohol drinking, skin injury especially due to exposure during pork processing, and underlying diseases contributing to immunocompromised conditions [2], [5].

Apparently, the incubation period of *S. suis* infection in humans varies depending on the route of transmission. A shorter incubation period occurs when infection happens through wounds on the skin which rapidly spread hematogenously, while the incubation period is longer if infected through oral consumption [7], [15], [16]. In the case of outbreaks in China, the

incubation period ranged from 3 hours to 14 days (median 2.2 days) [15]. *S. suis* infection often occurs in predominantly healthy adults, men with an average age of 51 years, but very rarely found in children [2], [16].

Microbiological examination plays an important role in establishing the diagnosis of meningitis, especially for identification of causative agents and antimicrobial sensitivity testing. The microbiological examinations of CSF fluid that were performed in this case — including the Gram staining, colony appearance on blood agar with alpha-hemolytic, and negative catalase test — matched the characteristics of *Streptococcus*. The result of bacterial identification using biochemical examination (VITEK 2 Compact) is *S. mitis*. However, the patient's history and clinical symptoms did not match with the bacterial identification result, *S. mitis*. This bacterium has never been reported as the causative agent of meningitis. Suspicion of other causative agents such as *S. suis* bacteria arose due to a history of contact with animals as hosts and hearing impairment that occur in the patient.

S. suis is a coccus Gram-positive bacterium, facultatively anaerobic, negative catalase, and forms alpha hemolysis in blood agar [17], [18]. If we review the results of the microbiological examination, it can be concluded that the bacteria have similarities with other *Streptococcus* bacteria such as *Streptococcus viridans*, *Streptococcus equid*, *Streptococcus mitis*, and other *Streptococcus* which often cause misidentification [4], [6], [11], [19]. In a study performed by Donsakul et al., from 1993 to 1999 (reported in 2003), 5 of the 8 cases were initially identified as *Streptococcus viridans* infections. The conventional biochemical examination has lower specificity than an examination of molecular biology [12].

In this case, PCR was performed using the specific primer to amplify two specific genes in *S. suis*, *gdh*, and *recN* [12], [13] and a positive result was shown, indicating that the bacteria isolated from the CSF is *S. suis*. The PCR method provides more specific result based on gene identification that encodes glutamate dehydrogenase (*gdh*) and the gene that encodes recombination, or repairs protein (*recN*) [11], [12], [20]. In 2015, Okwumabua et al. conducted a study that identified 306 *S. suis* isolates using PCR method through identification of the encoding gene *gdh*. The gene identification has been used in diagnosis of other bacterial infections and reported to have a very low mutation rate [12]. Identification of the *recN* gene is also an easy and accurate examination for *S. suis*. PCR based on the *recN* sequence is a better method for identification and detection of *S. suis* compared to *gdh* gene identification [11], because it has been shown to have a lower level of similarity at the species level and a higher divergence value at the subspecies level than other genes [21].

The diagnosis of *S. suis* meningitis is highly important, because it is one of the zoonotic diseases that can cause public health problems and potentially leads to an outbreak. It requires comprehensive prevention of the causative agents by breaking the chain of transmission through intermediary animals.

In conclusion, the low prevalence of *S. suis* meningitis cases reported in Indonesia, especially in Bali, does not indicate the actual condition of the cases. This could happen due to the misidentification of *S. suis* from other *Streptococcus sp.* from conventional cultural method. The choice of a more accurate type of examination such as PCR is highly important to consider as a confirmation test of CSF examination, especially in cases with clinical symptoms that were suspected as *S. suis* infection.

Acknowledgments

We would like to thank the staff of the laboratory of Clinical Microbiology at Sanglah Hospital Denpasar and staff of the Clinical Microbiology Department, faculty of Medicine Udayana University for their supports in this study. We also would like to thank the Research and Community Services Unit, Faculty of Medicine, Udayana University, since this research was partly funded by grant number 230.5/UN14.2.2/PD/2017.

References

1. van Samkar A, Brouwer MC, Schultsz C, van der Ende A, van de Beek D. Streptococcus suis Meningitis: A Systematic Review and Meta-analysis. *PLoS Negl Trop Dis*. 2015; 9(10):1-10. <https://doi.org/10.1371/journal.pntd.0004191> PMID:26505485 PMID:PMC4624688
2. Rayanakorn A, Goh BH, Lee LH, Khan TM, Saokaew S. Risk factors for Streptococcus suis infection: A systematic review and meta-analysis. *Sci Rep*. 2018; 8(1):1-9. <https://doi.org/10.1038/s41598-018-31598-w> PMID:30190575 PMID:PMC6127304
3. Goyette-Desjardins G, Auger JP, Xu J, Segura M, Gottschalk M. Streptococcus suis, an important pig pathogen and emerging zoonotic agent—an update on the worldwide distribution based on serotyping and sequence typing. Vol. 3, *Emerging Microbes and Infections*. 2014. <https://doi.org/10.1038/emi.2014.45> PMID:26038745 PMID:PMC4078792
4. Gottschalk M, Xu J, Calzas C, Segura M. Streptococcus suis, 2010; 371-91. <https://doi.org/10.2217/fmb.10.2> PMID:20210549
5. Ho DTN, Le TPT, Wolbers M, Cao QT, Nguyen VMH, Tran VTN, et al. Risk factors of Streptococcus suis infection in Vietnam. A case-control study. *PLoS One*. 2011; 6(3). <https://doi.org/10.1371/journal.pone.0017604> PMID:21408132 PMID:PMC3050921
6. Dutkiewicz J, Sroka J, Zając V, Wasiński B, Cisak E, Sawczyn A, Kloc A, Wójcik-Fatla A. Streptococcus suis: a re-emerging

- pathogen associated with occupational exposure to pigs or pork products. Part I-Epidemiology. *Annals of Agricultural and Environmental Medicine*. 2017; 24(4):683-95. <https://doi.org/10.26444/aaem/79813> PMID:29284248
7. Wertheim HFL, Nghia HDT, Taylor W, Schultz C. *Streptococcus suis*: An Emerging Human Pathogen. *Clin Infect Dis*. 2009; 48(5):617-25. <https://doi.org/10.1086/596763> PMID:19191650
8. Susilawathi NM, Tarini NMA, Sudewi AAR. Bacterial Meningitis *Streptococcus suis* With Bilateral Sensorineural Hearing Loss. *Neurona*. 2016; 34(1).
9. Feng Y, Zhang H, Wu Z, Wang S, Cao M, Hu D, et al. *Streptococcus suis* infection An emerging / reemerging challenge of bacterial infectious diseases ? *Streptococcus suis* infection. 2017; 5594(January):477-97. <https://doi.org/10.4161/viru.28595> PMID:24667807 PMCID:PMC4063810
10. Huang Y, Teng L, Ho S, Hsueh P. *Streptococcus suis* infection, 2005.
11. Okura M, Osaki M, Nomoto R, Arai S, Osawa R, Sekizaki T, Takamatsu D. Current taxonomical situation of *Streptococcus suis*. *Pathogens*. 2016; 5(3):45. <https://doi.org/10.3390/pathogens5030045> PMID:27348006 PMCID:PMC5039425
12. Okwumabua O, Connor MO, Shull E. A polymerase chain reaction (PCR) assay speci c for *Streptococcus suis* based on the gene encoding the glutamate dehydrogenase. 2015; 218(2003):3-8. [https://doi.org/10.1016/S0378-1097\(02\)01127-8](https://doi.org/10.1016/S0378-1097(02)01127-8)
13. Ishida S, Tien LHT, Osawa R, Tohya M, Nomoto R, Kawamura Y, et al. Development of an appropriate PCR system for the reclassification of *Streptococcus suis*. *J Microbiol Methods*. 2014; 107:66-70. <https://doi.org/10.1016/j.mimet.2014.09.003> PMID:25229648
14. Vu Thi Lan H, Ngo H, Nguyen Tien H, Horby P, Ho Dang Trung N, Vu Dinh T, et al. Epidemiology, Clinical Manifestations, and Outcomes of *Streptococcus suis* Infection in Humans. *Emerg Infect Dis*. 2014; 20(7):1105-14.
15. Yu H, Jing H, Chen Z, Zheng H, Liu L, Luo L, et al. Human *Streptococcus suis*. *Emerg Infect Dis*. 2006; 12(6):914-20. <https://doi.org/10.3201/eid1206.051194> PMID:16707046 PMCID:PMC3373052
16. Fong IW. Zoonotic *Streptococci*: A Focus on *Streptococcus suis*. In: *Emerging Zoonosis*. Springer International Publishing AG, 2017:189-210. https://doi.org/10.1007/978-3-319-50890-0_10 PMID:28039571
17. Isrina S, Salasia O, Mada UG, Nugroho W, Timika DP, Sandi NA, et al. Phenotypic and Genotypic Characterization of *Streptococcus suis* Isolated from Pigs in Papua. 2016.
18. Zaccaria E. Discovery, characterization and applications of natural DNA transformation in *Streptococcus suis* (Doctoral dissertation, Wageningen University).
19. Donsakul K, Dejthevaporn C, Witoonpanich R, Hospital R. Southeast Asian J Trop Med Public Health *Streptococcus suis* Infection : Clinical Features. 2003; 34(1).
20. Mahalaya S, Sandi NA, Cargill C, Nugroho W, Salasia SIO, Slipranata M. Phenotypic and Genotypic Characterization of *Streptococcus suis* Isolated from Pigs in Papua. *Asian J Anim Vet Adv*. 2016; 11(5):303-8. <https://doi.org/10.3923/ajava.2016.303.308>
21. Glazunova OO, Raoult D, Roux V. Partial recN gene sequencing: A new tool for identification and phylogeny within the genus *Streptococcus*. *Int J Syst Evol Microbiol*. 2010; 60(9):2140-8. <https://doi.org/10.1099/ijs.0.018176-0> PMID:19880633