

Dietary Ethanolic Extract of *Mangosteen pericarp* Reduces VCAM-1, Perivascular Adipose Tissue and Aortic Intimal Medial Thickness in Hypercholesterolemic Rat Model

Titin Andri Wihastuti¹, Fitria Nugraha Aini^{2,3}, Cholid Tri Tjahjono⁴, Teuku Heriansyah^{5*}

¹Department of Basic Science in Nursing, Faculty of Medicine, University of Brawijaya, Malang, Indonesia; ²Departement of Biomedicine, Faculty of Medicine, University of Brawijaya, Malang, Indonesia; ³Faculty of Medicine, University of Islam Malang, Malang, Indonesia; ⁴Department of Cardiology, Faculty of Medicine, Brawijaya University, Malang, Indonesia; ⁵Department of Cardiology, Faculty of Medicine, University of Syiah Kuala, Banda Aceh, Aceh, Indonesia

Abstract

Citation: Wihastuti TA, Aini FN, Tjahjono CT, Heriansyah T. Dietary Ethanolic Extract of *Mangosteen pericarp* Reduces VCAM-1, Perivascular Adipose Tissue and Aortic Intimal Medial Thickness in Hypercholesterolemic Rat Model. Open Access Maced J Med Sci. <https://doi.org/10.3889/oamjms.2019.717>

Keywords: Atherosclerosis; Dyslipidemia; Mangosteen Pericarp Ethanolic Extract; VCAM-1; PVAT; IMT

***Correspondence:** Teuku Heriansyah, Department of Cardiology, Faculty of Medicine, University of Syiah Kuala, Banda Aceh, Aceh, Indonesia. E-mail: teuku_hery@unsyiah.ac.id

Received: 21-May-2019; **Revised:** 21-Sep-2019; **Accepted:** 22-Sep-2019; **Online first:** 10-Oct-2019

Copyright: © 2019 Titin Andri Wihastuti, Fitria Nugraha Aini, Cholid Tri Tjahjono, Teuku Heriansyah. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Funding: This research was supported by the Ministry of Research, Technology and Higher Education of the Republic of Indonesia

Competing Interests: The authors have declared that no competing interests exist

BACKGROUND: High-fat diet (HFD) is associated with dyslipidemia which is a risk factor for atherosclerosis. Dyslipidemia causes oxidative stress which induces vascular cell adhesion molecule-1 (VCAM-1). Oxidative stress also triggers the thickening of tunica intima-media (IMT) and Perivascular Adipose Tissue (PVAT). Xanthone compound in ethanolic extract of *Mangosteen pericarp* (EEMP) has an antioxidant property to overcome the oxidative stress.

AIM: The objective of this study is to investigate the effect of dietary EEMP administration on the expression of VCAM-1 and thickness of PVAT and IMT in atherosclerotic rat model fed with HFD.

METHODS: This experimental laboratory study uses 25 Wistar strain *Rattus norvegicus* which were divided into 5 study groups. Negative Control group (GT1) was given a normal diet, Positive Control group (GT2) was treated with HFD, and three treatment groups were each treated with HFD with *Mangosteen pericarp* extract of 200 mg/kg BW (GT3), 400 mg/kg BW (GT4), and 800 mg/kg BW (GT5). Measurements of VCAM-1 expression were performed using immunofluorescence. PVAT and IMT measurements were performed on rat aortic preparations.

RESULTS: One-way ANOVA test showed the addition of dietary EEMP significantly ($p < 0.05$) decreased the expression of VCAM-1 and decreased the thickness of PVAT and IMT in treatment groups as compared with both negative and positive controls. Tukey HSD test showed a dose of 800 mg/kg BW was the most effective dose for decreasing VCAM-1 level, PVAT and IMT.

CONCLUSION: Dietary EEMP significantly decreases the expression of VCAM-1, as well as the thickness of PVAT and IMT in Wistar strain *Rattus norvegicus* treated with HFD.

Introduction

Cardiovascular disease such as hypertension, stroke, coronary heart disease and heart failure, is the leading cause of global death. In 2015, it was estimated that 17.7 million people or about 31% of the world's deaths were due to this disease [1]. Coronary heart disease develops from the atherosclerotic process. The exact cause of atherosclerosis still unknown, but research suggests that atherosclerosis

starts when certain factors damage the inner layers of the arteries [2].

Atherosclerosis is an irreversible process, causing disease by the slow development of narrowing of the arterial lumen, mainly located in the intima of middle-large size arteries [3], [4]. Atherosclerosis is progressive and leads to the formation of atherosclerotic plaques that causes obstruct in blood vessels. Atherosclerotic plaque formation begins from chronic inflammation which triggers oxidative stress. Oxidative stress is a state of

reactive oxygen species (ROS) and anti-oxidants imbalance [5]. Physiologically; ROS acts as a regulator in various processes of defence mechanisms, differentiation, proliferation, and migration of body cells. Conversely, an excess of ROS levels in the body is a sign of a pathological condition characterised by excessive inflammation and contributes to the pathogenesis of atherosclerosis [6], [7].

High ROS production can increase the production of PVAT and thickened IMT in dyslipidemia [8]. ROS can activate peroxisome proliferator-activated receptor- γ (PPAR- γ), which is a master regulator of adipogenesis, thus causing expansion and dysfunction of PVAT [9]. ROS also contribute to endothelial dysfunction characterised by increased expression of adhesion molecules such as vascular cell adhesion molecule 1 (VCAM-1) [10].

Garcinia mangostana L., or commonly known as mangosteen, is rich in phenol components, such as xanthenes, tannins, and anthocyanins [11]. These components have been shown to have many biological activities, and in particular, mangosteen pericarp extract has anti-inflammatory, anti-cancer, anti-microbial, and anti-oxidant effects both *in vivo* and *in vitro* [12]. Xanthone has a role as a scavenger antioxidant capable of inhibiting ROS [13]. Based on these facts, ethanolic extract of mangosteen pericarp (EEMP) is expected to inhibit VCAM-1 expression and decrease PVAT and IMT thickness in the aorta of atherosclerotic rat model treated with high-fat diet (HFD).

Material and Methods

Laboratory Animals

This research used 8-week-old male Wistar strain *Rattus norvegicus* weighing about 1.5-2 kg which was kept in Pharmacology Laboratory of Faculty of Medicine, Brawijaya University, Malang, East Java, Indonesia. Before the treatment, the rats underwent an acclimatization process for 2 weeks, and later were divided into 5 treatment groups (n = 5), i.e. Negative Control with normal diet (G1), Positive Control with HFD diet (G2), the group with HFD and 200 mg/kg BW EEMP administration (G3), the group with HFD and 400 mg/kg BW EEMP administration (G4), and the group with HFD and 800 mg/kg BW EEMP administration (G5).

Feeding and Creating Dyslipidemic Rat Model

Rats were fed according to their respective

treatment groups. The normal group was given a normal diet of PARS 62%. Groups with HFD were given 2% cholesterol, 0.2% cholic acid, and 5% lard oil supplements, 30 grams daily, *ad libitum* [1].

EEMP (Ethanolic extract of mangosteen pericarp) Process

The EEMP was obtained through drying, extraction, and evaporation processes. *Mangosteen pericarps* were washed and then dried in an oven at 80 degrees until dry or free of water content. The dried mangosteen pericarp was refined evenly into 100 grams of dried sample, and placed into a 1-litre Erlenmeyer flask. It was soaked in 70% ethanol until the volume reached 1000 mL, shaken for 30 minutes, and settled for 1 night until a sediment form. The next step was the evaporation process using *Rotary Evaporator*. The evaporation process is completed when the ethanol solution is separated from the active ingredient. The EEMP were administered to the research animals through a feeding tube with doses of 200 mg/kg BW, 400 mg/kg BW and 800 mg/kg BW respectively.

Rat Dissection

Dissection was performed 2 months subsequent of EEMP administration. Surgery was performed after the animal had been given ketamine as an anaesthetic agent. Later, the aorta was retrieved and preserved using 10% formalin for subsequent aortic preparation with haematoxylin-eosin staining.

Measurement of IMT and PVAT Thickness

Measurement of IMT and PVAT thickness was conducted at Anatomy Pathology Laboratory, Faculty of Medicine, Brawijaya University, Malang, East Java, Indonesia. Aortic preparation was done using paraffin block and *HE*. The measurements were made by drawing perpendicular vertical lines of the intima tunica and outline of the media tunica. Measurements were made in 5 clockwise zones and then averaged. The measurement of PVAT thickness was performed by measuring the mean of the lowest, medium, and highest thickness of PVAT on aortic preparations. Measurements with 400 x magnification and were made using *scan dot slide Olyvia* software.

VCAM-1 Measurement

Aortic tissue was fixated with PHEMO buffer (Pipes 0.068 M, HEPES 0.0025 M, EGTA 0.005 M, MgCl 0.003 M, 10% DMSO, PH 6.8), which contained 3.7% formaldehyde, 0.05% glutaraldehyde, 0.5 triton x-100, for 10 minutes at room temperature. Coverslip was blocked by 10% goat serum/PBS for 10 minutes.

VCAM-1 expression on aortic tissue was identified through double-staining immunofluorescence using anti-rat VCAM-1 with rhodamine as the secondary antibody, and α -actin was coloured by fluorescein isothiocyanate as the secondary antibody, and then when observed with Confocal Laser Scanning Microscope (CLSM), the double-stained result would show VCAM-1 expression in smooth muscle cells. Lastly, the collected data were quantitatively analysed with Olympus fluoview 1.7A software.

Ethics

The medical research ethics committee of the Faculty of Medicine of Brawijaya University has stated that this research is ethically detailed with No.211 / EC / KEPK-S1-PD / 06 / 2017

Statistical Analysis

Statistical analysis was performed using SPSS version 16 software with significance level of 0.05 ($p = 0.05$) and 95% confidence interval ($\alpha = 0.05$). This research used One Way ANOVA Parametric Test to determine the effect of EEMP in 5 groups of an atherosclerotic rat model. Data analysis was further conducted using Tukey HSD Test to understand the difference between groups.

The presentation of ANOVA and Post Hoc Tukey HSD data tables refer to Seng *et al.*, (2018) [15].

Results

Histopathological Findings of IMT and PVAT

Measurements were done with 400 x magnification and were made using *scan dot slide Olyvia* software. Blackline that is pointed by red arrow shows the tunica intima-media thickness and black arrow of each image shows the PVAT thickness. Histopathological features of the aorta between 5 groups show a clear difference in tunica intima-media and PVAT thickness. Intima-media of picture D has a similar thickness to intima-media of picture A as the negative control. Tukey HSD test shows that the dosage of EEMP 200 mg/kg BW can lower the BMI, with the mean dosage value closest to negative control is 400 mg/kg BW. The thickness of PVAT in figure E is thinner than others, shown as the length of the black arrow. Tukey HSD Test shows that PVAT thickness in 800 mg/kg BW EEMP group was significantly different compared to the HFD group.




Figure 1: Images of aortic preparation with tunica intima-media and PVAT thickness measurement on each research animal treatment group; A) Normal diet group (GT1); B) HFD group (GT2); C) the group with HFD and 200 mg/kg BW EEMP administration (GT3); D) the group with HFD and 400 mg/kg BW EEMP administration (GT4); and (E) the group with HFD and 800 mg/kg BW EEMP administration (GT5)

Immunofluorescence Findings of VCAM-1

On this study, statistically significant differences were seen in all of the parameters; IMT ($p < 0.001$), PVAT ($p = 0.008$), VCAM-1 ($p = 0.032$), HDL ($p = 0.010$), LDL ($p < 0.001$), TG ($p = 0.007$), and TC ($p < 0.001$). From the data presented in Table 1, it appears that there was a worsening of lipid profiles in the group of rats with HFD administration.




Figure 2: A) VCAM-1; B) α -actin expression; C) expression of VCAM-1 in Smooth Muscle Cell (SMC); Immunofluorescence findings of VCAM-1 in Smooth Muscle Cell (SMC), labelled with α -actin; A) VCAM-1 – rhodamine; B) α -actin – fluorescein isothiocyanate (FITC). The colour green shows α -actin expression (SMC marker); C) Double staining of VCAM-1 and α -actin showed the expression of VCAM-1 in smooth muscle cells of aortic tissue. The colour yellow shows expression of VCAM-1 in SMC

Then with EEMP treatment, there was a significant improvement of lipid profile from ONEWAY ANOVA test. Not only improved lipid profiles, but EEMP administration can also reduce IMT, VCAM-1, and PVAT thickness with significant ONE-WAY ANOVA test results.

Table 1: IMT, PVAT thickness, and VCAM-1 Expression Calculation, and Lipid Profile variables at Groups of Treatment (GT1-GT5) with ANOVA and Tukey HSD results

| Variables | Groups of Treatment | Mean \pm SD | F-statistics (df) | P | Tukey HSD | 95% CI | adj p | |
|-----------------|---------------------|---------------------|-------------------|---------|-----------|--------|-------|---------|
| IMT (μ m) | GT1 | 61.05 \pm 2.01 | 13.998 (24) | <0.001 | GT1-GT2 | 0.98 | 3.21 | < 0.001 |
| | GT2 | 78.19 \pm 6.33 | | | GT1-GT3 | -1.02 | 3.35 | 0.128 |
| | GT3 | 62.76 \pm 6.71 | | | GT1-GT4 | -0.67 | 1.59 | 0.468 |
| | GT4 | 61.10 \pm 3.35 | | | GT1-GT5 | -0.47 | 0.65 | 0.263 |
| | GT5 | 60.59 \pm 1.31 | | | GT2-GT3 | 1.68 | 3.27 | < 0.001 |
| PVAT (μ m) | GT1 | 547.48 \pm 152.10 | 4.608 (24) | 0.008 | GT2-GT4 | 1.22 | 3.14 | < 0.001 |
| | GT2 | 744.24 \pm 115.00 | | | GT2-GT5 | 0.54 | 3.06 | < 0.001 |
| | GT3 | 737.00 \pm 58.27 | | | GT3-GT4 | -1.15 | 2.93 | 0.206 |
| | GT4 | 711.64 \pm 112.11 | | | GT3-GT5 | -0.83 | 2.25 | 0.068 |
| | GT5 | 554.77 \pm 33.95 | | | GT4-GT5 | -0.33 | 1.76 | 0.104 |
| VCAM-1 (AU) | GT1 | 477.01 \pm 81.35 | 3.542 (24) | 0.032 | GT1-GT2 | 3.42 | 6.28 | 0.006 |
| | GT2 | 628.51 \pm 18.00 | | | GT1-GT3 | 1.68 | 3.76 | 0.012 |
| | GT3 | 497.52 \pm 64.70 | | | GT1-GT4 | 2.82 | 4.08 | 0.020 |
| | GT4 | 451.26 \pm 100.84 | | | GT1-GT5 | -4.42 | 1.36 | 0.075 |
| | GT5 | 447.71 \pm 101.30 | | | GT2-GT3 | -3.84 | 0.87 | 0.187 |
| HDL (mg/dL) | GT1 | 30.00 \pm 7.52 | 4.865 (24) | 0.010 | GT2-GT4 | -4.86 | 2.48 | 0.245 |
| | GT2 | 22.25 \pm 6.50 | | | GT2-GT5 | 1.27 | 5.08 | 0.046 |
| | GT3 | 28.75 \pm 7.50 | | | GT3-GT4 | -2.63 | 1.34 | 0.591 |
| | GT4 | 37.25 \pm 7.13 | | | GT3-GT5 | 2.74 | 4.92 | 0.024 |
| | GT5 | 41.50 \pm 5.25 | | | GT4-GT5 | 1.04 | 3.86 | 0.012 |
| LDL (mg/dL) | GT1 | 60.25 \pm 16.19 | 15.474 (24) | <0.001 | GT1-GT2 | 0.38 | 2.74 | 0.026 |
| | GT2 | 126.25 \pm 17.03 | | | GT1-GT3 | -3.26 | 1.82 | 0.143 |
| | GT3 | 89.50 \pm 9.84 | | | GT1-GT4 | -2.84 | 2.49 | 0.281 |
| | GT4 | 78.75 \pm 12.94 | | | GT1-GT5 | -3.04 | 1.86 | 0.309 |
| | GT5 | 65.75 \pm 8.26 | | | GT2-GT3 | 2.32 | 4.68 | 0.027 |
| TG (mg/dL) | GT1 | 101.75 \pm 6.80 | 5.331 (24) | 0.007 | GT2-GT4 | 2.28 | 5.86 | 0.018 |
| | GT2 | 146.25 \pm 33.38 | | | GT2-GT5 | 1.96 | 4.70 | 0.009 |
| | GT3 | 102.50 \pm 12.17 | | | GT3-GT4 | -2.46 | 1.68 | 0.074 |
| | GT4 | 101.75 \pm 11.70 | | | GT3-GT5 | -3.86 | 2.92 | 0.065 |
| | GT5 | 97.50 \pm 10.66 | | | GT4-GT5 | -3.58 | 3.28 | 0.073 |
| TC (mg/dL) | GT1 | 107.75 \pm 10.53 | 18.442 (24) | < 0.001 | GT1-GT2 | -2.38 | 3.31 | 0.086 |
| | GT2 | 181.25 \pm 23.72 | | | GT1-GT3 | 0.86 | 5.18 | < 0.001 |
| | GT3 | 113.25 \pm 12.55 | | | GT1-GT4 | -0.98 | 4.32 | 0.012 |
| | GT4 | 104.00 \pm 18.31 | | | GT1-GT5 | -2.45 | 1.72 | 0.087 |
| | GT5 | 102.50 \pm 7.18 | | | GT2-GT3 | 1.36 | 4.27 | < 0.001 |
| | | | | | GT2-GT4 | 0.46 | 2.62 | < 0.001 |
| | | | | | GT2-GT5 | 2.38 | 3.86 | < 0.001 |
| | | | | | GT3-GT4 | 1.26 | 2.76 | < 0.001 |
| | | | | | GT3-GT5 | 0.26 | 2.39 | < 0.001 |
| | | | | | GT4-GT5 | -1.41 | 3.84 | 0.065 |
| | | | | | GT1-GT2 | 0.97 | 4.05 | 0.013 |
| | | | | | GT1-GT3 | -2.25 | 5.72 | 0.067 |
| | | | | | GT1-GT4 | -1.36 | 3.84 | 0.248 |
| | | | | | GT1-GT5 | -1.72 | 3.24 | 0.357 |
| | | | | | GT2-GT3 | 4.84 | 8.63 | 0.004 |
| | | | | | GT2-GT4 | 4.81 | 8.97 | 0.002 |
| | | | | | GT2-GT5 | 5.94 | 9.08 | 0.001 |
| | | | | | GT3-GT4 | -0.45 | 4.64 | 0.492 |
| | | | | | GT3-GT5 | -2.76 | 3.86 | 0.074 |
| | | | | | GT4-GT5 | -3.46 | 4.06 | 0.062 |
| | | | | | GT1-GT2 | 2.76 | 5.87 | < 0.001 |
| | | | | | GT1-GT3 | -4.64 | 1.08 | 0.083 |
| | | | | | GT1-GT4 | -5.26 | 2.21 | 0.074 |
| | | | | | GT1-GT5 | -5.46 | 1.98 | 0.059 |
| | | | | | GT2-GT3 | 2.43 | 4.66 | 0.001 |
| | | | | | GT2-GT4 | 1.24 | 3.57 | < 0.001 |
| | | | | | GT2-GT5 | 0.69 | 2.74 | 0.001 |
| | | | | | GT3-GT4 | -4.79 | 3.27 | 0.056 |
| | | | | | GT3-GT5 | -4.02 | 1.85 | 0.086 |
| | | | | | GT4-GT5 | -2.52 | 0.82 | 0.074 |

Footnote: GT1: Negative control with normal diet; GT2: Positive control with HFD diet; GT3: the group with HFD and 200 mg/kg BW EEMP administration; GT4: the group with HFD and 400 mg/kg BW EEMP administration; GT5: the group with HFD and 800 mg/kg BW EEMP administration. Abbreviation: HFD, High Fatty Diet; IMT, Intima-Media Thickness; PVAT, Peri Vascular Adipocyte Tissue; VCAM-1, Vascular Cell Adhesion Molecule, HDL, High-Density Lipoprotein, LDL, Low-Density Lipoprotein; TG, Triglyceride; TC, Total Cholesterol.

Tukey HSD test was performed on VCAM-1, IMT and PVAT showed that a dose of 400 mg/kg BW was a significant dose to decrease IMT and VCAM-1 thickness because approached with normal diet group and a dose of 800 mg/kg BW was a significant dose to decrease PVAT thickness approached with normal diet group.

Discussion

The peel of mangosteen fruit has long been used by residents in various countries as an ingredient in traditional medicine. Several experimental studies have proved that EEMP could be used as anti-tumour, anti-inflammatory, anti-bacterial, anti-viral, and anti-oxidant agents. In previous studies, the ethanolic extract of mangosteen pericarp was mentioned to have the highest antioxidant effect [14]. The EEMP as a source of antioxidants works by releasing electrons to free radicals to form stable products to prevent the generation of chain reactions [15]. Xanthone compound in the pericarp of mangosteen is needed by the body for the balance of pro-oxidants. Xanthone has binding properties towards unstable free oxygen. This free oxygen acts as free radical and destroying body cells; therefore, xanthone can inhibit the degeneration process of cell damage [15]. Several *in vitro* studies have shown that mangosteen pericarp extract can absorb free radicals [16]. Xanthones are divided into α -mangosteen, β -mangosteen, γ -mangosteen, and methoxybetamangosteen, but the most common form is α -mangosteen [17]. The α -mangosteen antioxidant may increase lipoprotein lipase enzyme activity and improve very-low-density lipoprotein (VLDL) catabolism. As a result, the total cholesterol, triglyceride, and LDL levels fall, and HDL or good cholesterol level rises [18].

In Table, HFD administration has been shown to worsen the lipid profiles of research animals significantly. This worsening of lipid profile shows the presence of dyslipidemia. Dyslipidemia is one of the risk factors for atherosclerosis [17]. HFD causes dyslipidemia, and manifestations are elevated LDL level in the body; in a condition of excessive fat intake, LDL may accumulate in IMT and PVAT of the blood vessels. On the other hand, dyslipidemia will cause adipocyte cells hypertrophy. Adipocyte growth is associated with an increase of pro-oxidant enzymes, such as NADPH oxidase, and a decrease of antioxidant enzymes, such as catalase [19]. These two conditions will increase the H_2O_2 production. H_2O_2 reacts with Fe_2^+ to release hydroxide (OH^-) via Fenton reaction. OH^- is a free radical that oxidises LDL accumulated in tunica intima; this process is called lipid peroxidation, and it will produce oxidised LDL (oxLDL); from this stage, the atherosclerosis process begins. OxLDL will later lead to an increase of proinflammatory cytokines, such as *Tumor Necrosis Factor- α* (TNF- α) [17]. Increased proinflammatory cytokines will trigger an increase in VCAM-1 expression [20]. Increased expression of VCAM-1 causes accumulation of monocytes in IMT. The adhesion of monocyte-endothelial cells to atherosclerotic lesions is affected by intercellular oxidative stress in Mononuclear Cells (MNC) and causes the thickening of IMT [21]. The administration of HFD which may cause thickening of PVAT will also

lead to an increase in PVAT proinflammatory property that is characterised by decreased expression of adiponectin and increased expression of IL-6, IL-8, and monocyte chemoattractant protein-1 (MCP-1). These parameters (VCAM-1, PVAT, and IMT) play a role in atherosclerosis pathogenesis.

VCAM-1 is also a molecule that plays a role in atherogenesis. One component in atheroma plaque is accumulated leukocytes. Adhesion molecules, such as VCAM-1, activates endothelial cell signal transduction, which in turn alters the form of endothelial cells to pave the way for leukocyte migration [22]. Increased expression of VCAM-1 in HFD group showed a positive tendency. This result is accordance with previous research by Huang *et al.*, reported that after treating rats with HFD for 12 weeks, VCAM-1 expression in the aorta and the levels of sVCAM-1 were increased significantly [23]. After EEMP administration, VCAM-1 expression displayed a significant decrease. Aside from being an anti-oxidant, xanthone also has anti-inflammatory properties. In an experiment conducted by Wihastuti *et al.*, the administration of EEMP exhibit the ability to lower TNF- α and IL-1 levels in high-cholesterol diet-induced rats compared to the positive control group. The two cytokines are known to be related to the expression of VCAM-1 [24]. These findings will contribute to further improving our understanding of the underlying mechanism regulating adhesion molecule levels by exploring the effects of EEMP on the pathogenesis of atherosclerosis.

The thickness of the research animal's PVAT in the HFD group (seen in Table) experienced a significantly different increase, as evidenced by Tukey HSD *post-hoc* test results. HFD administration may induce a pro-inflammatory phenotype in PVAT with low adiponectin expression and high IL-6, IL-8, and MCP-1 expression [25]. This thickening of PVAT illustrates a condition of inflammation and endothelial dysfunction as a result of an increase in ROS [25]. Consistent with our studies, others reported that PVAT plays a role in the pathogenesis of vascular lesion formation. Ketonen *et al.* reported that obesity-induced endothelial dysfunction is caused by increased oxidative stress and enhanced expression of inflammatory cytokine in PVAT [26]. In this research, the administration of EEMP that contains xanthone as an anti-oxidant agent was proved by ANOVA parametric test to be able to decrease the thickness of PVAT significantly. The EEMP dose that significantly lowered PVAT thickness was 800 mg. This EEMP dose could decrease PVAT thickness almost until the level of normal group rats. PVAT thickness in 800 mg/kg BW EEMP group was significantly different compared to the HFD group. The increase of free radical and oxidative stress conditions in atherosclerosis leads to the thickening of adipocyte tissue that produces proinflammatory and anti-inflammatory cytokines. This event is characterised by an increase in PVAT thickness observed using

histopathologic imaging. In addition to PVAT, oxidative stress is also evident from the increasing number of foam cells and narrowing of the lumen diameter of the blood vessel due to atherosclerotic plaque deposition [27].

Based on results, it can be concluded that administration of mangosteen pericarp ethanol extract (*Garcinia mangostana Linn*) may decrease the expression of VCAM-1, IMT and PVAT thickness in research animals with the high-fat diet group. 400 mg/kg BW of EEMP was proved to exhibit IMT and VCAM-1 expression value, and 800 mg/kg BW of EEMP was able to exhibit the value of PVAT thickness which was equivalent to a normal physiological condition in research animal groups. The results of this study were similar to other studies which proved that the administration of anti-oxidants might reduce oxidative stress, therefore inhibiting atherosclerosis process by inhibiting LDL oxidation and decreasing ROS in endothelial cells [9].

Acknowledgement

We want to thank Prof. Dr Djangan Sargowo, Sp.JP, Sp.PD, FIHA, FACC, FESC, FCAPC, FASC for support and advice towards the completion of this research. We would also like to thank the Head of the Medical Faculty for the kind collaboration of Laboratory of Pharmacology, Laboratory of Pathology Anatomy, and Laboratory of Biomedicine, Faculty of Medicine, Brawijaya University, Malang, East Java, Indonesia.

References

1. World Health Organization. Cardiovascular diseases, 2017. Available from: http://www.who.int/cardiovascular_diseases/world-heart-day-2017/en/. Accessed: 27.9.2017.
2. National Heart, Lung, and Blood Institute. At a glance : Atherosclerosis. Available from: https://www.nhlbi.nih.gov/files/docs/public/heart/atherosclerosis_at_glance.pdf
3. Frostegard J. SLE, Atherosclerosis and Cardiovascular Disease. *J Intern Med.* 2005; 257:485-495. <https://doi.org/10.1111/j.1365-2796.2005.01502.x> PMID:15910552
4. Jain PK, Joshi SC. Atherosclerosis, Inflammation, and Oxidative Stress. *Suresh Gyan Vihar University International Journal of Environment, Science and Technology.* 2016; 2(2):17-29.
5. Birben E, Sahiner UM, Sackesen C, Erzurum S, Kalayci O. Oxidative Stress and Antioxidant Defense. *The World Allergy Organization Journal.* 2012; 5(1):9-19. <https://doi.org/10.1097/WOX.0b013e3182439613> PMID:23268465 PMCid:PMC3488923
6. Handy DE, Loscalzo J. Responses to reductive stress in the cardiovascular system. *Free Radic Biol Med.* 2017; 109:114-24. <https://doi.org/10.1016/j.freeradbiomed.2016.12.006>

PMid:27940350 PMCID:PMC5462861

7. Di MS, Reed TT, Venditti P, Victor VM. Role of ROS and RNS Sources in Physiological and Pathological Conditions. *Oxid Med Cell Longev*. 2016; 2016. <https://doi.org/10.1155/2016/1245049> PMid:27478531 PMCID:PMC4960346

8. Szasz T, Bomfim GF, Webb RC. The influence of perivascular adipose tissue on vascular homeostasis. *Vasc Health Risk Manag*. 2013; 9:105-16. <https://doi.org/10.2147/VHRM.S33760> PMid:23576873 PMCID:PMC3616689

9. Wihastuti TA, Sargowo D, Heriansyah T, Aziza YE, Puspitarini D, Iwana AN, et al. The reduction of aorta histopathological images through inhibition of reactive oxygen species formation in hypercholesterolemia *rattus norvegicus* treated with polysaccharide peptide of *Ganoderma lucidum*. *Iran J Basic Med Sci*. 2015; 18(5):514-9.

10. Aldosari S, Awad M, Harrington EO, Sellke FW, Abid MR. Subcellular reactive oxygen species (ROS) in cardiovascular pathophysiology. *Antioxidants*. 2018; 7(1):14. <https://doi.org/10.3390/antiox7010014> PMid:29337890 PMCID:PMC5789324

11. Adiputro D, Khotimah H, Widodo M, Romdoni R, Sargowo D. Cathecins in ethanolic extracts of *Garcinia mangostana* fruit pericarp and anti-inflammatory effect in atherosclerotic rats. *J Exp Integr Med*. 2013; 2(1):61-4. <https://doi.org/10.5455/oams.170313.br.007>

12. Suthammarak W, Numpraphrut P, Charoensakdi R, Neungton N, Tunrungruangtavee V, Jaisupa N, et al. Antioxidant-Enhancing Property of the Polar Fraction of Mangosteen Pericarp Extract and Evaluation of Its Safety in Humans. *Oxid Med Cell Longev*. 2016; 2016:10. <https://doi.org/10.1155/2016/1293036> PMid:27703599 PMCID:PMC5040814

13. Francik R, Szkaradek N, Øelaszczyk D and Marona H. Antioxidant activity of Xanthone Derivatives. *Acta Poloniae Pharmaceutica-Drug Research*. 2016; 73(6):1505-1509.

14. Pedraza-Chaverri J, Cárdenas-Rodríguez N, Orozco-Ibarra M, Pérez-Rojas JM. Medicinal properties of mangosteen (*Garcinia mangostana*). *Food Chem Toxicol*. 2008; 46(10):3227-39. <https://doi.org/10.1016/j.fct.2008.07.024> PMid:18725264

15. Mahabusarakam W, Proudfoot J, Taylor W, Croft K. Inhibition of Lipoprotein Oxidation by Prenylated Xanthenes Derived from Mangostin. *Free Radic Res*. 2000; 33:643-659. <https://doi.org/10.1080/10715760000301161> PMid:11200095

16. Suttirak W, Manurakchinakorn S. In Vitro Antioxidant Properties of Mangosteen Peel Extract. *J Food Sci Technol*. 2014; 51(21):3546-3558. <https://doi.org/10.1007/s13197-012-0887-5> PMid:25477623 PMCID:PMC4252444

17. Wihastuti TA, Sargowo D, Soeharto S, Heriansyah T, Widyasih GR. Vasa Vasorum Angiogenesis through Increased Levels of H₂O₂, HIF-1 α , NF- κ B and iNOS: In Vivo Study of Atherosclerosis. *J Med Bioeng*. 2015; 4(5):342-9. <https://doi.org/10.12720/jomb.4.5.342-349>

18. Vikas K, Prakash CB, Gaurav K, Mohd R, Al-Abbasi FA, Jalaluddin AJ, Khan, Firoz A, Amita V. α -Mangostin Mediated

Pharmacological Modulation of Hepatic Carbohydrate Metabolism in Diabetic Induced Wistar Rat. *J Basic and Apply Science*. 2016; 5:255-276. <https://doi.org/10.1016/j.bjbas.2016.07.001>

19. Bryk D, Malecki M, Hajdukiewicz K, Sitkiewicz D. Trans Fatty Acids Induce a Proinflammatory Response in Endothelial Cells through ROS-Dependent Nuclear Factor- κ B Activation. *J Physiol Pharmacol*. 2011; (8):229-38.

20. Wihastuti TA, Heriansyah T, Soraya M, Wijayanti MD, Firani NK, Iskandar A, Anita KW, Agustin D, Pujiana L, Sandika N, Ardyanto E, Puji A, Wiji Y, Febrianto A, Nuraminudin A, Qurino F, Rachmawati S, Candra Y. Inhibition of Oxidative Stress in Hypercholesterolemic Rats by Soy Milk. *J Cardiovasc Disease Res*. 2016; 7(2):74-82. <https://doi.org/10.5530/icdr.2016.2.4>

21. Watanabe T, Yasunari K, Nakamura M, Maeda K. Carotid artery intima-media thickness and reactive oxygen species formation by monocytes in hypertensive patients. *Journal of Human Hypertens*. 2006; 20(5):336-40. <https://doi.org/10.1038/sj.jhh.1001990> PMid:16467862

22. Joan MCM, Michelle EM, Hiam AV. Vascular Cell Adhesion Molecule-1 Expression and Signaling During Disease: Regulation by Reactive Oxygen Species and Antioxidant. *J Antioxid Signal*. 2011; 15(6):1607-1638. <https://doi.org/10.1089/ars.2010.3522> PMid:21050132 PMCID:PMC3151426

23. Huang MZ, Lu XR, Yang YJ, Liu XW, Qin Z, Li JY. Cellular Metabolomics Reveal the Mechanism Underlying the Anti-Atherosclerotic Effects of Aspirin Eugenol Ester on Vascular Endothelial Dysfunction. *Int J Mol Sci*. 2019; 20(13):3165. <https://doi.org/10.3390/ijms20133165> PMid:31261711 PMCID:PMC6651823

24. Wihastuti TA, Sargowo D, Tjokroprawiro A, Permatasari N, Widodo MA, Soeharto S. Vasa Vasorum Anti-angiogenesis Through H₂O₂, HIF-1 α , NF- κ B, and iNOS Inhibition by Mangosteen Pericarp Ethanolic Extract (*Garcinia mangostana* Linn) in Hypercholesterol Diet-Given *Rattus norvegicus* Wistar Strain. *Vascular Health and Risk Management*. 2014; (10):523-531. <https://doi.org/10.2147/VHRM.S61736> PMid:25187725 PMCID:PMC4149390

25. Fernández-Alfonso MS, Somoza B, Tsvetkov D, Kuczmanski AD, Dashwood MR, Gil-Ortega M. Role of Perivascular Adipose Tissue in Health and Disease. *Comprehensive Physiology*. 2017; 8(1):23-59. <https://doi.org/10.1002/cphy.c170004> PMid:29357124

26. Ketonen J, Shi J, Martonen E, Mervaala E. Periadventitial adipose tissue promotes endothelial dysfunction via oxidative stress in diet-induced obese C57Bl/6 mice. *Circ J*. 2010; 74(7):1479-87. <https://doi.org/10.1253/circj.CJ-09-0661> PMid:20526041

27. Badimon L, Padró T, Vilahur G. Atherosclerosis, platelets and thrombosis in acute ischaemic heart disease. *European Heart Journal Acute Cardiovascular Care*. 2012; 1(1):60-74. <https://doi.org/10.1177/2048872612441582> PMid:24062891 PMCID:PMC3760546